1 **The fate of the Deep Western Boundary Current in the South Atlantic**

2 Silvia L. Garzoli ^{1,2}, Shenfu Dong ^{1,2}, Rana Fine³, Christopher S. Meinen², Renellys C.
3 Perez ^{1,2}, Claudia Schmid², Erik van Sebille^{4,5}, and Qi Yao ^{1,2}

4
5 ¹ Cooperative Institute for Marine and Atmospheric Studies, University of Miami

6 ² NOAA/Atlantic Oceanographic and Meteorological Laboratory

7 ³ Rosenstiel School of Marine and Atmospheric Sciences, University of Miami

8 ⁴ ARC Centre of Excellence for Climate System Science & Climate Change Research Centre,
9 UNSW, Sydney, New South Wales, Australia

10 ⁵ Now at Grantham Institute & Department of Physics, Imperial College London, London, United
11 Kingdom

13 **Abstract**

14 The pathways of recently ventilated North Atlantic Deep Water (NADW) are part of the
15 lower limb of the Atlantic Meridional Overturning Circulation (AMOC). In the South
16 Atlantic these pathways have been the subject of discussion for years, mostly due to the
17 lack of observations. Knowledge of the pathways of the AMOC in the South Atlantic is a
18 first order prerequisite for understanding the fluxes of climatically important properties.
19 In this paper, historical and new observations, including hydrographic and oxygen
20 sections, Argo data, and chlorofluorocarbons (CFCs), are examined together with two
21 different analyses of the Ocean general circulation model For the Earth Simulator (OFES)
22 to trace the pathway of the recently ventilated NADW through the South Atlantic.
23 CLIVAR-era CFCs, oxygen and salinity clearly show that the strongest NADW pathway
24 in the South Atlantic is along the western boundary (similar to the North Atlantic). In
25 addition to the western boundary pathway, tracers show an eastward spreading of NADW

between $\sim 17\text{--}25^\circ\text{S}$. Analyzed together with the results of earlier studies, the observations and model output presented here indicate that after crossing the equator, the Deep Western Boundary Current (DWBC) transports water with the characteristics of NADW and a total volume transport of approximately 14 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$). It crosses 5°S as a narrow western boundary current and becomes dominated by eddies further south. When this very energetic eddying flow reaches the Vitória-Trindade Ridge ($\sim 20^\circ\text{S}$), the flow follows two different pathways. The main portion of the NADW flow continues along the continental shelf of South America in the form of a strong reformed DWBC, while a smaller portion, about 22% of the initial transport, flows towards the interior of the basin.

1. Introduction

The role of the southward flowing Deep Western Boundary Current (DWBC) as a primary pathway for the cold, lower limb of the Atlantic Meridional Overturning Circulation (AMOC) has been well documented in the North Atlantic Ocean (e.g. Molinari et al., 1998; Schott et al., 2004; Johns et al., 2008; Meinen et al., 2012)¹. However the pathways and variability of the DWBC as the lower limb of the overturning

¹ Note: One exception to this is near and south of the Grand Banks of Newfoundland, where Lagrangian floats suggest that the DWBC flow breaks up and interacts significantly with the interior (e.g. Bower et al., 2009).

43 circulation in the South Atlantic Ocean are less well established. Much of what is known
44 about the deep flows in the South Atlantic has been inferred from the merging of
45 hydrographic sections in the region. They are used to trace the North Atlantic Deep
46 Water (NADW), the main water mass carried by the DWBC. For example, Reid's (1989)
47 maps of geopotential anomaly of the 3000 db surface with respect to the 3500 db surface
48 indicate that part of the NADW from the DWBC flows to the interior near 25°S
49 producing eastward tongues of high salinity and oxygen extending close to the Mid-
50 Atlantic Ridge. Reid's analysis of salinity, oxygen, phosphate, and silica also suggests
51 eastward flow towards the interior.

52 There are few direct observations of the DWBC in the South Atlantic. A two year-long
53 velocity and temperature time series from a moored array, shipboard observations, and
54 model simulations previously suggested that the southward flowing DWBC breaks up
55 into eddies before reaching 8°S (Dengler et al., 2004). They reported that between 8°S
56 and 16°S, the transport of NADW into the South Atlantic Ocean is accomplished by
57 southeastward migrating eddies, rather than by a continuous flow. Direct current meter
58 observations by Schott et al. (2005) supported the idea that a mean DWBC exists at 5°S
59 and dissolves into a sequence of deep eddies between 5°S and 11°S. Hogg and Owens
60 (1999) analyzed observations collected from Lagrangian floats launched as part of the

61 Deep Basin Experiment. They concluded that the line of sea mounts that extends
62 eastward from the South American coast just south of 20°S rising to depths of 2000 m or
63 less at places, (known as the Vitória-Trindade Ridge), interrupts the southward-flowing
64 DWBC and is the location for significant eddy exchange into the interior. Their float
65 trajectories provided little indication of a continuation of the DWBC flowing southward
66 beyond the Vitória-Trindade Ridge, although a single float trajectory did return to the
67 western boundary south of 20°S.

68 Further south, direct current measurements (Muller et al., 1998) and geostrophic current
69 estimates (Zangenberg and Siedler, 1998) have hinted at the existence of a DWBC along
70 the continental slope between 20°S and 28°S, with a southward transport that ranges
71 between 2 and 10 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3\text{s}^{-1}$). This might suggest the presence of at least a
72 portion of the DWBC along the western boundary south of the Vitória-Trindade Ridge. A
73 more recent study (Meinen et al., 2012) used data from a line of four pressure equipped
74 inverted echo sounders (PIES) as well as hydrographic and dissolved oxygen (O_2)
75 measurements to study the DWBC along 34.5°S on the South American continental
76 shelf/slope. These data have confirmed the presence of a DWBC with water mass
77 characteristics of NADW (elevated oxygen and salinity) at 34.5°S.

78 On the other side of the Atlantic basin at 30°S, hydrographic sections have also been used

79 to show southward flow of water with characteristics of NADW along the eastern
80 boundary and a mean transport of 3 Sv (Arhan et al., 2003). These observations are
81 consistent with a portion of the NADW perhaps having followed an interior pathway over
82 to the eastern boundary. Because of the paucity of observations to date, the details of this
83 interior pathway are still quite uncertain.

84 Enhancements to the suite of measurements made during hydrographic cruises over the
85 past decades allow for these hydrographic data to yield more information on ocean
86 pathways than were previously possible. For example, hydrographic sections of salinity
87 and dissolved oxygen from the WOCE Atlas Volume 3, Atlantic Ocean ([http://www-](http://www-pord.ucsd.edu/whp_atlas/atlantic/sections.htm)
88 [pord.ucsd.edu/whp_atlas/atlantic/sections.htm](http://www-pord.ucsd.edu/whp_atlas/atlantic/sections.htm)) show clear evidence of water with the
89 NADW characteristics (i.e., low temperature, $T \sim 3^{\circ}\text{C}$, and high salinity, $S > 34.8$ psu,
90 water with high levels of dissolved oxygen, $\text{O}_2 > 240 \mu\text{mol kg}^{-1}$) along the western
91 boundary at 11°S , 19°S , and 30°S (Koltermann et al., 2011). The NADW can also be
92 traced by looking for waters with high concentrations of chlorofluorocarbons (CFCs)
93 (Fine et al., 2002); recently ventilated NADW carries an elevated CFC concentration
94 because it has been more recently in contact with the atmosphere than the older waters at
95 the same depths that were last in contact with the atmosphere before the measurable CFC
96 transient ($\sim 1960\text{s}$). The high CFC NADWs flow equatorward from the high latitude

97 North Atlantic in the DWBC (e.g., Fine and Molinari, 1988; Smethie et al., 2000) and
98 along the equator (Weiss et al., 1985). Wanninkhof et al. (2010) analyzed repeat
99 observations from the meridional Atlantic section A16 along $\sim 25^\circ\text{W}$ from Iceland to
100 56°S . The composite quasi-meridional sections from the 2003 and 2005 A16 cruises for
101 salinity, apparent oxygen utilization, total dissolved inorganic carbon, nitrate, and silicate
102 show that water masses with NADW properties are present at 20°S in the center of the
103 basin ($\sim 25^\circ\text{W}$).

104 Numerical models have also been used to improve our understanding of the spreading of
105 the NADW into the South Atlantic. Speich et al. (2007) computed trajectories of the
106 particles that represent the southward flow of NADW at 44°N in the North Atlantic (i.e.,
107 characterized by a potential density anomaly greater than 27.7 kg m^{-3}). Trajectories were
108 computed in a coarse-resolution ocean general circulation model (2° uniform zonal
109 resolution, and meridional resolution that varies from 0.5° at the equator to $2^\circ \cos(\phi)$
110 poleward of the tropics). This coarse resolution Lagrangian simulation shows NADW
111 flowing southward along the coast of South America in the DWBC to approximately
112 45°S , after which it flows into the interior towards the east. Although analysis of a global
113 eddy-permitting (nominal horizontal resolution of 0.25°) Parallel Ocean Climate Model
114 (POCM) by Matano and Beier (2003) similarly demonstrated the main pathway for the

115 DWBC is along the coast of South America, there were also some indications of another
116 pathway flowing into the interior along approximately 27°S.

117 More recently, van Sebille et al. (2012) utilized the output from the global high-
118 resolution (0.1° horizontal resolution) Ocean general circulation model For the Earth
119 Simulator (OFES) model to study the NADW pathways in the South Atlantic. They
120 found 3.6 Sv of NADW crossing eastward at about 25°S underneath the trajectory
121 followed by the Agulhas rings that detach from the Agulhas Retroflection and then
122 slowly drift northwestward along the so-called Agulhas ring corridor. They analyzed the
123 velocity and density fields of the last 27-years of output from a 57-year (1950-2006) run
124 of the OFES model and found that the decay of the Agulhas rings, and their forward tilt
125 with depth, results in a southward velocity of 1 to 2 cm/s across isolines of planetary
126 vorticity in the deep waters. The associated stream function pattern yields a deep
127 circulation transporting about 4 Sv of NADW from the DWBC eastward at 25°S to the
128 southern tip of Africa. The pathway of the DWBC and how it reforms along the western
129 boundary south of 25°S were not the objective of the analysis conducted by van Sebille et
130 al. (2012), and will be analyzed in detail in what follows using the OFES model and
131 available observations.

132 Previous modeling and observational work has shown that important water mass

transformations occur in the South Atlantic (e.g., Garzoli and Matano, 2011, and references therein). For example, transformations occur in the very energetic regions near the boundaries such as the Brazil-Malvinas Confluence and the Cape Basin (Provost et al., 1999; Lumpkin and Speer, 2007; Jullion et al., 2010; Garzoli and Matano, 2011; Rimaud et al., 2012). These water mass transformation regions will both influence and be influenced by the pathways of the DWBC as it flows through the South Atlantic, and the transformations will have important impacts on the AMOC. The stability of the AMOC has been shown to be directly related to meridional salt fluxes between the South Atlantic and other ocean basins (e.g., Drijfhout et al., 2011; Bryden et al., 2011; Garzoli et al., 2013). Developing a detailed knowledge and understanding of the pathways of the lower limb of the AMOC will therefore be crucial for quantifying the fluxes of heat, fresh water, and CO₂ through the South Atlantic and for assessing the stability of the AMOC.

The objective of this paper is to utilize existing and recent hydrographic and tracer data sets, a new gridded subsurface Argo data product (Schmid, 2014), and Eulerian and Lagrangian analyses from the high-resolution OFES model, to extend the previous work conducted by van Sebille et al. (2012) and to learn more about the western boundary and interior pathways of the NADW in the South Atlantic. Furthermore the transports associated with these pathways will be quantified and will demonstrate the strength of the

flow along the western boundary pathway, which has historically been underestimated.

2. Observational Evidence

2.1 Pathways based on hydrographic and CFC tracers

The recently ventilated NADW carried by the lower limb of the AMOC is characterized by low temperature ($\sim 3^{\circ}\text{C}$), high salinity ($S > 34.8$ psu), and high levels of dissolved oxygen ($\text{O}_2 > 240 \mu\text{mol kg}^{-1}$). This NADW flows southward at neutral density levels between 27.9 and 28.1 kg m^{-3} in the South Atlantic near 35°S (Schmitz and McCartney, 1992; Meinen et al., 2012; Preu et al., 2013). Inspection of the CTD sections collected during the WOCE experiment at 11°S , 19°S and 30°S (Koltermann et al., 2011) show that water masses with these characteristics in the South Atlantic are concentrated in the western basin between 1500 and 3000 m depth. At 11°S these water are observed west of 20°W , and at 19°S and 30°S the recently ventilated NADW is found as far offshore as 5°W .

As mentioned in the introduction recently ventilated NADW can be tracked using other CFCs. Using the WOCE data era (1992-1997), maps of chlorofluorocarbon-11 (CFC-11) concentrations are generated via the data-interpolating variational analysis (DIVA)

169 gridding technique (Troupin et al. 2012) applied to the South Atlantic (Figure 1) on
170 neutral density level 27.9 kg m^{-3} , the level of the upper NADW between 1700 and 2000
171 m. High concentrations ($> 0.03 \text{ pmol kg}^{-1}$) of CFC-11 are observed along the western
172 boundary in the North Atlantic, along the equator, and extending into the South Atlantic
173 along the western boundary north of 5°N (Figure 1). On the WOCE era map,
174 concentrations above $0.05 \text{ pmol kg}^{-1}$ are found extending into the South Atlantic along
175 the western boundary only to $\sim 15^{\circ}\text{S}$. However, there is evidence of elevated
176 concentrations ($> 0.04 \text{ pmol kg}^{-1}$) of recently ventilated NADW in the interior near 23°S ,
177 35°W . Ventilated NADW is defined here as those waters of NADW origin that carry
178 above blank level CFC-11 (i.e. CFC-11 values $> 0.005 \text{ pmol kg}^{-1}$). CFCs are a transient
179 tracer, and they increased in NADW well after the mid-1990s (e.g., Kieke et al., 2007).
180 As a consequence, the southward and eastward spreading of ventilated NADW is
181 probably limited in the WOCE era data, as they had not yet had time to spread.

182

183 To investigate this point further, CLIVAR transect data (repeats of WOCE-era transects
184 obtained during the first two decades of the 2000s) have been analyzed in this study. CFC
185 data in the South Atlantic south of 10°S and north of 40°S were collected along the
186 following CLIVAR sections: A13 $\sim 5^{\circ}\text{--}10^{\circ}\text{E}$ in 2010, A15 20°W in 2000, A16S $\sim 25^{\circ}\text{W}$ in
187 2005 and 2013-2014, and A10 $\sim 30^{\circ}\text{S}$ in 2003 and 2011 (see brown lines in Figure 1).

188 Although other sections were occupied in the South Atlantic, they did not collect CFC
189 measurements. Thus, there are not enough CLIVAR data with which to make a map of
190 CFC concentrations, similar to what was done for WOCE data of the 1990s (Figure 1).
191 Instead, data from A10 (~30°S), A16S (~25°W) and A13 (~5°-10°E) are presented as
192 individual cross-sections of high CFCs, oxygen and salinity, which are used to show for
193 the first time the eastward spreading of NADW into the central South Atlantic Ocean
194 (Figure 2).

195

196 Between ~17°-25°S the NADW has reached as far east as 25°W during the 2005 (not
197 shown) and 2013-2014 (Figure 2a-c) occupations and there is a suggestion of elevated
198 CFCs at 5°-10°E south of 28°S during the 2010 occupation (Figure 2d). This can be more
199 clearly seen along 30°S (Figure 2e), as a large area with elevated CFCs $> 0.01 \text{ pmol kg}^{-1}$
200 is observed to extend east of Walvis Ridge between 5° and 15°E. Note that the high tracer
201 cores (CFC-11, salinity and oxygen) are not completely overlapping in space. If the high
202 CFC-11 core in Figure 2a is taken as $0.04 \text{ pmol kg}^{-1}$, then indeed it does occupy the same
203 depths as the oxygen and salt cores of Figures 2b and 2c. Furthermore, the high salinity
204 core reaches upward to almost 1500 m as does the CFC-11, but not oxygen. In addition,
205 highest concentrations of CFC-11 are observed at shallower depths 1500-2400 m in
206 Figure 2a, as compared with Figure 2e where the highest CFC-11 is observed at 2000-
207 2400 m. The upper part of the high CFC-11 core in Figure 2a at approximately 22°S,
208 25°W is of Labrador Sea origin. A probable explanation for the difference in signatures at

the two latitudes is that at 30°S near the western boundary the shallower Labrador Sea Water has been largely replaced with waters of southern origin and lower CFC-11, for example, Upper Circumpolar Deep Water.

Figure 2e shows the highest subsurface CFC concentrations ($> 0.2 \text{ pmol kg}^{-1}$) in NADW along the western boundary at 30°S. High CFCs spreading southward in the WBC is consistent with recent dissolved oxygen measurements collected during hydrographic cruises along the western boundary near 35°S (Schmitz and McCartney, 1992; Meinen et al., 2012; Preu et al., 2013). In time, as the CFC transient penetrates the South Atlantic, on future occupations of the $\sim 25^\circ\text{W}$, $\sim 5^\circ\text{--}10^\circ\text{E}$, and 30°S sections, it may be possible to observe interior eastward flowing NADW that has turned southward near the eastern boundary east of Walvis Ridge. To summarize, the CLIVAR-era CFCs, oxygen and salinity demonstrate that the strongest NADW signal in the South Atlantic (similar to the North Atlantic) is along the western boundary and that there is eastward spreading of NADW between $\sim 17^\circ\text{--}25^\circ\text{S}$, and NADW is observed east of the Walvis Ridge at 30°S.

2.2 Argo data

The Argo array is a global set of more than 3,000 free-drifting floats that sample the upper ocean to a maximum pressure level of 2000 dbar, which is within the upper layer of the NADW. Schmid (2014) developed a procedure to obtain three-dimensional fields

229 of absolute geostrophic velocities from Argo and altimeter data. Results were published
230 for the period ending in June 2012. In an expansion of Schmid (2014), 66,264 profiles of
231 temperature and salinity collected in the South Atlantic in the years 2000 to 2013, were
232 used to generate an updated product for this study. The 3-D velocity field extends from
233 the surface to 2000 dbar with a vertical resolution of 10 dbar. Velocity fields are derived
234 by first combining *in situ* dynamic height profiles from floats with the sea surface height
235 from AVISO (CLS, 1996) to generate synthetic dynamic height profiles with better
236 spatial and temporal coverage than could be achieved using only the *in situ* observations.
237 This process yields 162 monthly mean dynamic heights covering the period March 2000
238 to August 2013. Dynamic heights are used to calculate geostrophic velocities relative to
239 the reference level of 1000 dbar on a 0.5° longitude by 0.5° latitude grid from 5°S to
240 50°S. Absolute geostrophic velocity is then estimated using drift velocities obtained from
241 the trajectories of 825 Argo and WOCE floats, which were active from December 9, 1992
242 to February 20, 2014 and drifted in the pressure range of 800 to 1100 dbar. These drift
243 velocities, and the associated pressures, are used to derive gridded fields of the absolute
244 velocity (Schmid, 2014). The gridded monthly mean profiles of geostrophic relative
245 velocity are then referenced to the monthly mean float-derived absolute velocities, when
246 available. If no monthly mean absolute velocity estimate is available at a given grid point,
247 then the climatological absolute velocity is used. The resulting monthly mean absolute

geostrophic velocity fields are then averaged to generate a 3-D field of the long-term mean of the horizontal velocity.

The long-term mean gridded Argo-derived absolute velocity field at 2000 dbar (within the upper layer of the NADW) is shown in Figure 3 (top panel). This high-resolution velocity field is a bit overwhelming, so for illustration purposes polygons were defined around each of the two bands of deep flow discussed herein and the vectors within those polygons were colored to highlight the predominant flow of interest for the present discussion. In the boundary current polygon, the grid cells where the meridional velocity is negative (southward) were highlighted in red. In the polygon for the interior pathway, grid cells where the zonal velocity is positive (eastward) were highlighted in blue. In order to provide a measure of the significance of the velocity field, a t-test is applied to detect which zonal velocity component is too small to be significant with respect to a 95% confidence interval. Vectors where the zonal component of the velocity fails this criteria are excluded from the top panel of Figure 3. A similar t-test has also been applied to the meridional velocity component to ensure that the southward flows along the western boundary are significant. This was found to be the case with the exception of a very small band about 37°S, where the influence of the Brazil Malvinas confluence is large and the flow near the boundary has a relatively large (and significant) zonal

267 component.

268

269 South of the Vitória-Trindade Ridge (Figure 3, top panel), two branches of the flow are
270 evident. The dominant branch is a strong southward flow (highlighted in red) along the
271 coast of South America. This flow separates from the boundary at 38°S, the region of the
272 Brazil-Malvinas Confluence, and continues southward all the way to 45°S. The Argo
273 velocities also show eastward flow into the interior along roughly 22°S consistent with
274 the spreading of the NADW towards the center of the basin; this signal is not as clear as
275 the southward flow, however it is strongly supported by the CFC data (Figure 2). Some
276 details of the meridional and vertical structure of the eastward flow along this pathway
277 (highlighted in blue) are given in Figure 3 (bottom panel) which shows the zonal velocity
278 in cm sec^{-1} obtained from the Argo product along 25°W from 10° to 30°S. The figure
279 shows the evidence of two eastward jets at this longitude between 18° and 25°S. A jet
280 with a higher eastward velocity (maximum of 3.4 cm sec^{-1} at 2000 m) is observed
281 between 18°S and 20°S while the second eastward jet to the south of it, between 21°S and
282 24°S, is observed to have a maximum eastward velocity of 1.7 cm sec^{-1} at 2000 m at
283 22.5°S. These two eastward jets are located in the region where the CLIVAR water
284 properties indicate the presence of NADW crossing 25°W (Figure 2). The analysis also
285 shows two westward flows at around 12° and 16°S, the southern one possibly
286 representing part of a recirculation that may result in enhanced mixing of waters on this
287 pathway. Also there is a very weak indication (at around 19°S, 8°W) of a northward
288 transport that may feed back a small portion of the water into these westward flows
289 (Figure 3 top panel). However this small pathway is not connected by significant vectors

and as such it is not clear how it fits in the overall circulation pattern. Nevertheless it appears that most of the significant flow continues towards the east. Eastward of the Walvis Ridge ($\sim 0^\circ\text{E}$), the Argo velocity product does not show a clear picture of a continuous jet (Figure 3, top panel). This may be likely due to the high mesoscale motion in and northwest of the Cape Basin.

3. The Ocean General Circulation Model for the Earth Simulator (OFES)

Even with the improved data sets currently available for this study (updated hydrographic data, Argo float drift velocities), the data that can be used to study NADW pathways in the South Atlantic is still fairly limited. As such, a high quality ocean model can provide valuable circulation information. OFES is a massively parallelized implementation of version 3 of the NOAA/GFDL Modular Ocean Model (MOM3) run by the Japan Agency for Marine-Earth Science and Technology (JAMSTEC). The model equations have been discretized on a Mercator B grid with a horizontal resolution of 0.1° and 54 vertical z-levels. The simulation used in this study was spun up for 50 years with a monthly climatology of atmospheric forcing derived from the National Center for Environmental Prediction–National Center for Atmospheric Research (NCEP–NCAR) reanalysis (Masumoto et al., 2004), and then forced with daily mean NCEP–NCAR reanalysis data from January 1950 to December 2006 (Sasaki et al., 2008). OFES fields are publically available for a 27-year time period from January 1980 to December 2006 as snapshots at

310 3-day intervals. OFES reproduces the large-scale sea surface height variability in the
311 South Atlantic well when compared to altimetric data (e.g., Dong et al., 2011; Perez et al.,
312 2011). It has recently been used in various studies aimed at characterizing the time-mean
313 and time-variability of components of the AMOC in the South Atlantic (e.g. Dong et al.,
314 2011; Perez et al., 2011, van Sebille et al., 2012).

315

316 The OFES model has been validated in different regions of the global ocean by different
317 authors (Masumoto, 2010, and references therein). In the context of NADW pathways in
318 the South Atlantic, van Sebille et al. (2012) validated the velocity fields at the western
319 boundary of the South Atlantic between 5°S and 11°S by comparing sections of 27-year
320 mean meridional velocity in OFES in the DWBC with data collected by moored current
321 meters (Schott et al., 2005) and, at 30°S with data from Lagrangian floats collected by
322 Hogg and Thurnherr (2005). They concluded that the extent, position, and strength of the
323 DWBC in OFES at these locations compare well with the available *in situ* observations at
324 that time.

325

326 In the present study, further validation is done by comparing fields from the OFES model
327 with hydrographic data collected on zonal sections along 5°S and 30°S, and the model
328 output is also compared to the gridded World Ocean Atlas 2009 temperature and salinity

329 climatology (WOA09; Locarnini et al., 2010), and the Argo data product discussed
330 earlier (Section 2.2). These data were used to validate the accuracy of the location of the
331 different water masses in the South Atlantic in the model and the total volume transport
332 of NADW that enters and exits the South Atlantic at various latitudes between 4.6°S and
333 35°S.

334

335 Comparisons of WOA09 and OFES salinity sections in the South Atlantic show that the
336 boundaries of the neutral density levels (27.9 and 28.1 kg m⁻³), which bracket the NADW,
337 are very similar in the model and in the observations. An example is Figure 4, which
338 shows zonal sections of the climatological mean salinity from the WOA09 (left), and
339 from the 27-year mean OFES model section (right), both along 30°S . The lower panel of
340 Figure 4 shows the corresponding potential temperature/salinity diagram. In the interior
341 of the basin, the upper boundary of the NADW (neutral density 27.9) is at 1700 m both in
342 the model and the observations. Near the continental boundaries, the upper boundary of
343 the NADW is slightly deeper in the observations (2000 m in the west and 1500 m in the
344 east for the observations, compared with 1800 m and 1500 m for the model) (Figure 4
345 upper panel). Differences in the depth of the lower boundary of NADW (neutral density
346 28.1 kg m⁻³) in the observations and the model are mostly related to differences in their
347 representations of bottom topography in particular at the Mid Atlantic Ridge (deeper

white contours in Figure 4, upper panels). In addition, while the high salinity water on the eastern side in the observations is captured well in the model, the agreement is not as good in the west. The area bounded by the $S = 34.8$ contour in the west is thicker in the observations than in the model. A potential temperature-salinity (T/S) comparison (Figure 4, lower panels) indicates that at 30°S the model is more saline (35.1) than the observations (34.8) for upper NADW with neutral density between 27.9 and 28 kg m^{-3} , but with overall similar T/S structure along the 28.1 kg m^{-3} neutral density contour.

3.1 Lagrangian analysis

As mentioned in the Introduction, van Sebille et al. (2012) examined the pathways of NADW in the South Atlantic in the OFES model using Lagrangian pathways. In their experiment, van Sebille et al. (2012) focused their analysis on the connectivity between the DWBC and the Cape Basin.

In this paper, we used the simulated float database produced by van Sebille et al. (2012) to further examine the primary NADW pathway in the DWBC along the South American coast as well as the interior pathway. A brief description of the procedure they used is as follows: Float deployments were simulated along a grid at 5°S whenever the local instantaneous transport within the grid cell is southward. The grid spanned the full width

366 of the South Atlantic between 1000 m and 3500 m depth and had 100 m vertical spacing.
367 Since most southward transport is carried by the DWBC, the horizontal grid spacing was
368 0.1° in the DWBC west of 34°W and 0.5° in the rest of the basin. For a total of 5 years,
369 one simulated float was released every 3 days. The vast majority of these floats are within
370 the NADW layer (van Sebille et al., 2012). Each float was assigned a transport equal to
371 the volume flux (velocity times area) through the grid cell in which it starts. After release,
372 the floats were tracked forward with a fourth order Runge-Kutta integration scheme as
373 implemented in the Connectivity Modeling System (CMS, Paris et al., 2013) using
374 snapshots of the OFES velocity fields with 3-day resolution. Advection of a float was
375 terminated when the float rises above 500 m depth, or when the float left the region
376 bounded on the north by 5°S , at the south by 45°S , and on the east by 17°E . At the end of
377 the 27-year run, the model output was looped to produce a longer time series (300 years)
378 for forward integration of the simulated float tracks.

379 In this section, these 100,000+ synthetic float tracks are further analyzed to show the
380 main pathway of the NADW along the continental slope of South America as well as the
381 interior pathway indicated by hydrographic, tracers, and Argo observations discussed in
382 Section 2. Transport estimates indicate that approximately 13.2 Sv (with a standard
383 deviation of 3.1 Sv) are carried southward of 5°S across the entire basin in the depth

384 range where the float were deployed (1000 to 3500 m). This result is in agreement with
385 the value obtained from the hydrographic data collected along the WOCE A07 line (12.6
386 Sv southward between neutral densities 27.9 to 28.1 kg m⁻³, see section 4). Lagrangian
387 particle backtracking (i.e. tracing particle pathways backward in time to determine where
388 the trajectories started) from points further south indicates that most of this south-
389 southeastward flow originates west of 30°W along 5°S (90% of the total virtual floats).

390

391 The synthetic float trajectories illustrate the very eddy-rich nature of the flows in the
392 South Atlantic. To focus on the primary pathways, the trajectories of the synthetic floats
393 that cross 5°S are then followed for two different pathways. In the first pathway, the
394 Lagrangian flow that crosses 30°S west of 30°W and continues south west of 40°W is
395 tracked and the transport is estimated. Examples of some of the floats that follow this
396 pathway are shown in Figure 5, top panel. The total transport carried by those
397 trajectories estimated following the procedure outlined in van Sebille et al. (2012) is
398 approximately 9.4 Sv (with a standard deviation of 4.2 Sv) or 71% of the total southward
399 transport across 5°S. A second pathway is defined by synthetic floats crossing 5°S west
400 of 30°W that continue towards the Cape Basin directly beneath the Agulhas Ring
401 corridor, crossing 17°E with the constraint of previously passing through the region
402 bounded by 20° and 30°S and by 10° and 9°W (examples shown in Figure 5, bottom

panel). This is the interior pathway observed by the hydrographic, tracers, and Argo observations. The transport associated with this pathway is approximately 2.9 Sv (with a standard deviation of 1.4 Sv). This is approximately 22% of the simulated float transport in the NADW. These estimates are in agreement with those previously obtained by van Sebille et al. (2012) who calculated that of the 15.0 Sv net southward transport crossing 5°S, almost all of which is in the NADW density range, 10.2 Sv leaves the South Atlantic across the southern boundary at 45°S. They also calculated that 3.6 Sv exits across the Agulhas section at 17°E with the remaining floats recirculating in the basin and staying in the domain for a very long time (more than 300 years).

The present analysis of the synthetic Lagrangian floats leads to the conclusion that the main pathway along which the NADW spreads into the South Atlantic is along the continental shelf of South America, in the southward DWBC. The second pathway crosses the Atlantic in agreement with that described by van Sebille et al. (2012) and with the observational findings presented in Section 2.

3.2 Eulerian analysis

In this present analysis, OFES fields are also analyzed using more traditional Eulerian methods that allow direct comparison with observations. The objective of the analysis is

422 to quantify the associated Eulerian transports and to look at the variability of the NADW
423 pathways at different levels in the model and compare and contrast the Eulerian results
424 with the Lagrangian results.

425

426 **3.2.1 Kinetic energy variability**

427 In order to provide a better picture of the energy distribution and the characteristic of the
428 flow, the mean velocity magnitude and eddy kinetic energy (EKE) in the model were
429 calculated at different depths within the NADW layer: 1700 m (the upper boundary of
430 NADW), 2500 m (the NADW core depth), and 3600 m (the lower boundary of NADW).
431 Examples are shown for the two upper depths in Figure 6. The top panels show the
432 annual mean (model years 2001 to 2006) of the velocity magnitude at 1700 m (left) and
433 2500 m (right). The two bottom panels show the mean EKE at 1700 m (left) and at 2500
434 m (right). Note, although not shown, the distribution of velocity and EKE at 3600 m
435 figures are similar to those at 2500 m.

436

437 The mean velocity magnitude (Figure 6, top panels) shows a steady and strong DWBC
438 flow following the continental shelf of South America at 1700 m depth from the equator
439 down to the Vitória-Trindade Ridge. At 2500 m, the mean DWBC flow is weaker but still
440 identifiable. This pattern is in agreement with the mean Argo horizontal velocity map

441 (Figure 3). Analysis of the EKE fields (lower panels) reveals that the very energetic flow
442 variability entering the South Atlantic from the tropics is fairly tightly confined along the
443 western boundary (e.g. near 5°S). Traveling southward, the region of flow variability
444 extends a little further offshore beginning around 8°S . This increase in the EKE is
445 consistent with the DWBC breaking up into rings at this latitude as previously suggested
446 by Dengler et al. (2004) and Schott et al. (2005). When this flow reaches the Vitória-
447 Trindade Ridge at approximately 20°S the region of high variability stretches even
448 further offshore, which is consistent with the flow becoming even more energetically
449 eddy-dominated due to the interaction with the coastline and topographic ridge extending
450 offshore.

451

452 There is a signature of enhanced EKE stretching eastward across the basin towards the
453 Walvis Ridge (Figure 6, bottom panels), suggesting that these eddies may play a role in
454 transporting NADW into the interior (consistent with the simulated Lagrangian pathways,
455 the Argo velocity map, and the water mass maps). This feature is most clearly seen in the
456 EKE maps at 1700 m, although there are hints at 2500 m as well. The EKE maps also
457 show a narrow tongue of higher variability close to the southern side of the Vitória-
458 Trindade Ridge and continental slope to the south, consistent with a more stable and
459 stationary DWBC flow reforming on the continental slope moving southward to about

460 25°S. This is confirmed by an increase in the mean horizontal velocity (top panels) south
461 of 25°S that remains confined along the western boundary until 40°S (again the signals
462 are stronger at 1700 m than 2500 m). High levels of energy are also observed at the
463 eastern side of the basin in the region of the retroflexion of the Agulhas Current and the
464 Agulhas Ring corridor (blue area in bottom panels of Figure 6 extending northwest from
465 the southern tip of Africa), and in the western side of the basin at the confluence of the
466 Brazil/Malvinas currents (blue area south of about 38°S near the western boundary). At
467 the core of the NADW, 2500 m (Figure 6 lower right panel), the EKE along the western
468 boundary is lower than at 1700 m, but the western boundary signals are still much higher
469 than is found in the interior. At the lower boundary of the NADW, 3600 m (not shown),
470 the fields are less energetic than at 2500 m (Figure 6), however, the distribution of EKE
471 is similar.

472

473 Animations of the flows in the South Atlantic at all three levels analyzed in the model
474 (not shown) indicate that energetic mesoscale features propagating westward are
475 prevalent in the South Atlantic basin at these depths. In particular, signals generated in
476 the Agulhas Current region propagate northwestward into the interior along the model
477 ring corridor and in some cases reach the western boundary. To look at the westward
478 component of the propagation of the mesoscale signals in the basin, speed and EKE were

479 plotted as a function of longitude for different latitudes and at different depths. An
480 example of this westward propagation can be seen in Figure 7 that shows a Hovmoller
481 diagram of the speed and the EKE along 25°S at 2500 m from 2001 to 2006 as a function
482 of longitude and time. The observed westward propagation has the characteristics of a
483 Rossby wave-like feature with estimated speeds of about 5 to 7 cm/s. The feature is
484 consistent with previous satellite and *in situ* observations of first baroclinic mode Rossby
485 waves (e.g., Osychny and Cornillon, 2004; Meinen and Garzoli, 2014).

486

487 **3.2.2. Volume transports**

488 During WOCE, a hydrographic section was occupied along 4.6°S (A07; Oudot, 1993;
489 Fajar et al., 2011). This latitude is close to the latitude of the simulated float deployments
490 in the experiment by van Seville et al. (2012) discussed in Section 3.1. Data from the
491 WOCE section is used to obtain additional Eulerian validation of the OFES model flow
492 crossing this latitude, particularly within NADW density levels. Temperature and salinity
493 data from the WOCE section are used in conjunction with Argo drift velocities, at
494 nominally 1000 m depth, to calculate the absolute volume transport of NADW between
495 neutral density levels. From the temperature and salinity profiles, the geostrophic
496 velocities are calculated with a reference level of 1000 m. Using the 3-D velocity
497 profiles derived from Argo in the South Atlantic (Section 2.2), the absolute velocity at

1000 m between each pair of hydrographic sections is added to the geostrophic relative velocities in order to obtain the absolute velocity. From the latter, the total integrated NADW transport across the basin-wide section is estimated to be -12.6 Sv (southward) between neutral density layers 27.9 and 28.1 kg m⁻³. The corresponding mean transport in the OFES model across 5°S between neutral densities 27.9 and 28.1 kg m⁻³ is -13.9 Sv (southward). The standard deviation (σ) of the mean values obtained from the 27-year model output is 0.95 Sv. This total integrated southward flow is mostly concentrated west of 30°W.

506

From the model velocity fields, meridional volume transports at different latitudes are obtained between neutral density levels 27.9 and 28.1 kg m⁻³ corresponding to the boundaries of the NADW. Results focusing on the meridional transport along the western and eastern boundaries are summarized in Figure 8 and Table I. West of 30°W, there is a southward flow of -13.9 Sv ($\sigma = 1.2$ Sv) at 10°S that is reduced to -11.8 Sv ($\sigma = 0.7$ Sv) at 20°S, the latitude of the Vitória-Trindade Ridge. Between 20°S and 30°S, the coastline makes an abrupt shift westward of roughly 5° of longitude, so south of 25°S different integration limits were used. The strong southward mean flow continues to be concentrated on the western boundary following the continental shelf of South America. From the -12 Sv flowing southward at 25°S, -9.0 Sv ($\sigma = 0.5$ Sv) can be found

517 concentrated west of 40°W (as shown in Figure 8) while a smaller -2.5 Sv is flowing
518 south more broadly between 30°W and 40°W (not shown in the figure for visual impact).
519 The southward transport at 35°S is -7.5 Sv ($\sigma = 0.3$ Sv) and there is little indication of
520 additional southward flow offshore of 40°W. The loss of transport between 30°S and
521 35°S seems to be primarily associated with a significant recirculation offshore of the
522 DWBC. Analysis of the model zonal velocity shows that the mean zonal transport is
523 westward -1.3 Sv at 30°S and increase to -3.0 Sv at 35°S which is an indication of the
524 recirculation consistent with the velocity field from the Argo floats shown in Figure 3.
525 Detailed analysis of the model (not shown) indicates that by 35°S, there is a northward
526 volume transport of 1.5 Sv between 45°W and 40°W, or roughly 20% of the strength of
527 the southward DWBC transport at that latitude.
528
529 Along the eastern boundary, east of 0°E, only a very small volume of water crosses 5°S (-
530 0.2 Sv with a standard deviation of 0.03 Sv) and the meridional volume transport near the
531 eastern boundary remains small with values of -0.4 ($\sigma = 0.01$ Sv) at 10°S and 0.2 Sv at
532 20°S ($\sigma = 0.03$ Sv). At 25°S, the contributions from the interior of the basin begin to
533 augment the southward volume transport, reaching a maximum of -4.6 Sv ($\sigma = 0.2$ Sv) at
534 35°S.

535

536 Figure 8 can also be used to revisit a concept discussed earlier, namely that the DWBC
537 reforms in a coherent manner as a continuous boundary current south of the Vitória-
538 Trindade ridge at 20°S. To illustrate this, the zero-lag correlations between the DWBC
539 meridional volume transport at 34.5°S integrated from coast to 44.5°W and 1700 m to
540 3000 m (over the 27-year record) and the meridional velocity at 2500 m depth were
541 determined at each OFES grid-point in the South Atlantic (color shading in Figure
542 8). Correlations are significant with 95% confidence, if they are larger than ± 0.11
543 (assuming there are 12 independent samples per year over 27 years, or 324 independent
544 samples indicated with black and grey contours). Positive correlations (gray shading)
545 along the coast between roughly 22°S and 42°S indicate that the meridional flow between
546 these latitudes is coherent with the DWBC at 34.5°S. This is consistent with the DWBC
547 having reformed into a steady flow in this region, as opposed to the eddy dominated flow
548 where the horizontal scales would be small enough that transport would be poorly
549 correlated beyond an eddy length scale. The negative correlations (cyan shading)
550 observed offshore of the strong positive correlation band are consistent with recirculation
551 of the DWBC waters in the model between roughly 37°S to 22°S, just south of the
552 Vitória-Trindade ridge. Note that the pattern of the zero-lag correlations is not
553 significantly different from that of the maximum lagged correlations (not shown).

554

555 In what follows, Eulerian volume transports obtained from the OFES model and available
556 observations across the basin and at the boundaries are analyzed at 5°S, 30°S, and 34.5°S
557 and compared with the Lagrangian transports estimated by van Sebille et al. (2012).
558 Estimates of the Eulerian volume transports are computed in two ways: integrated within
559 the neutral density range 27.9 and 28.1 kg m⁻³, the defined boundaries of NADW; and
560 integrated between 1000 and 3500 m, the depth span over which the synthetic Lagrangian
561 floats were launched by van Sebille et al. (2012). Results are shown in Table I; negative
562 numbers indicate southward flow. Column 1 (black numbers) are the Eulerian transports
563 estimated from the OFES model between density levels 27.9 to 28.1 kg m⁻³. In the same
564 column results are compared with estimates previously discussed obtained from WOCE
565 A07 CTD data coupled with Argo data collected along nominally 5°S, and with those
566 obtained by other observational systems in the area: from a CLIVAR A10 CTD section
567 collected along 30°S, and from the average of the values obtained from a series of
568 expendable bathythermograph (XBT) transects conducted nominally along 35°S and
569 analyzed jointly with hydrographic and Argo data (see Garzoli et al., 2013) (called
570 “combined observations” in the Table). Column 2 show the volume transports estimated
571 from 1000 to 3500 m in a Eulerian reference frame, and column 3 the Lagrangian
572 transports from simulated floats deployed between 1000 and 3500 m following van
573 Sebille et al. (2012). Overall, there is agreement between model results and observations.
574 Across 5°S basin-wide, there is very good agreement between the basin wide transports
575 determined between neutral densities 27.9 and 28.1 kg m⁻³ (-13.9 Sv, $\sigma = 0.96$ from the
576 model, -12.6 Sv from observations). The simulated Lagrangian estimate of the flow at

577 5°S in the 1000 to 3500 m layer (-13.2 Sv , $\sigma = 3.2 \text{ Sv}$) is similar to the Eulerian
578 simulated transport between neutral density levels (13.9 Sv , $\sigma = 0.96 \text{ Sv}$). However, the
579 Eulerian estimates from observations between 1000 and 3500 m (-21.4 Sv) are noticeably
580 larger than those obtained from the model (-16.8 Sv , $\sigma = 1.1 \text{ Sv}$); this could be an
581 indication of the model not capturing the circulation in the deep ocean (below 3000 m)
582 correctly.

583

584 When transports are estimated west of 40°W there is very good agreement between
585 estimates from the model and the observations. The difference between Eulerian and
586 Lagrangian results can be explained by analyzing the depth of the simulated float
587 trajectories, which suggests that even though the synthetic floats were launched between
588 1000 and 3500 m, the majority of the floats tended to occupy a more restricted depth
589 range from 1000 to less than 3000 m.

590

591 At the eastern boundary there is good agreement between the model and observations in
592 Eulerian coordinates (Table I), but the relationship is the opposite of those observed near
593 the western boundary (see also Figure 8). For example, southward volume transport
594 across 34.5°S and east of 0°E in the model is larger when estimated between density
595 layers (-4.6 Sv , $\sigma = 0.2 \text{ Sv}$) than between 1000 and 3500 m (-2.8 Sv , $\sigma = 0.23 \text{ Sv}$); a
596 similar, albeit smaller, difference is found in the observations (-4.2 Sv , $\sigma = 1.9 \text{ Sv}$ versus

597 -3.7 Sv, $\sigma = 1.0$ Sv). This is because at the eastern boundary the fixed 1000-3500 m depth
598 range includes some of the northward flow of Antarctic Intermediate Water (AAIW)
599 above the NADW, and the result is a smaller net southward flow in the fixed depth
600 integral (e.g., Schmid et al., 2003).

601

602 In addition to the comparisons shown in the Table, one additional data set that can be
603 compared is from the pressure-equipped inverted echo sounders (PIES) deployed along
604 34.5°S west of 44.5°W (Meinen et al., 2012). The PIES transports calculated between
605 density levels 27.9 and 28.1 kg m⁻³ (-10.9 Sv) is larger than the model result (-5.7 Sv, not
606 shown in the Table) when integrated to 44.5°W. About 2 Sv of this difference can be
607 explained by a northward flow observed in the model, between 47°W and 49°W that is
608 not present in the PIES observations. Furthermore the agreement improves when the
609 volume transports are calculated between 1000 and 3500 m (PIES: -16.7 Sv; model: -14.0
610 Sv).

611

612 **4. Discussion and conclusions**

613 Lagrangian and Eulerian analysis of the OFES model, validated against
614 hydrographic/tracer, XBT, Argo and PIES data, provides a description of the pathways of
615 the DWBC as it flows through the South Atlantic, carrying NADW southward as the

616 lower limb of the AMOC. From the Eulerian model analysis, it is estimated that
617 approximately -13.9 Sv of water crosses 5°S flowing southward into the South Atlantic in
618 a layer limited by the neutral density surfaces 27.9 and 28.1 kg m⁻³. These density levels
619 correspond approximately to depths between 1700 m and 3000 m, and are the boundaries
620 of the NADW ventilated with CFCs in the South Atlantic.

621

622 Re-analysis of the WOCE-era hydrographic data collected along 5°S in conjunction with
623 the Argo drift velocity field is in agreement with further analysis of the Lagrangian floats
624 advected for 300 years in OFES model following van Sebille et al. (2012). This analysis
625 indicates that 90% of the total floats that constitute the southward flow at 5°S originate
626 west of 30°W. These results are obtained by backtracking the particles, and by the
627 Eulerian estimates of volume transport integrated across 5°S.

628

629 Crossing 5°S and mostly concentrated west of 30°W, the DWBC carries NADW along
630 the coast of South America and, in the model in agreement with previous results, breaks
631 up into rings near 8°S, a region where the flow is mostly “non-steady.” Analyses of the
632 eddy kinetic energy (EKE) at different depth levels in the model indicate high levels of
633 energy spreading further offshore south of 8°S, consistent with the formation of energetic
634 rings. This “ring-dominated” flow continues southward to the Vitória-Trindade Ridge

635 (near 20°S). When the ring-dominated flow reaches the Vitória-Trindade Ridge, the
636 NADW is forced to flow to the east due to the topography. The branching of the flow in
637 this area is due to the preservation of potential vorticity (Zangenberg and Siedler, 1998).
638 Theory suggests that part of the along-slope southward flow turns into a zonal flow to the
639 east at the latitudes where the change in planetary vorticity is sufficiently large to
640 compensate for the change in potential vorticity due to the initial depth changes. In the
641 Southern Hemisphere, the Coriolis parameter is negative and the water column rotates
642 clockwise with increasing depth.

643

644 The model suggests that about 22% of the NADW flows eastward from the western
645 boundary into the interior South Atlantic. WOCE and CLIVAR-era CFC observations
646 give a clear pathway eastward roughly underlying the Agulhas Ring corridor, and
647 extending over a fairly broad latitude range (Figures 1 and 2). The Argo velocity field in
648 the upper NADW layer does not indicate a robust mean pathway for this eastward flow;
649 rather it is characterized by several meanders (Figure 3) and relatively weak mean
650 eastward jets. As there is a broad latitudinal distribution of properties extending eastward
651 into the interior and the complex Argo velocity mean picture is not particularly
652 suggestive of a narrow corridor of NADW transfer under the Agulhas Ring corridor, it is
653 possible that more than one mechanism is involved in the transfer of ventilated NADW

properties to the east near 20°S. Therefore, it is hypothesized that this southeastward spreading of properties could result from a combination of the mechanism of eddy-thickness flux divergence due to Agulhas ring decay, previously proposed and examined by van Sebille et al. (2012), enhanced by the clearly highly energetic eddy field at the Vitória-Trindade ridge and eastward jets observed in the velocity field.

The most prevalent motion noted in the animations of the ocean model velocity fields at three depths (not shown, but this can be inferred from the connectivity of the elevated EKE regions in the Cape Basin to Vitoria-Trindade Ridge in Figure 6, and Hovmoller diagrams in Figure 7) associated with the upper boundary, core, and lower boundary of NADW is a strong northwestward propagation consistent with the westward propagation characteristics of Rossby Waves. It is hypothesized that this westward propagation interacting with the mean circulation field may be the reason why south of the Vitória-Trindade Ridge most of the flow (~71% in the model) returns to the continental slope, as a more steady flow, forming the main branch of the DWBC south of the ridge. However, a detailed analysis of this complicated eddy-mean flow interaction is beyond the scope of the present study and is a topic for future research.

At 34.5°S, the DWBC meridional volume transport between the coast and 44.5°W and

673 from 1000 to 3500 m is about -14 Sv in the model. This model transport is close to an
674 estimate of -14.5 Sv from hydrographic data collected at 30°S. Further analysis of the
675 model trajectories (not shown) indicates that after crossing 34.5°S, a small amount (about
676 1 Sv) of NADW flows towards the east at the latitude where the Malvinas Current turns
677 offshore (~38°S), in agreement with previous results from van Sebille et al. (2012). The
678 rest of the flow continues south to approximately 45°S where it joins with the eastward
679 flowing Antarctic Circumpolar Current.

680

681 Eulerian volume transport estimates obtained from the velocity field of the OFES model
682 between 27.9 and 28.1 neutral density levels (Figure 8) indicate that about 2 Sv of
683 NADW does not leave the South Atlantic as NADW. This is consistent with previous
684 modeling and observational results (Garzoli and Matano, 2011), which found that in the
685 Brazil-Malvinas Confluence region approximately 2.6 Sv of water is transformed from
686 NADW to AAIW.

687

688 Based on the available observations and the model analysis, we conclude that the NADW
689 follows two different pathways south of 5°S. The main pathway (~71%) is southward in
690 the DWBC flowing along the coast of South America. A smaller portion (~22%) flows
691 eastward towards the interior of the basin. Better understanding of the pathways of the

692 lower limb of the AMOC will be crucial for quantifying the fluxes of heat, fresh water
693 and CO₂ through the South Atlantic. Furthermore, for understanding variations of the
694 AMOC, detailed knowledge of temporal changes of the pathways as well as what
695 mechanisms control those variations will be necessary.

696

697 **Acknowledgements**

698 This research was carried out in part under the auspices of the Cooperative Institute for
699 Marine and Atmospheric Studies (CIMAS), a Cooperative Institute of the University of
700 Miami and the National Oceanic and Atmospheric Administration (NOAA), cooperative
701 agreement #NA10OAR4320143. Additional support was provided by NOAA's Climate
702 Program Office under grant # NA13OAR4310131, by the NOAA Southwest Atlantic
703 MOC project, and the Physical Oceanography Division of NOAA's Atlantic
704 Oceanographic and Meteorological Laboratory. Rana Fine thanks the Physical
705 Oceanography Program of the National Science Foundation for support. Erik van Sebille
706 was supported by the Australian Research Council via grant DE130101336. The OFES
707 simulation was conducted on the Earth Simulator under the support of JAMSTEC. The
708 authors are indebted to Yoshikazu Sasai and Ricardo Matano for providing output from
709 the OFES simulations. The authors would like the anonymous reviewers whose
710 comments helped to the submission of an improved manuscript.

711

712 **References**

713 Arhan M., H. Mercier and Y-H Park, 2003: On the deep water circulation of the eastern
714 South Atlantic Ocean, *Deep-Sea Res. I*, **50**, 889–916, doi:10.1016/S0967-
715 0637(03)00072-4.

716 Bower, A. S, Lozier, M.S, Gary, S.F. and Böning, C.W, 2009: Interior pathways of the
717 North Atlantic meridional overturning circulation. *Nature*, vol. 459, no. 7244, pp
718 243-247.

719 Bryden, H. L., B. A. King, G. D. McCarthy, 2011: South Atlantic overturning circulation
720 at 24°S. *J. Mar. Res.* 69 (1), 38–55.

721 CLS (1996): AVISO User Handbook for Merged TOPEX/POSEIDON products. CLS

722 Ref.: AVI-NT-02-101. See www.aviso.altimetry.fr/en/data/product-

723 information/aviso-user-handbooks.html

724 Dengler, M., F. A. Schott, C. Eden, P. Brandt, J. Fischer, and R. J. Zantopp, 2004: Break-
725 up of the Atlantic deep western boundary current into eddies at 8°S, *Nature*, 432,
726 1018–1020.

727 Dong, S., M. Baringer, G. Goni, and S. Garzoli, 2011: Importance of the assimilation of
728 Argo Float Measurements on the Meridional Overturning Circulation in the South
729 Atlantic. *Geophys. Res. Lett.*, 38, L18603, doi:10.1029/2011GL048982.

730 Drijfhout, S. S., S. L. Weber, E. van der Swaluw, 2011: The stability of the MOC as
 731 diagnosed from model projections for pre-industrial, present and future climates.
 732 Clim. Dyn. 37, 1575–1586, [doi:10.1007/s00382-010-0930-z](https://doi.org/10.1007/s00382-010-0930-z).
 733 Fajar, N. M., F. F. Perez, A. Velo, and A. F. Rios, 2011: Data recovery of A06 and A07
 734 WOCE cruises. Earth Syst. Sci. Data Discuss., 4, 99–122, [doi:10.5194/essdd-4-99](https://doi.org/10.5194/essdd-4-99).
 735 Fine, R.A. and R. Molinari, 1988: A continuous Deep Western Boundary Current be-
 736 tween Abaco (26.5°N) and Barbados (13°N). A Rapid Response paper in Deep-Sea
 737 Research, 35: 1441-1450.
 738 Fine, R.A., M. Rhein and C. Andri, 2002: Using a CFC effective age to estimate
 739 propagation and storage of climate anomalies in the Deep Western North Atlantic
 740 Ocean. Geophys. Res. Letts., 29(24), 2227, [doi:10.1029/2002GL015618](https://doi.org/10.1029/2002GL015618).
 741 Garzoli, S. L. and R. Matano, 2011: The South Atlantic and the Atlantic Meridional
 742 Overturning Circulation. Deep-Sea Res., Part II, 58(17-18):1837-1847,
 743 [doi:10.1016/j.dsr2.2010.10.063](https://doi.org/10.1016/j.dsr2.2010.10.063).
 744 Garzoli, S. L., M. O. Baringer, S. Dong, R. Perez, and Q. Yao, 2013: South Atlantic
 745 meridional fluxes. Deep-Sea Res., Part I, 71:21-32, [doi:10.1016/j.dsr.2012.09.003](https://doi.org/10.1016/j.dsr.2012.09.003).
 746 Hogg, N. G. and W. B. Owens, 1999: Direct measurement of the deep circulation within
 747 the Brazil Basin. Deep-Sea Research II, 46, 335-353, [doi:10.1029/2004JC002311](https://doi.org/10.1029/2004JC002311).
 748 Hogg, N. G. and A. M. Thurnherr, 2005: A zonal pathway for NADW in the South

749 Atlantic. Journal of Oceanography, 61(3): 493-507.

750 Johns, W. E., L. M. Beal, M. O. Baringer, J. R. Molina, S. A. Cunningham, T. Kanzow,
751 and D. Rayner, 2008: Variability of shallow and deep western boundary currents off
752 the Bahamas during 2004-05: Results from the 26°N RAPID-MOC array.
753 J.Phys.Oceanogr., **38**, 605– 623.

754 Jullion, L., K. J. Heywood, A. C. Naveira Garabato, and D. P. Stevens, 2010: Circulation
755 and water mass modification in the Brazil-Malvinas confluence. J. Phys. Oceanogr.,
756 **40**, 845-864, doi: 10.1175/2009JPO4174.1.

757 Kieke, D., M. Rhein, L. Stramma, W. M. Smethie, Jr., J. L. Bullister and D. A. LeBel,
758 2007: Changes in the pool of Labrador Sea Water in the subpolar North Atlantic
759 related to changes in the upper North Atlantic circulation Geophysical Research
760 Letters 34, LO6605, doi:10.1029/2006GL028959.

761 Koltermann, K. P., V. V. Gouretski and K. Jancke, 2011: Hydrographic Atlas of the
762 World Ocean Circulation Experiment (WOCE). Volume 3: Atlantic Ocean (eds. M.
763 Sparrow, P. Chapman and J. Gould). International WOCE Project Office,
764 Southampton, UK, ISBN 090417557X.

765 Locarnini, R. A., A. V. Mishonov, J. I. Antonov, T. P. Boyer, H. E. Garcia, O. K.
766 Baranova, M. M. Zweng, and D. R. Johnson, 2010: World Ocean Atlas 2009, Volume

767 1: Temperature. S. Levitus, Ed. NOAA Atlas NESDIS 68, U.S. Government Printing
 768 Office, Washington, D.C., 184 pp.

769 Lumpkin, R., and Speer, K., 2007: Global ocean meridional overturning. *J. Phys.*
 770 *Oceanogr.* **37**, 2550–2562.

771 Masumoto, Y., H. Sasaki, T. Kagimoto, N. Komori, A. Ishida, Y. Sasai, T. Miyama, T.
 772 Motoi, H. Mitsudera, K. Takahashi, H. Sakuma, and T. Yamagata, 2004: A fifty-year
 773 eddy-resolving simulation of the world ocean: preliminary outcomes of OFES
 774 (OGCM for the Earth Simulator), *J. Earth Sim.*, 1, 35-56.

775 Masumoto, Y., 2010: Sharing the results of a high-resolution ocean general circulation
 776 model under a multi-discipline framework a review of OFES activities. *Ocean Dyn.*,
 777 60, 633–652.

778 Matano, R. P., and E. J. Beier, 2003: A kinematic analysis of the Indian/Atlantic
 779 interocean exchange. *Deep-Sea Res. II.* 50 (1), 229-249.

780 Meinen, C. S., W. E. Johns, S. L. Garzoli, E. van Sebille, D. Rayner, T. Kanzow, and M.
 781 O. Baringer, 2012: Variability of the Deep Western Boundary Current at 26.5°N
 782 during 2004-2009. *Deep-Sea Res. Part II*, 85:154-168, doi:10.1016/
 783 j.dsr2.2012.07.036.

784 Meinen, C. and S. L. Garzoli, 2014: Attribution of Deep Western Boundary Current
 785 variability at 26.5°N. *Deep Sea Research I*, 90, 81-90, doi: 10.1016/

786 j.dsr.2014.04.016.

787 Molinari, R. L., R. A. Fine, W. D. Wilson, R. G. Curry, J. Abell, and M. S. McCartney,
788 1998: The arrival of recently formed Labrador Sea Water in the Deep Western
789 Boundary Current at 26.5°N. *Geophysical Research Letters*, 25: 2249-2252.

790 Müller, T. J., Y. Ikeda, N. Zangenber, and L. V. Nonato, 1998: Direct measurements of
791 western boundary currents of Brazil between 20°S and 28°S. *J. Geophys. Res.*, **103**,
792 5429-5437.

793 Oudot, C., 1993: Total CO₂ and total alkalinity data obtained during the R/V L'Atalante
794 cruise in the Atlantic Ocean, (WOCE Section A06, 2 January–10 February 1993)
795 <http://cdiac.ornl.gov/ftp/oceans/a06woce/>, Carbon Dioxide Information Analysis
796 Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge,
797 Tennessee, doi:10.3334/CDIAC/otg.WOCE A06 1993.

798 Osynny, V., and P. Cornillon, 2004: Properties of Rossby waves in the North Atlantic
799 estimated from satellite data, *J. Phys. Oceanogr.*, 34(1), 61-76.

800 Paris, C. Helgers, J., van Sebille, E. and A. Srinivasan, 2013: Connectivity Modeling
801 System: A probabilistic modeling tool for the multi-scale tracking of biotic and
802 abiotic variability in the ocean. *Environ. Modell. Softw.*, 42, 47-54.

803 Perez, R.C., S.L. Garzoli, C.S. Meinen, and R.P. Matano, 2011: Geostrophic Velocity
804 Measurement Techniques for the Meridional Overturning Circulation and

805 Meridional Heat Transport in the South Atlantic. *J. Atmos. Ocean. Techn.*, 28:1504-
806 1521.

807 Preu, B., F. J. Hernández-Molina, R. Violante, A. R. Piola, C. M. Paterlini, T. Schwenk, I.
808 Voigt, S. Krastel and V. Speiss, 2013: Morphosedimentary and hydrographic
809 features of the northern Argentine margin: the interplay between erosive,
810 depositional and gravitational processes and its conceptual implications, *Deep-Sea*
811 *Res. I*, 75, 157–174, doi: 10.1016/j.dsr.2012.12.013.

812 Provost, C., C. Escoffier, K. Maamaatuaiahutapu, A. Kartavtseff, and V. Garçon, 1999:
813 Subtropical mode waters in the South Atlantic Ocean, *J. Geophys. Res.*, **104**,
814 21,033–21,049, doi:10.1029/1999JC900049.

815 Reid, J. L., 1989: On the total geostrophic circulation of the South Atlantic Ocean: Flow
816 patterns, tracers, and transports, *Prog. Oceanogr.*, 23, 149–244.

817 Rimaud, J., S. Speich, B. Blanke, and N. Grima, 2012: The exchange of Intermediate
818 Water in the South East Atlantic: Water mass transformations diagnosed from the
819 Lagrangian analysis of a regional ocean model. *J. Geophys. Res.*, **117**, C08034,
820 doi:10.1029/2012JC008059.

821 Sasaki, H., M. Nonaka, Y. Masumoto, Y. Sasai, H. Uehara, and H. Sakuma, 2008: An
822 eddy-resolving hindcast simulation of the quasiglobal ocean from 1950 to 2003 on
823 the Earth Simulator, in *High Resolution Numerical Modelling of the Atmosphere*

824 and Ocean, edited by K. Hamilton and W. Ohfuchi, pp. 157–185, Springer, New
 825 York.

826 Smethie, W.M., R.A. Fine, A. Putzka, E.P. Jones, 2000: Tracing the flow of North
 827 Atlantic Deep Water using chlorofluorocarbons. *Journal of Geophysical Research*,
 828 105: 14,297-14,234, 2000.

829 Schmid, C., Z. D. Garraffo, E. Johns, and S. L. Garzoli, 2003: Pathways and Variability
 830 at intermediate depths in the tropical Atlantic. ed. by G. J. Goni and P. Malanotte-
 831 Rizzoli. EOS, 68, 233-268.

832 Schmid, C., 2014: Mean vertical and horizontal structure of the subtropical circulation in
 833 the South Atlantic from three-dimensional observed velocity fields. *Deep-Sea Res.I*,
 834 91 (9), 50-71. DOI: 10.1016/j.dsr.2014.04.015.

835 Schmitz, W. J. and M. S. McCartney, 1992: On the North Atlantic circulation. *Rev. of*
 836 *Geophysics*, 31, 29-50.

837 Schott, F. A., J. P. McCreary, and G. C. Johnson, 2004: Shallow overturning circulations
 838 of the tropical-subtropical oceans. *Earth Climate: The ocean-atmosphere interaction*.
 839 C. Wang, S.- P. Xie, and J.A. Carton, Eds, *Geophys. Monogr.*, Amer. Geophys.
 840 Union, 261-304.

841 Schott, F. A., M. Dengler, R. J. Zantopp, L. Stramma, J. Fischer, and P. Brandt, 2005:
 842 The shallow and deep western boundary circulation of the South Atlantic at 5_–11_S,

843 J. Phys. Oceanogr., 35, 2031–2053.

844 Speich, S., B. Blanke, and W. Cai, 2007: Atlantic meridional overturning circulation and
845 the Southern Hemisphere supergyre. *Geophysical Research Letters*, Vol. 34, L23614,
846 doi:10.1029/2007GL031583.

847 Troupin, C., A. Barth, D. Sirjacobs, M. Ouberdous, J.-M. Brankart, P. Brasseur, M.
848 Rixen, A. Alvera Azcarate, M. Belouni, A. Capet, F. Lenartz, M.-E. Toussaint,
849 and J.-M. Beckers, 2012: Generation of analysis and consistent error fields using
850 the Data Interpolating Variational Analysis (Diva), *Ocean Modelling*, 52-53, 90-
851 101.

852 van Sebille, E., W. E. Johns, and L. M. Beal, 2012: Does the vorticity flux from Agulhas
853 rings control the zonal pathway of NADW across the South Atlantic?, *J. Geophys.*
854 *Res.*, 117, C05037, doi:10.1029/2011JC007684.

855 Wanninkhof, R., S. C. Doney, J. L. Bullister, N. M. Levine, M. Warner, and N. Gruber,
856 2010: Detecting anthropogenic CO₂ changes in the interior Atlantic Ocean between
857 1989 and 2005. *Journal of Geophysical Research*, Vol. 115, C11028,
858 doi:10.1029/2010JC006251.

859 Weiss, R. F., J. L. Bullister, R. H. Gammon, M. J. Warner, 1985: Atmospheric
860 chlorofluoromethanes in the deep equatorial Atlantic, *Nature*, 314, 608–610.

861 Zangenberg, N., and G. Siedler, 1998: Path of North Atlantic Deep Water in the Brazil

Basin, J. Geophys. Res., 103, 5419–5428, doi:10.1029/ 97JC03287.

Captions for the Figures

Figure 1: Map of CFC-11 in pmol kg^{-1} on neutral density level 27.9 kg m^{-3} , the level of the upper NADW, generated from WOCE-era data from 1992 to 1997. Brown lines indicate highlighted WOCE lines: A10 zonal line along 30°S , A16S meridional line at 25°W and A13 meridional line at 0°E . The black dots indicate locations of all of the other WOCE stations used to make this map.

Figure 2: Partial CLIVAR sections demonstrating the spread of NADW eastward into the central South Atlantic Ocean. Vertical axis is depth in meters. Panel a) partial A16S section of CFC-11 from 15° - 30°S along about 25°W from 1000-5000 m. The lowest contour for CFC-11 concentration is $0.01 \text{ pmol kg}^{-1}$. Color scale given on the right; two neutral density surfaces are 27.9 and 28.1 kg m^{-3} are identified as white lines. Panel b) Same as panel a) except for oxygen concentration in $\mu\text{mol kg}^{-1}$. Panel c) Same as panel a) except for salinity. Panel d) partial A13 section of CFC-11 from 15° - 30°S along about 5° - 10°E from 1000-5000 m. The lowest contour for CFC-11 concentration is 0.005 pmol/kg . Panel e) full cross-basin section of CFC-11 along A10, 30°S from 1000-5000 m. Note for a, d, and e that 0.3 pmol kg^{-1} is the highest concentration CFC-11 contoured and higher concentrations will appear in the same shades as for 0.3 pmol kg^{-1} .

Figure 3: Top panel: Velocity field at 2000 dbar derived from the Argo data. Red vectors

highlight the southward to southwestward flow along the western boundary; blue indicates the eastward velocity originating near the Vitória-Trindade ridge. Isobaths: 2000, 2500 and 3000 m. Solid curves highlight the pathway of the DWBC along the South American coast. Dashed lines indicate regions where the pathway is less well defined as it moves to the interior of the basin. The meridional line indicates the location of the vertical section displayed in the lower panel. Areas with no vectors or shading indicate that velocities are too small to be significant with respect to a 95% confidence interval. Lower panel: Meridional-vertical structure of the eastward pathway in the top panel showing the zonal velocity in cm sec^{-1} along 25°W from 10° to 30°S .

Figure 4: Top panels: zonal sections of the climatological mean salinity from World Ocean Atlas 2009 (WOA09, left), and from the 27-year mean OFES model section (right), both along 30°S . White color contours in the upper panels indicate neutral density levels 27.9 and 28.1 kg m^{-3} . The lower panels show the corresponding potential temperature/salinity diagrams from the WOA09 (left) and the model (right). Black contours indicate neutral density. Red indicates temperature and salinity values corresponding to the characteristics of the NADW (neutral densities 27.9 and 28.1 kg m^{-3}).

Figure 5: Some examples of the two main pathways for synthetic floats launched in the NADW layer at 5°S west of 30°W . Top panel: Floats that follow the western boundary, defined as floats that cross 30°S west of 40°W and leave the South Atlantic across 45°S . Bottom panel: Floats that follow an interior pathway by leaving the South Atlantic across 17°E after having passed through a box

defined by 9° to 10° W and 20° to 30° S. The color shading indicates the depth of the float in meters.

Figure 6: Mean velocity magnitude in cm sec^{-1} (top panels) and eddy kinetic energy (EKE) in $\text{cm}^2 \text{sec}^{-2}$ (bottom panels) calculated from the 2001 to 2006 OFES fields for two different depths: the upper boundary of the DWBC (1700 m, left) and the core of the DWBC (2500 m, right). Color bars are velocity magnitude in cm sec^{-1} (top panels) and eddy kinetic energy (EKE) in $\text{cm}^2 \text{sec}^{-2}$ (bottom panels).

Figure 7: Hovmoller diagrams of the horizontal velocity magnitude in cm sec^{-1} (left) and eddy kinetic energy (EKE) in $\text{cm}^2 \text{sec}^{-2}$ (right) calculated from the 2001 to 2006 OFES fields across 25° S and at 2500 m depth, as a function of longitude and time. Color bars are velocity magnitude in cm sec^{-1} (left) and eddy kinetic energy (EKE) in $\text{cm}^2 \text{sec}^{-2}$ (right). The vertical line at approximately 14° W indicate no data due to the presence of the Mid Atlantic Ridge.

Figure 8: Volume transports between neutral density levels 27.9 and 28.1 kg m^{-3} corresponding to NADW, and obtained from the OFES velocity fields. At 5° S the transport is integrated across the basin. At other latitudes, transports are calculated west of 30° W (from 5° to 25° S) and west of 40° W at 30° and 35° S. At the eastern side of the ocean, volume transports are calculated east of 0° E. The color shading show the zero-lag correlation between meridional velocity in the South Atlantic at a depth of 2500 m with the DWBC meridional volume transport along 34.5° S. The color scale is the correlation value from 1.0 to -1.0. Black and cyan contours show positive and negative correlations that are

significant with 95% confidence.

Caption for the Table

Table I: Meridional volume transport in Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ sec}^{-1}$, negative indicates southward), Numbers in parentheses correspond to the standard deviation. Observed values (red) were obtained from: CTD and Argo observations at 5°S , CTD observations at 30°S , XBT observations combined with CTD and Argo floats data at $\sim 35^\circ\text{S}$, and a line of PIES (pressure-equipped inverted echo sounder) moorings along 34.5°S . See text for more details.

Column 1: Eulerian transports estimated from the OFES velocity field between neutral densities 27.9 and 28.1 kg m^{-3} (black) and from observations (red) between the same density levels. Column 2: Eulerian transports estimated from the OFES velocity field between 1000 and 3500 m (black) and from observations (red) between the same levels. Column 3: Lagrangian transports estimated from the simulated float trajectory field for floats deployed between 1000 and 3500 m .

	Eulerian transports estimated between neutral density surfaces 27.9 to 28.1 (~1700 to 3000 m)	Eulerian transports estimated between depth surfaces 1000 to 3500 m	Lagrangian transports estimated from simulated floats deployed between depths of 1000 to 3500 m
From model results from coast to coast at 5°S. From CTD and Argo observations from coast to coast at 5°S	-13.9 Sv (0.96) -12.6 Sv	-16.8 Sv (1.1) -21.4 Sv	-13.2 Sv (3.2) n/a
	From 40.0° W to the west coast	From 40.0° W to the west coast	From 40.0° W to the west coast
Model results at 34.5°S From CTD observations at 30°S From combined observations at ~35°S	-7.5 Sv (0.3) -8.2 Sv -9.0 Sv (3.1)	-16.4 Sv (0.6) -14.5 Sv -19.5 Sv(5.4)	-9.4 Sv (4.2) n/a n/a
	From 0°E to east coast	From 0°E to east coast	From 0°E to east coast
Model results at 34.5°S, From combined observations at ~35°S	-4.6 Sv (0.2) -4.2 Sv (1.9)	-2.8 Sv (0.23) -3.7 Sv (1.0)	-3.0 Sv (1.4) n/a

Figure 1

Figure 2

Figure 3

Figure 4:

Figure 5

Figure 6

Figure 7

Figure 8

Table I

	Eulerian transports estimated between neutral density surfaces 27.9 to 28.1 (~1700 to 3000 m)		Eulerian transports estimated between depth surfaces 1000 to 3500 m	Lagrangian transports estimated from simulated floats deployed between depths of 1000 to 3500 m
From model results from coast to coast at 5°S. From CTD and Argo observations from coast to coast at 5°S	-13.9 Sv (0.96) -12.6 Sv		-16.8 Sv (1.1) -21.4 Sv	-13.2 Sv (3.2) n/a
	From 40.0° W to the west coast		From 40.0° W to the west coast	From 40.0° W to the west coast
Model results at 34.5°S From CTD observations at 30°S From combined observations at ~35°S	-7.5 Sv (0.3) -8.2 Sv -9.0 Sv (3.1)		-16.4 Sv (0.6) -14.5 Sv -19.5 Sv(5.4)	-9.4 Sv (4.2) n/a n/a
	From 0°E to east coast		From 0°E to east coast	From 0°E to east coast
Model results at 34.5°S, From combined observations at ~35°S	-4.6 Sv (0.2) -4.2 Sv (1.9)		-2.8 Sv (0.23) -3.7 Sv (1.0)	-3.0 Sv (1.4) n/a