CAMBRIDGE
UNIVERSITY PRESS

N-Myristoyltransferase as a potential drug target in malaria and leishmaniasis

Journal:	<i>Parasitology</i>
Manuscript ID:	PAR-2013-0006.R2
Manuscript Type:	Special Issue Review (invited contributions only)
Date Submitted by the Author:	n/a
Complete List of Authors:	Tate, Edward; Imperial College London, Chemistry Bell, Andrew; Imperial College London, Chemistry Rackham, Mark; Imperial College London, Chemistry Wright, Megan; Imperial College London, Chemistry
Key Words:	malaria, leishmaniasis, medicinal chemistry, chemical biology, plasmodium vivax, leishmania donovani, plasmodium falciparum, high throughput screening

SCHOLARONE™
Manuscripts

view

1 *Title: N-Myristoyltransferase as a potential drug target in malaria and leishmaniasis*

2 Authors: EDWARD W. TATE, ANDREW S. BELL, MARK D. RACKHAM and MEGAN H.
3 WRIGHT

4 Address: Institute of Chemical Biology, Department of Chemistry, Imperial College London, London
5 SW7 2AZ

6 Running Title: NMT as a drug target

7 Contact: Edward W Tate, Institute of Chemical Biology, Department of Chemistry, Imperial College
8 London, London SW7 2AZ Phone: 020 7594 3752 Fax: 020 7594 1139 E-mail e.tate@imperial.ac.uk

9

10 Keywords: malaria, leishmaniasis, medicinal chemistry, chemical biology, *Plasmodium vivax*,
11 *Leishmania donovani*, *Plasmodium falciparum*, high throughput screening

For Peer Review

SUMMARY

Infections caused by protozoan parasites are among the most widespread and intractable transmissible diseases affecting the developing world, with malaria and leishmaniasis being most costly in terms of morbidity and mortality. Although new drugs are urgently required against both diseases in the face of ever-rising resistance to frontline therapies, very few candidates passing through development pipelines possess a known and novel mode of action. Set in the context of drugs currently in use and under development, we present the evidence for *N*-myristoyltransferase (NMT), an enzyme that *N*-terminally lipidates a wide range of specific target proteins through post-translational modification, as a potential drug target in malaria and the leishmaniases. We discuss the limitations of current knowledge regarding the downstream targets of this enzyme in protozoa, and our recent progress towards potent cell-active NMT inhibitors against the most clinically-relevant species of parasite. Finally, we outline the next steps required in terms of both tools to understand *N*-myristoylation in protozoan parasites, and the generation of potential development candidates based on the output of our recently-reported high-throughput screens.

INTRODUCTION

Malaria

Malaria is a disease caused by infection of a human host with protozoan parasites of the genus *Plasmodium*, and is a devastating global health issue with approximately 200 million cases and 1 million deaths in 2010 alone (Murray *et al.* 2012). The complex life cycle of malaria parasites spreads across two hosts and five host tissues whilst undergoing at least ten distinct morphological transitions (Sturm *et al.* 2006; Mackinnon and Marsh 2010). Replication of parasites and subsequent rupture of erythrocytes in the intra-erythrocytic stages are responsible for the clinical symptoms of malaria, and the majority of drugs target these asexual (human-host) stages of the life cycle. Some species of malaria, most notably *Plasmodium vivax*, can exist in a latent liver hypnozoite form that can cause relapse even after clearance of bloodstream parasites (Derbyshire *et al.* 2012; Rodrigues *et*

38 *al.* 2012). Of the five relevant species of human parasite, the vast majority of deaths occur from *P.*
39 *falciparum* infections, which is the typical cause of severe malaria (Claessens *et al.* 2012). This has
40 led to the majority of drug discovery efforts focussing on *P. falciparum*, typically at the expense of
41 other species. Although the demand for new *P. falciparum* drugs is in no doubt, *P. vivax* is
42 responsible for the majority of worldwide malaria endemicity (Price *et al.* 2009; WHO 2011).
43 However, difficulties culturing the parasite (Udomsangpetch *et al.* 2007) along with challenges
44 imaging and targeting the hypnozoite liver stages (Meister *et al.* 2011) have led to a dearth of new *P.*
45 *vivax* drugs (Price *et al.* 2009). Medications capable of targeting all relevant species of parasite, and
46 crucially clearing liver-stage parasites, are in great demand.

47 For the latter half of the 20th century, antimalarial drug discovery was a success story for natural
48 product-inspired therapies, by far the most widely used of which are chloroquine (Loeb *et al.* 1946)
49 and artemisinin (Miller and Su 2011).

50 Chloroquine was first discovered as a derivative of antimalarial natural product quinine (Krafts *et al.*
51 2012), and has been a first-line antimalarial for over sixty years (Loeb *et al.* 1946). Studies over the
52 past twenty years have shown that this class of compounds (with the exception of primaquine) is
53 involved in the disruption of haem detoxification by the parasite (Weissbuch and Leiserowitz 2008).
54 Artemisinin is a highly effective antimalarial natural product, isolated from *Artemisia annua* (Miller
55 and Su 2011); its antimalarial action is still under debate, but most hypotheses involve reductive
56 activation of the endoperoxide moiety resulting in parasite death from oxidative damage (O'Neill and
57 Posner 2004; Li and Zhou 2010; Slack *et al.* 2012). Other drug classes used in the treatment of
58 malaria to varying extents include pyrimidine biosynthesis disruptors, drugs that target the apicoplast
59 (Botté *et al.* 2012) and drugs discovered by phenotypic screening with unknown targets (e.g.
60 primaquine) (Kappe *et al.* 2010). Despite the apparent plethora of antimalarials, drug resistance is a
61 major issue and new medications with distinct mechanisms are constantly required to combat the
62 continued evolution of the parasite (Fidock 2010; Mackinnon and Marsh 2010). This is compounded
63 by the emergence of resistance to the artemisinins in Asia, (Dondorp *et al.* 2009; Phyto *et al.* 2012)
64 reinforcing the urgent requirement for new therapies. Fortunately, a great deal of resource has been

65 directed towards antimalarial drug discovery in the past few decades. Elaboration of the artemisinin
66 pharmacophore has resulted in multiple clinical candidates (Vennerstrom *et al.* 2004; O'Neill *et al.*
67 2010; Charman *et al.* 2011), although the potential effectiveness of these compounds in artemisinin-
68 resistant regions remains a concern due to the shared mechanism of action. In addition, novel
69 inhibitors of the pyrimidine biosynthetic pathway are in development (Painter *et al.* 2007) and
70 phenotypic high-throughput screens have resulted in a wealth of information on relevant scaffolds
71 (Plouffe *et al.* 2008; Gamo *et al.* 2010; Guiguemde *et al.* 2010), and yielded promising clinical
72 candidates (Rottmann *et al.* 2010). Among the numerous clinical and preclinical candidates for the
73 treatment for malaria, the vast majority work by existing or unknown mechanisms and are based on
74 known pharmacophores; indeed many are new combinations of existing marketed drugs. Although
75 some have shown efficacy against resistant strains, concerns remain that resistance may develop
76 quickly against already vulnerable mechanisms. New drugs that work by distinct novel biological
77 mechanisms are therefore highly desirable.

78 *The leishmaniasis*

79 The leishmaniasis are the second most prevalent class of parasitic infection after malaria, giving rise
80 to >2 million new cases each year. The disease occurs in three forms, cutaneous (CL), muco-
81 cutaneous (MCL) and the most fatal form, visceral leishmaniasis (VL). The latter is associated with
82 infection by the species *Leishmania donovani*, while the cutaneous forms are due to infection by
83 multiple species including *L. major*, *L. braziliensis* and *L. mexicana*. The leishmaniasis are endemic
84 in more than 90 countries around the world, being particularly prevalent in India, East Africa,
85 Bangladesh and Brazil. Additional clinical issues include post-kala-azar dermal leishmaniasis
86 (PKDL), occurring after the apparent drug cure of VL in certain geographical regions (e.g. Sudan) and
87 difficult to cure with pentavalent antimonials. Combination therapies with therapeutic vaccines
88 (Maroof *et al.* 2012) or immune-response activating drugs, such as imiquimod (Arevalo *et al.* 2001),
89 show some promise. In contrast to the wealth of treatments and drugs in development for malaria, the
90 leishmaniasis are poorly provided for. None of the currently available drugs (Fig. 2) were discovered
91 by a rational design process for this neglected disease, and suffer from drawbacks including lack of an

oral formulation, prolonged treatment times, high cost of treatment, toxicity, teratogenicity and/or increasing drug resistance. In addition, all work through unknown mechanisms, by disruption of cell membranes or through unspecific antibiotic effects. There are very few drugs in development for these conditions with the most advanced being another antibiotic, fexinidazole (Winkelmann and Raether 1978), discovered as part of a repurposing initiative (Wyllie *et al.* 2012).

PROTEIN *N*-MYRISTOYLTRANSFERASE

The post-translational modification (PTM) of proteins contributes hugely to the chemical and functional diversity of the cellular proteome and results in the incorporation of molecular motifs not directly encoded by the genome. Protein *N*-myristoylation is the attachment of a 14-carbon saturated fatty acid, myristate, to the N-terminal glycine residue in a specific set of cellular proteins, catalysed by the enzyme myristoyl CoA:Protein *N*-myristoyltransferase, NMT (Fig. 3) (Wright *et al.* 2010). Whilst *N*-myristoylation is often referred to as a PTM, it most commonly occurs co-translationally. Post-translational myristoylation is less well documented but is known to occur following exposure of an internal glycine after cleavage of proteins by caspases during the apoptotic cascade (Zha *et al.* 2000). *N*-Myristoylation can be involved in protein stability, protein-protein interaction interfaces and association of proteins with membranes.

NMT appears to be ubiquitous in eukaryotes, including fungi (Towler *et al.* 1987; Lodge *et al.* 1994; Shaw *et al.* 2002), insects (Ntwasa *et al.* 1997), plants (Boisson *et al.* 2003), mammals (including mouse, rat, cow and human) and the parasitic protozoa *P. falciparum* (Gunaratne *et al.* 2000), *L. major* (Price *et al.* 2003), *L. donovani* (Brannigan *et al.* 2010) and *Trypanosoma brucei* (Price *et al.* 2003). NMT has been shown to be essential for survival in the bloodstream form of *T. brucei* (Price *et al.* 2003; Price *et al.* 2010), in insect stages of *L. major* (Price *et al.* 2003) and *L. donovani* (Brannigan *et al.* 2010) and most recently in the rodent malaria parasite *P. berghei* (Pino *et al.* 2012).

NMT structure and mechanism

The enzyme catalytic cycle has been well-studied in yeast, and follows a Bi-Bi mechanism (Towler *et al.* 1987; Rudnick *et al.* 1991). Myristoyl-CoA (Myr-CoA) binds to the apo-enzyme, inducing a

118 conformational change that allows the protein substrate to bind. Myristate is transferred by attack of
119 the N-terminal glycine amine of the peptide on the thioester carbonyl of Myr-CoA (Rudnick *et al.*
120 1991); CoA is released followed by the myristoylated substrate. The first crystal structure of an NMT
121 to be published was *Candida albicans* NMT (CaNMT) (Weston *et al.* 1998). Crystal structures of
122 *Saccharomyces cerevisiae* (ScNMT) (Bhatnagar *et al.* 1998; Farazi *et al.* 2001b), and more recently
123 *Leishmania donovani* NMT, (LdNMT) (Brannigan *et al.* 2010), *L. major*, (LmNMT) (Frearson *et al.*
124 2010), and the *P. vivax* enzyme (Goncalves *et al.* 2012b), followed. These structures and others
125 provide insight into the binding sites of Myr-CoA, peptide substrates and inhibitors. Published
126 structures are consistent with a highly conserved Myr-CoA binding mode, with Myr-CoA binding in a
127 bent 'question mark' conformation (Fig. 4b). The thioester carbonyl is placed into an 'oxyanion hole',
128 which activates it for nucleophilic attack, and the fatty acyl chain of Myr-CoA inserts into a deep,
129 hydrophobic pocket. The peptide N-terminal glycine ammonium interacts electrostatically with the
130 buried carboxylate of the C-terminal enzyme residue (Farazi *et al.* 2001b), which is responsible for
131 deprotonation of the ammonium so that the generated nucleophilic amine can attack the Myr-CoA
132 thioester (Farazi *et al.* 2001a; Farazi *et al.* 2001b).

133 *NMT substrate specificity*

134 NMT appears to be highly specific for transfer of C14 fatty acid, tolerating only slight changes to
135 chain length (reviewed in (Wright *et al.* 2010)). However, peptide substrate specificity is complex and
136 there is no definitive “myristoylation motif”, beyond the requirement for an N-terminal glycine. This
137 requirement may be mechanism-based: Gordon and co-workers have suggested that rotation of the
138 peptide N-terminal amine about the peptide backbone aligns it for attack on the thioester, and that
139 such a rotation may be hindered for residues with β -substituents, i.e. any amino acid except for
140 glycine (Farazi *et al.* 2001b). Maurer-Stroh and co-workers used crystal structures and biochemical
141 data to develop a myristoylation predictive tool: the MYR predictor (Maurer-Stroh *et al.* 2002a;
142 Maurer-Stroh *et al.* 2002b), which suggested that as many as 17 residues may be involved in substrate
143 recognition by NMT. A second tool for predicting *N*-myristoylation, the Myristoylator, is based upon
144 a different model for prediction (neural networks, or machine learning) but generates similar error
145 rates compared to the MYR predictor (Bologna *et al.* 2004). Both predictors are necessarily based on
146 the set of proteins annotated or predicted to be *N*-myristoylated using sequence similarity in
147 SwissProt. Unlike the acyl-CoA binding site, the peptide pocket is not well conserved across NMTs
148 from different species (Maurer-Stroh *et al.* 2002b). This, together with the observation that NMT is
149 essential for survival, means that this pocket is a target for selective NMT inhibitors
150 (Georgopapadakou 2002; Price *et al.* 2003).

151 **PROTEIN *N*-MYRISTOYLATION**

152 Myristoylated proteins have been estimated to make up between 0.5 % and 3 % of eukaryotic cellular
153 proteomes, depending on the species and *in silico* model used (Maurer-Stroh *et al.* 2002a; Martinez *et*
154 *al.* 2008). In the majority of myristoylated proteins studied so far, myristate has a role in transient
155 membrane localisation. The ‘two-signal’ membrane binding model suggests that strong membrane
156 localisation is only achieved when a second feature of the protein complements myristoylation (Resh
157 1994). This may be a second acyl group near the N-terminus, such as palmitate (which is usually
158 attached to the side-chain of cysteine residues), a polybasic cluster of amino acids that interacts with

membrane phospholipid acidic head groups, or a domain that interacts with another membrane-bound protein (Resh 2006a). Membrane binding in some myristoylated proteins is dynamically regulated via so-called myristoyl-switches, as in the ARF-GTPases, where a change in the ligand bound (from GTP to GDP) causes a conformational change that exposes a hydrophobic pocket that binds myristate, sequestering the fatty acyl chain so that it can no longer interact with the membrane bilayer (Goldberg 1998). Myristate-mediated membrane binding is clearly an important mechanism and numerous studies have concluded that it often contributes to the function or regulation of the target protein.

Detecting N-myristoylation in protozoan parasites

Predicting and validating the *N*-myristoylation of potential substrate proteins in general is challenging due to the complex substrate specificity of NMT and difficulties inherent in detecting protein lipidation. Some proteins can be assigned as likely NMT substrates based on homology; an example being the ARF-GTPases, a class of proteins present in all eukaryotes, having common roles and known to be *N*-myristoylated in many organisms (Donaldson and Jackson 2011). For protozoan parasite proteins sharing no sequence identity with generic eukaryotic proteins the main recourse is bioinformatic prediction. However, these tools are necessarily based on known *N*-myristoylated proteins, of which few have been reported in protozoan parasites, and many of these predictions still require experimental proof. Demonstrating *N*-myristoylation of a protein in its native context is non-trivial, and thus non-native approaches predominate. A candidate protein is often over-expressed as a GFP or other tagged construct, or the protein of interest is co-expressed with NMT in *E. coli*, and metabolic radiolabelling with myristate or mass spectrometry is used to demonstrate myristoylation. Lipidation of the protein of interest shown by mass spectrometry, radiolabelling or well characterised chemical probes in the wild type parasite constitute the only methods for direct proof of *N*-myristoylation in the native context, whereas lipidation of an overexpressed construct is good evidence. Other data, based on co-expression of the protein with NMT in *E. coli* or the effects of mutagenesis on membrane localisation are merely suggestive of *N*-myristoylation.

Protein myristoylation in Plasmodium species

185 Relatively little is known about which proteins are *N*-myristoylated in *Plasmodium* species. *P.*
186 *falciparum* possesses a single NMT isoform (Gunaratne *et al.* 2000) which is able to transfer
187 myristate from Myr-CoA to a peptide substrate based on *Pf*ARF1 (Gunaratne *et al.* 2000).
188 Experimentally studied substrates have roles in life cycle regulation or progression (calcium
189 dependent protein kinase 1 [CDPK1] and Calpain) (Moskes *et al.* 2004; Russo *et al.* 2009a; Russo *et*
190 *al.* 2009b), host cell invasion (45 kDa glideosome associated protein [GAP45]) (Rees-Channer *et al.*
191 2006), trafficking (ARF1) (Leber *et al.* 2009), Golgi function (GRASP1) (Struck *et al.* 2005) and
192 energy metabolism (Adenylate kinase 2, AK2) (Rahlfs *et al.* 2009). However, only GAP45 and
193 CDPK1 have been shown to be *N*-myristoylated in their native context (Moskes *et al.* 2004; Rees-
194 Channer *et al.* 2006), whilst evidence for *N*-myristoylation of Calpain is based on radiolabelling of a
195 Calpain-GFP construct (Russo *et al.* 2009a). For other potential targets such as Armadillo repeats only
196 protein (ARO), AK2 and GRASP1, evidence is limited (Struck *et al.* 2005; Rahlfs *et al.* 2009;
197 Cabrera *et al.* 2012). CDPK1 requires *N*-myristoylation for membrane localisation (Moskes *et al.*
198 2004), and its gene cannot be knocked out in *P. falciparum* or the rodent parasite *P. berghei*, implying
199 essentiality (Kato *et al.* 2008; Tewari *et al.* 2010). In addition, CDPK1 has key functions in multiple
200 stages of the parasite life cycle and is involved in translational activation during sexual development
201 (Sebastian *et al.* 2012). Another potentially myristoylated kinase, CDPK4, has been shown to be
202 essential for sexual reproduction and mosquito transmission in *P. berghei* (Billker *et al.* 2004).
203 GAP45 is an *N*-myristoylated protein with a direct role in parasite invasion of RBCs. It is localised at
204 the inner membrane complex (IMC), a series of membrane structures lying beneath the parasite
205 plasma membrane (PM) (Jones *et al.* 2006). An actomyosin motor, located between the IMC and the
206 PM, drives merozoite invasion, allowing the parasite to enter the RBC (Baum *et al.* 2006). In
207 *Toxoplasma gondii*, GAP45 is essential to the function of the motor, and therefore for host cell egress,
208 motility and invasion: it has a role in the recruitment of the motor complex and there is evidence for a
209 structural role in maintaining pellicle cohesion during invasion, presumably holding the IMC and PM
210 together (Frenal *et al.* 2010). Recent data on the localisation of *Pf*GAP45 and N- or C-terminal
211 mutants are consistent with this role in spanning the IMC-PM gap (Ridzuan *et al.* 2012).

212 *Protein myristoylation in Leishmania species*

213 As with *Plasmodium*, very few proteins have been experimentally validated as *N*-myristoylated in
214 *Leishmania* species. Known NMT substrates include proteins involved in trafficking (ARL1 (Sahin *et al.*
215 *et al.* 2008)) and proteins of unknown function, such as HASPB, a member of a family of hydrophilic
216 acylated surface proteins expressed in the host and required for parasite development in the insect
217 vector (Denny *et al.* 2000; Sadlova *et al.* 2010). A family of ‘small myristoylated proteins’ with
218 probable functions at the flagellum (Tull *et al.* 2004; Tull *et al.* 2012) and a protein phosphatase
219 (PPEF) (Mills *et al.* 2007) have also been reported. In the related trypanosomatid parasites
220 *Trypanosoma brucei* and *T. cruzi*, known likely substrates include *Tb*ARF1 (Price *et al.* 2007),
221 *Tb*ARL1 (Price *et al.* 2005), *Tb*ARL6 (Price *et al.* 2012), cytoskeletal protein *Tb*CAP5.5 (Hertz-
222 Fowler *et al.* 2001), which is involved in cell morphogenesis (Olego-Fernandez *et al.* 2009), a
223 flagellar-localised protein (*Tc*FCaBP) (Godsel and Engman 1999) and a metacaspase implicated in
224 virulence (Proto *et al.* 2011). *N*-Myristoylation of native protein has only been demonstrated directly
225 for *Ld*SMP1 (Tull *et al.* 2004), *Tc*FCaBP (Godsel and Engman 1999), *Tb*CAP5.5 (Hertz-Fowler *et al.*
226 2001) and *Tb*ARL6 (Price *et al.* 2012), but there is evidence for lipidation of *Lm*PPEF (Mills *et al.*
227 2007) and HASPB (Denny *et al.* 2000) in *Leishmania*. In all other cases evidence is limited.
228 Myristoylation of *Ld*ARL1 is essential for localisation to the Golgi (Sahin *et al.* 2008) and ARL3,
229 which is involved in maintaining the flagellum of promastigotes, is also thought to be myristoylated
230 (Cuvillier *et al.* 2000). A number of other proteins have been shown to carry a dual acylation motif,
231 including HASPB, which localises to the outer leaflet of the plasma membrane in infective stages
232 (Denny *et al.* 2000). *N*-terminal *N*-myristoylation and internal *S*-palmitoylation are both required for
233 this targeting. Similarly, SMP-1 is flagellum-targeted by myristoylation and palmitoylation (Tull *et al.*
234 2004). A bioinformatic approach predicted around 60 *N*-myristoylated proteins in *Leishmania* (Mills
235 *et al.* 2007), many of which are of unknown function and share little identity with other eukaryotic
236 proteins, suggestive of parasite-specific roles.

237 *Current challenges in defining the N-myristoylated parasite proteome*

238 The handful of identified NMT substrates reflects what is known in other organisms: that *N*-
239 myristoylation is involved in crucial cellular processes. However, a comprehensive understanding of
240 the *N*-myristome and the roles of *N*-myristoylated proteins in protozoa, particularly in infective stages,
241 is still lacking. This is partly due to technical limitations associated with these organisms – for
242 example, challenges in the genetic manipulation of *P. falciparum* and *L. donovani* intracellular human
243 infective stages, and a lack of functional analysis of many of the target proteins – but also because
244 detecting lipidation of a native protein is inherently difficult. Radiolabelling with [³H]myristate or
245 other fatty acids such as palmitate with detection by fluorography is the most common traditional
246 method for studying protein acylation, but is a laborious process due to the very long (weeks or
247 months) exposure times required. Mass spectrometry is a powerful and continually advancing
248 technique that can be used to detect PTMs such as fatty acylation, but generally the protein of interest
249 must first be highly enriched and lipophilic fatty acylated proteins can be lost during preparation
250 (Resh 2006b). The concentration range of proteins within cells is huge – around 5-6 orders of
251 magnitude (Tyers and Mann 2003) – and this further hinders detection of rare proteins in complex
252 mixtures. Chemical proteomic approaches have revolutionised the field of post-translational
253 modification in the past decade, particularly for low abundance PTMs such as *N*-myristoylation. Acyl
254 biotin exchange chemistry (ABE), where palmitate is exchanged site-specifically for biotin at the site
255 of PTM (Roth *et al.* 2006), is widely used for analysing *S*-palmitoylation and was recently applied in
256 *P. falciparum* blood stages with the identification of several thousand new potential targets (Jones *et al.*
257 *et al.* 2012). Another methodology, the bioorthogonal probe approach, involves the metabolic
258 incorporation of a PTM substrate analogue containing a small, biologically inert chemical tag into
259 proteins in live cells; the tag is subsequently functionalised with useful labels for detection or
260 identification (Wright *et al.* 2010; Hang *et al.* 2011). These approaches have been used to great effect
261 for the profiling of acylated proteins in mammalian cells and tagged lipid analogues were recently
262 applied to identify *S*-palmitoylated proteins in *P. falciparum* (in parallel with ABE) (Jones *et al.* 2012)
263 and to demonstrate *N*-myristoylation of *T. brucei* ARL6, a protein with a putative role in flagellum
264 biogenesis (Price *et al.* 2012). Chemical probes have the potential to greatly expand the list of *N*-

myristoylated proteins in parasitic protozoa and contribute to our understanding of NMT as a drug target.

TOWARDS NMT INHIBITORS AS ANTIMALARIAL OR ANTILEISHMANIAL AGENTS

Several lines of evidence suggest that NMT is a promising drug target for malaria and leishmaniasis; it is a monomeric enzyme carrying out a specific modification on substrates involved in diverse and essential pathways, it is essential for viability where genetic validation has been possible, and is constitutively expressed. Furthermore it is genetically and chemically validated as a drug target in *T. brucei*, where small molecule inhibitors have been shown to be effective in animal models (Frearson *et al.* 2010). The wide variety of NMT substrates may limit the potential for resistance to develop against inhibitors targeting the protein binding site, since mutations in this site could inhibit correct myristoylation of substrates.

Initial research within our group focused on the discovery of plasmodial NMT inhibitors as chemical probes and potential therapeutic agents for the treatment of malaria, due to the availability of chemical starting points for this indication. Drug repositioning, the adaptation of an existing drug for a new indication, is often used to bypass the significant cost of clinical trials as the safety/pharmacokinetic data for these compounds has already been established (Sleigh and Barton 2010). A similar approach can be used for hit discovery (the “piggy-back approach”) to discover promising hit series without the cost of an HTS campaign (Gelb *et al.* 2003). In the case of NMT, a wealth of information has been collated with the purpose of generating antifungal NMT inhibitors. It was hypothesised that this information could be used as a valuable resource in the generation of parasitic NMT inhibitors, since *Ca*NMT and *Pf*NMT display 38% identity and 65% similarity.

In the process of validation of a radioactive assay for monitoring *N*-myristoylation, a library of 43 *Ca*NMT inhibitors containing a benzothiazole scaffold (provided by Pfizer) were screened against *Pf*NMT and *Homo sapiens* NMT1 (*Hs*NMT1) (Bowyer *et al.* 2007). Of these 43 compounds, dose-response curves were generated for 7 of the most promising inhibitors, four of which reduced parasitaemia *in vitro*. These compounds had weak enzyme affinity and cellular potency, and displayed

very high molecular weight and lipophilicity for compounds with this level of activity. Ligand Efficiency (LE) is a commonly-used measure of how tightly a compound binds to a target protein, relative to its overall size (Hopkins *et al.* 2004; Bembenek *et al.* 2009); LE > 0.35 is considered favourable, as it suggests scope for substantial optimisation. However, LE in this series was around 0.21, limiting the potential for future development. A distinct small library of 25 previously described *Ca*NMT and *Tb*NMT inhibitors was also tested against *Pf*NMT *in vitro* and compound **RO-09-4609** emerged as a promising hit compound, with weak but selective *Pf*NMT affinity. This hit was optimised by iterative medicinal chemistry, resulting in a moderate affinity and highly selective compound **Example 26** (Yu *et al.* 2012), representing a 100-fold affinity improvement over the initial hit (Fig. 5A). The binding mode of these compounds was validated in *Pv*NMT (81% sequence identity to *Pf*NMT), confirming the hypothesis that these compounds are competitive with the peptide substrate; however, LE (0.29) remained sub-optimal. A scaffold hopping approach was then applied to yield 2,3-benzothiophene **Example 9** (Rackham *et al.* 2012), resulting in the most ligand efficient inhibitor discovered at this stage of development against *Pf*NMT (Fig. 5A). Crystallography of this inhibitor series bound to *Pv*NMT confirmed that the 2,3-benzothiophene inhibitors occupy a distinct binding mode to the 2,3,4 benzofurans exemplified by **Example 26** (Yu *et al.* 2012), representing a novel inhibitor series for further development (Fig. 5B) (Rackham *et al.* 2012).

Whilst the “piggy-back” approach provided a highly ligand-efficient series of *Pf*NMT inhibitors, cross-screening the same set of compounds against *L. major* NMT failed to identify any starting points (Panethymitaki *et al.* 2006), necessitating alternative hit generation strategies. High-throughput screening has been successful in identifying multiple series of NMT inhibitors, including *C. albicans* (Masubuchi *et al.* 2001; Ohtsuka 2003), *Aspergillus fumigatus* (Bowyer *et al.* 2008), *Cryptococcus neoformans* (Bowyer *et al.* 2008) and *T. brucei* (Frearson *et al.* 2010). Although compounds from the latter publication have been co-crystallised with *L. major* NMT and display excellent enzyme affinity, the cellular potency of this series has not been disclosed. Consequently, we initiated a broader screening programme to identify additional selective inhibitors of *Plasmodium* and *Leishmania* NMTs. We screened the 150,000 compound Pfizer Global Diverse Representative Set against

*Ld*NMT, anticipating that hits would be likely to possess pan-*Leishmania* NMT activity since the residues involved in the binding pocket are completely conserved (Brannigan *et al.* 2010). We also screened the same set against *Pf*NMT, and the initial hit set was supplemented by analogues selected from the remainder of the Pfizer file (~ 2.5 million compounds at the time of the high-throughput screen) (Bell *et al.* 2012). In addition, a separate library of 60,000 compounds was screened against *Pv*NMT, in collaboration with MRC Technology, in the hope of identifying distinct hit compounds (Goncalves *et al.* 2012a).

Comparison of the hits from each HTS (Fig. 6) shows a remarkably wide range of structural features. Although many possess a basic centre, a common pharmacophore of several previously described NMT inhibitors, the hits feature primary, secondary and tertiary amines, which may be interacting with the conserved C-terminal Leu of the enzyme. However, the MRCT HTS hit (**MRT00057965**) is not basic and makes an alternative H-bond interaction with Ser319 (Goncalves *et al.* 2012b). Based on the diversity of known binding modes for NMT inhibitors, it is difficult to predict those for novel inhibitor series, and structures of the remaining HTS hits are, as yet, unreported. A limited medicinal chemistry optimization programme based on the MRT hit has been published (Goncalves *et al.* 2012b), resulting in a confirmed hit with improved physical properties and moderate selectivity over the human isoforms. In order to address the enzyme selectivity issue, the Pfizer screen hit set was also tested in dose-response assays against both human NMT isoforms and against *T. brucei* NMT (Bell *et al.* 2012). Consistent with our previous findings, *Ld*NMT seems distinct from the other NMTs in our screening panel, as most hits show excellent selectivity. In contrast, selectivity for *Pf*NMT over either human NMT is more elusive, though the screen did identify two hit series with encouraging profiles.

The structural basis for the observed selectivities remains unclear. Analysis of the peptide-binding pocket of *Ld*NMT identified two residue differences with *T. brucei* NMT (Brannigan *et al.* 2010), and the same residues are also points of differentiation between *Ld*NMT and both human NMTs. In contrast, the residues lining the binding pocket are conserved between *Pf*NMT and human NMT with the exception of a conservative F334Y change, which is also a point of difference between *Pf*NMT and *P. vivax* NMT. *Plasmodium*/human NMT selectivity has been attributed to differences in their

ability to tolerate a conformational change in Y296 (*Hs*NMT1 numbering) (Yu *et al.* 2012). It remains to be seen whether this difference in conformation is observed with other *Plasmodium*-selective inhibitors.

FUTURE PERSPECTIVES

Two significant hurdles remain for the validation of NMT as a drug target in malaria and leishmaniasis: the challenge of understanding the role of *N*-myristoylation through understanding of its downstream protein substrates, and proving its essentiality in clinically-relevant parasites.

Protein *N*-myristoylation is a mechanism used universally by eukaryotes to direct protein localisation and hence function, but detecting protein lipidation by traditional methods presents challenges, particularly in *P. falciparum* and *L. donovani* which are not easily genetically manipulated and go through distinct life cycle stages in a variety of host environments. Bioinformatic prediction suggests that many proteins with diverse and unknown roles are likely to be myristoylated in parasitic protozoa, but in very few cases has acylation been experimentally validated. The development of new techniques to profile protein lipidation, such as the application of bioorthogonal chemical probes (Heal and Tate 2010), may allow us to gain a much wider and in-depth understanding of the myristoylated proteome and to explore the downstream effects of NMT inhibition in these important human pathogens.

Demonstration of essentiality and druggability of NMT in *P. falciparum* and *L. donovani* requires selective inhibitors, and proof that these compounds act on-target in parasites. Progress towards selective and potent parasite NMT inhibitors is at an exciting stage; **Example 9** (Fig. 5) derived from the “piggy-back” approach represents a promising starting point for inhibitor development, demonstrating excellent ligand efficiency and selectivity over the human orthologues. Future development of this series will focus on affinity enhancements whilst maintaining optimum physicochemical properties for drug-like molecules, with the aim of generating a high value lead series for clinical development. Replacement of the alkyl ester in **Example 9** with a more biologically stable isostere is a paramount objective for further development since oral administration is a

prerequisite for a malaria medication. In addition, the other *Plasmodium*-selective hits identified (Fig. 6) represent highly promising series for further development, demonstrating a range of chemotypes with varying physicochemical properties. Successful elaboration of any of these hit series into a potent, selective and drug-like inhibitor of NMT would enable investigation in cellular and *in vivo* models of malaria, providing a chemical tool for the validation of NMT as a drug target in malaria infections. The need for new drugs against leishmaniasis is still more pressing, particularly in view of the relatively neglected nature of this disease. Our recent screening initiative has also opened the door to the development of potent, selective inhibitors of *Leishmania* NMT, and we anticipate that the available chemical matter will enable both the development of tools to explore the role and importance of NMT in this challenging organism, and new starting points for discovery of antileishmanial drugs.

The historic track record of targeted approaches to treatment of parasitic infections is in general relatively poor, since there are significant barriers to overcome in achieving good translation of enzyme to cellular activity and *in vivo* efficacy. Future work in our laboratory is focused on overcoming these hurdles to confirm the relevance of NMT inhibition as a valid target for treating parasitic infections.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge members of the Tate, Leatherbarrow, Holder, Smith and Wilkinson labs at Imperial College, MRC National Institute for Medical Research, and the University of York for their advice and assistance with proof reading, and the reviewers for their constructive comments during the preparation of this manuscript.

FINANCIAL SUPPORT

The authors would like to acknowledge the Wellcome Trust (grant no. 087792), the Medical Research Council (grant nos. 0900278 and U117532067), Imperial College, EPSRC and Pfizer for supporting this work.

REFERENCES

- Arevalo, I., Ward, B., Miller, R., Meng, T.-C., Najar, E., Alvarez, E., Matlashewski, G. and Alejandro, L.-C.** (2001). Successful Treatment of Drug-Resistant Cutaneous Leishmaniasis in Humans by Use of Imiquimod, an Immunomodulator. *Clinical Infectious Diseases* **33**, 1847-1851.
- Baum, J., Richard, D., Healer, J., Rug, M., Krnajski, Z., Gilberger, T. W., Green, J. L., Holder, A. A. and Cowman, A. F.** (2006). A conserved molecular motor drives cell invasion and gliding motility across malaria life cycle stages and other apicomplexan parasites. *Journal of Biological Chemistry* **281**, 5197-5208.
- Bell, A. S., Mills, J. E., Williams, G. P., Brannigan, J. A., Wilkinson, A. J., Parkinson, T., Leatherbarrow, R. J., Tate, E. W., Holder, A. A. and Smith, D. F.** (2012). Selective Inhibitors of Protozoan Protein *N*-myristoyltransferases as Starting Points for Tropical Disease Medicinal Chemistry Programs. *PLoS Neglected Tropical Diseases* **6**, e1625.
- Bembenek, S. D., Tounge, B. A. and Reynolds, C. H.** (2009). Ligand efficiency and fragment-based drug discovery. *Drug Discovery Today* **14**, 278-283.
- Bhatnagar, R. S., Futterer, K., Farazi, T. A., Korolev, S., Murray, C. L., Jackson-Machelski, E., Gokel, G. W., Gordon, J. I. and Waksman, G.** (1998). Structure of *N*-myristoyltransferase with bound myristoylCoA and peptide substrate analogs. *Nature Structural Biology* **5**, 1091-1097.
- Billker, O., Dechamps, S., Tewari, R., Wenig, G., Franke-Fayard, B. and Brinkmann, V.** (2004). Calcium and a calcium-dependent protein kinase regulate gamete formation and mosquito transmission in a malaria parasite. *Cell* **117**, 503-514.
- Boisson, B., Giglione, C. and Meinnel, T.** (2003). Unexpected protein families including cell defense components feature in the *N*-myristoylome of a higher eukaryote. *Journal of Biological Chemistry* **278**, 43418-43429.
- Bologna, G., Yvon, C., Duvaud, S. and Veuthey, A. L.** (2004). N-Terminal myristoylation predictions by ensembles of neural networks. *Proteomics* **4**, 1626-1632.
- Botté, C. Y., Dubar, F., McFadden, G. I., Maréchal, E. and Biot, C.** (2012). *Plasmodium falciparum* Apicoplast Drugs: Targets or Off-Targets? *Chemical Reviews* **112**, 1269-1283.

- 424 **Bowyer, P. W., Gunaratne, R. S., Grainger, M., Withers-Martinez, W., Wickramasinghe, S. R.,**
425 **Tate, E. W., Leatherbarrow, R. J., Brown, K. A., Holder, A. A. and Smith, D. F. (2007).**
426 Molecules incorporating a benzothiazole core scaffold inhibit the *N*-myristoyltransferase of
427 *Plasmodium falciparum*. *Biochemical Journal* **408**, 173–180.
- 428 **Bowyer, P. W., Tate, E. W., Leatherbarrow, R. J., Holder, A. A., Smith, D. F. and Brown, K. A.**
429 (2008). *N*-myristoyltransferase: a prospective drug target for protozoan parasites. *ChemMedChem* **3**,
430 402–408.
- 431 **Brannigan, J. A., Smith, B. A., Yu, Z., Brzozowski, A. M., Hodgkinson, M. R., Maroof, A.,**
432 **Price, H. P., Meier, F., Leatherbarrow, R. J., Tate, E. W., Smith, D. F. and Wilkinson, A. J.**
433 (2010). *N*-myristoyltransferase from *Leishmania donovani*: structural and functional characterisation
434 of a potential drug target for visceral leishmaniasis. *Journal of Molecular Biology* **396**, 985–999.
- 435 **Cabrera, A., Herrmann, S., Warszta, D., Santos, J. M., John Peter, A. T., Kono, M., Debrouver,**
436 **S., Jacobs, T., Spielmann, T., Ungermann, C., Soldati-Favre, D. and Gilberger, T. W. (2012).**
437 Dissection of minimal sequence requirements for rhoptry membrane targeting in the malaria parasite.
438 *Traffic* **13**, 1335–1350.
- 439 **Charman, S. A., Arbe-Barnes, S., Bathurst, I. C., Brun, R., Campbell, M., Charman, W. N.,**
440 **Chiu, F. C. K., Chollet, J., Craft, J. C., Creek, D. J., Dong, Y., Matile, H., Maurer, M., Morizzi,**
441 **J., Nguyen, T., Papastogiannidis, P., Scheurer, C., Shackleford, D. M., Sriraghavan, K.,**
442 **Stingelin, L., Tang, Y., Urwyler, H., Wang, X., White, K. L., Wittlin, S., Zhou, L. and**
443 **Vennerstrom, J. L. (2011). Synthetic ozonide drug candidate OZ439 offers new hope for a single-**
444 **dose cure of uncomplicated malaria. *Proceedings of the National Academy of Sciences* **108**, 4400-**
445 **4405.**
- 446 **Claessens, A., Adams, Y., Ghumra, A., Lindergard, G., Buchan, C. C., Andisi, C., Bull, P. C.,**
447 **Mok, S., Gupta, A. P., Wang, C. W., Turner, L., Arman, M., Raza, A., Bozdech, Z. and Rowe, J.**
448 **A. (2012). A subset of group A-like var genes encodes the malaria parasite ligands for binding to**
449 **human brain endothelial cells. *Proceedings of the National Academy of Sciences* **109**, E1772–E1781.**

- 450 **Cuvillier, A., Redon, F., Antoine, J. C., Chardin, P., DeVos, T. and Merlin, G.** (2000). LdARL-
451 3A, a *Leishmania promastigote*-specific ADP-ribosylation factor-like protein, is essential for
452 flagellum integrity. *Journal of Cell Science* **113** (Pt 11), 2065-2074.
- 453 **Denny, P. W., Gokool, S., Russell, D. G., Field, M. C. and Smith, D. F.** (2000). Acylation-
454 dependent protein export in *Leishmania*. *Journal of Biological Chemistry* **275**, 11017-11025.
- 455 **Derbyshire, E. R., Prudêncio, M., Mota, M. M. and Clardy, J.** (2012). Liver-stage malaria
456 parasites vulnerable to diverse chemical scaffolds. *Proceedings of the National Academy of Sciences*
457 **109**, 8511-8516.
- 458 **Donaldson, J. G. and Jackson, C. L.** (2011). ARF family G proteins and their regulators: roles in
459 membrane transport, development and disease. *Nature Reviews Molecular Cell Biology* **12**, 362-375.
- 460 **Dondorp, A. M., Nosten, F., Yi, P., Das, D., Physo, A. P., Tarning, J., Lwin, K. M., Ariey, F.,
461 Hanpithakpong, W., Lee, S. J., Ringwald, P., Silamut, K., Imwong, M., Chotivanich, K., Lim, P.,
462 Herdman, T., An, S. S., Yeung, S., Singhasivanon, P., Day, N. P. J., Lindegardh, N., Socheat, D.
463 and White, N. J.** (2009). Artemisinin Resistance in *Plasmodium falciparum* Malaria. *New England*
464 *Journal of Medicine* **361**, 455-467.
- 465 **Farazi, T. A., Manchester, J. K., Waksman, G. and Gordon, J. I.** (2001a). Pre-steady-state kinetic
466 studies of *Saccharomyces cerevisiae* myristoylCoA:protein *N*-myristoyltransferase mutants identify
467 residues involved in catalysis. *Biochemistry* **40**, 9177-9186.
- 468 **Farazi, T. A., Waksman, G. and Gordon, J. I.** (2001b). Structures of *Saccharomyces cerevisiae* *N*-
469 myristoyltransferase with bound myristoylCoA and peptide provide insights about substrate
470 recognition and catalysis. *Biochemistry* **40**, 6335-6343.
- 471 **Fidock, D. A.** (2010). Drug discovery: Priming the antimalarial pipeline. *Nature* **465**, 297-298.
- 472 **Frearson, J. A., Brand, S., McElroy, S. P., Cleghorn, L. A., Smid, O., Stojanovski, L., Price, H.
473 P., Guther, M. L., Torrie, L. S., Robinson, D. A., Hallyburton, I., Mpamhanga, C. P.,
474 Brannigan, J. A., Wilkinson, A. J., Hodgkinson, M., Hui, R., Qiu, W., Raimi, O. G., van Aalten,
475 D. M., Brenk, R., Gilbert, I. H., Read, K. D., Fairlamb, A. H., Ferguson, M. A., Smith, D. F. and
476 Wyatt, P. G.** (2010). *N*-myristoyltransferase inhibitors as new leads to treat sleeping sickness. *Nature*
477 **464**, 728-732.

- 478 **Frenal, K., Polonais, V., Marq, J. B., Stratmann, R., Limenitakis, J. and Soldati-Favre, D.**
479 (2010). Functional dissection of the apicomplexan glideosome molecular architecture. *Cell Host &*
480 *Microbe* **8**, 343-357.
- 481 **Gamo, F.-J., Sanz, L. M., Vidal, J., de Cozar, C., Alvarez, E., Lavandera, J.-L., Vanderwall, D.**
482 **E., Green, D. V. S., Kumar, V., Hasan, S., Brown, J. R., Peishoff, C. E., Cardon, L. R. and**
483 **Garcia-Bustos, J. F.** (2010). Thousands of chemical starting points for antimalarial lead
484 identification. *Nature* **465**, 305-310.
- 485 **Gelb, M. H., Van Voorhis, W. C., Buckner, F. S., Yokoyama, K., Eastman, R., Carpenter, E. P.,**
486 **Panethymitaki, C., Brown, K. A. and Smith, D. F.** (2003). Protein farnesyl and *N*-myristoyl
487 transferases: piggy-back medicinal chemistry targets for the development of antitrypanosomatid and
488 antimalarial therapeutics. *Molecular and Biochemical Parasitology* **126**, 155-163.
- 489 **Georgopapadakou, N. H.** (2002). Antifungals targeted to protein modification: focus on protein *N*-
490 myristoyltransferase. *Expert Opin Investig Drugs* **11**, 1117-1125.
- 491 **Godsel, L. M. and Engman, D. M.** (1999). Flagellar protein localization mediated by a calcium-
492 myristoyl/palmitoyl switch mechanism. *EMBO Journal* **18**, 2057-2065.
- 493 **Goldberg, J.** (1998). Structural basis for activation of ARF GTPase: mechanisms of guanine
494 nucleotide exchange and GTP-myristoyl switching. *Cell* **95**, 237-248.
- 495 **Goncalves, V., Brannigan, J. A., Thinon, E., Olaleye, T. O., Serwa, R., Lanzarone, S.,**
496 **Wilkinson, A. J., Tate, E. W. and Leatherbarrow, R. J.** (2012a). A fluorescence-based assay for *N*-
497 myristoyltransferase activity. *Analytical Biochemistry* **421**, 342-344.
- 498 **Goncalves, V., Brannigan, J. A., Whalley, D., Ansell, K. H., Saxty, B., Holder, A. A., Wilkinson,**
499 **A. J., Tate, E. W. and Leatherbarrow, R. J.** (2012b). Discovery of *Plasmodium vivax N*-
500 Myristoyltransferase Inhibitors: Screening, Synthesis, and Structural Characterization of their Binding
501 Mode. *Journal of Medicinal Chemistry* **55**, 3578-3582.
- 502 **Guiguemde, W. A., Shelat, A. A., Bouck, D., Duffy, S., Crowther, G. J., Davis, P. H., Smithson,**
503 **D. C., Connelly, M., Clark, J., Zhu, F., Jiménez-Díaz, M. B., Martinez, M. S., Wilson, E. B.,**
504 **Tripathi, A. K., Gut, J., Sharlow, E. R., Bathurst, I., Mazouni, F. E., Fowble, J. W., Forquer, I.,**
505 **McGinley, P. L., Castro, S., Angulo-Barturen, I., Ferrer, S., Rosenthal, P. J., DeRisi, J. L.,**

- 506 Sullivan, D. J., Lazo, J. S., Roos, D. S., Riscoe, M. K., Phillips, M. A., Rathod, P. K., Van
507 Voorhis, W. C., Avery, V. M. and Guy, R. K. (2010). Chemical genetics of *Plasmodium*
508 *falciparum*. *Nature* **465**, 311-315.
- 509 Gunaratne, R. S., Sajid, M., Ling, I. T., Tripathi, R., Pachebat, J. A. and Holder, A. A. (2000).
510 Characterization of *N*-myristoyltransferase from *Plasmodium falciparum*. *Biochemical Journal* **348 Pt**
511 **2**, 459-463.
- 512 Hang, H. C., Wilson, J. P. and Charron, G. (2011). Bioorthogonal chemical reporters for analyzing
513 protein lipidation and lipid trafficking. *Accounts of Chemical Research* **44**, 699-708.
- 514 Heal, W. P. and Tate, E. W. (2010). Getting a chemical handle on protein post-translational
515 modification. *Organic & Biomolecular Chemistry* **8**, 731-738.
- 516 Hertz-Fowler, C., Ersfeld, K. and Gull, K. (2001). CAP5.5, a life-cycle-regulated, cytoskeleton-
517 associated protein is a member of a novel family of calpain-related proteins in *Trypanosoma brucei*.
518 *Molecular and Biochemical Parasitology* **116**, 25-34.
- 519 Hopkins, A. L., Groom, C. R. and Alex, A. (2004). Ligand efficiency: a useful metric for lead
520 selection. *Drug Discovery Today* **9**, 430-431.
- 521 Jones, M. L., Collins, M. O., Goulding, D., Choudhary, J. S. and Rayner, J. C. (2012). Analysis
522 of protein palmitoylation reveals a pervasive role in *Plasmodium* development and pathogenesis. *Cell*
523 *Host & Microbe* **12**, 246-258.
- 524 Jones, M. L., Kitson, E. L. and Rayner, J. C. (2006). *Plasmodium falciparum* erythrocyte invasion:
525 a conserved myosin associated complex. *Molecular and Biochemical Parasitology* **147**, 74-84.
- 526 Kappe, S. H. I., Vaughan, A. M., Boddey, J. A. and Cowman, A. F. (2010). That Was Then But
527 This Is Now: Malaria Research in the Time of an Eradication Agenda. *Science* **328**, 862-866.
- 528 Kato, N., Sakata, T., Breton, G., Le Roch, K. G., Nagle, A., Andersen, C., Bursulaya, B.,
529 Henson, K., Johnson, J., Kumar, K. A., Marr, F., Mason, D., McNamara, C., Plouffe, D.,
530 Ramachandran, V., Spooner, M., Tuntland, T., Zhou, Y., Peters, E. C., Chatterjee, A., Schultz,
531 P. G., Ward, G. E., Gray, N., Harper, J. and Winzeler, E. A. (2008). Gene expression signatures
532 and small-molecule compounds link a protein kinase to *Plasmodium falciparum* motility. *Nature*
533 *Chemical Biology* **4**, 347-356.

- 534 **Krafts, K., Hempelmann, E. and Skórska-Stania, A.** (2012). From methylene blue to chloroquine:
535 a brief review of the development of an antimalarial therapy. *Parasitology Research* **111**, 1-6.
- 536 **Leber, W., Skippen, A., Fivelman, Q. L., Bowyer, P. W., Cockcroft, S. and Baker, D. A.** (2009).
537 A unique phosphatidylinositol 4-phosphate 5-kinase is activated by ADP-ribosylation factor in
538 *Plasmodium falciparum*. *International Journal of Parasitology* **39**, 645-653.
- 539 **Li, J. and Zhou, B.** (2010). Biological Actions of Artemisinin: Insights from Medicinal Chemistry
540 Studies. *Molecules* **15**, 1378-1397.
- 541 **Lodge, J. K., Johnson, R. L., Weinberg, R. A. and Gordon, J. I.** (1994). Comparison of myristoyl-
542 CoA:protein *N*-myristoyltransferases from three pathogenic fungi: *Cryptococcus neoformans*,
543 *Histoplasma capsulatum*, and *Candida albicans*. *Journal of Biological Chemistry* **269**, 2996-3009.
- 544 **Loeb, F., Clark, W. M., Coatney, G. R., Coggeshall, L. T., Dieuaide, F. R., Dochez, A. R.,**
545 **Hakansson, E. G., Marshall, E. K., Marvel, C. S., McCoy, O. R., Saper, J. J., Sebrell, W. H.,**
546 **Shannon, J. A. and Carden, G. A.** (1946). ACTIVITY OF A NEW ANTIMALARIAL AGENT,
547 CHLOROQUINE (SN 7618) Statement Approved by the Board for Coordination of Malarial Studies.
548 *Journal of the American Medical Association* **130**, 1069-1070.
- 549 **Mackinnon, M. J. and Marsh, K.** (2010). The Selection Landscape of Malaria Parasites. *Science*
550 **328**, 866-871.
- 551 **Maroof, A., Brown, N., Smith, B., Hodgkinson, M. R., Maxwell, A., Losch, F. O., Fritz, U.,**
552 **Walden, P., Lacey, C. N. J., Smith, D. F., Aebischer, T. and Kaye, P. M.** (2012). Therapeutic
553 Vaccination With Recombinant Adenovirus Reduces Splenic Parasite Burden in Experimental
554 Visceral Leishmaniasis. *Journal of Infectious Diseases* **205**, 853-863.
- 555 **Martinez, A., Traverso, J. A., Valot, B., Ferro, M., Espagne, C., Ephritikhine, G., Zivy, M.,**
556 **Giglione, C. and Meinel, T.** (2008). Extent of N-terminal modifications in cytosolic proteins from
557 eukaryotes. *Proteomics* **8**, 2809-2831.
- 558 **Masubuchi, M., Kawasaki, K., Ebiike, H., Ikeda, Y., Tsujii, S., Sogabe, S., Fujii, T., Sakata, K.,**
559 **Shiratori, Y., Aoki, Y., Ohtsuka, T. and Shimma, N.** (2001). Design and synthesis of novel
560 benzofurans as a new class of antifungal agents targeting fungal *N*-myristoyltransferase. Part 1.
561 *Bioorganic & Medicinal Chemistry Letters* **11**, 1833-1837.

- 562 **Maurer-Stroh, S., Eisenhaber, B. and Eisenhaber, F.** (2002a). N-terminal *N*-myristoylation of
563 proteins: prediction of substrate proteins from amino acid sequence. *Journal of Molecular Biology*
564 **317**, 541-557.
- 565 **Maurer-Stroh, S., Eisenhaber, B. and Eisenhaber, F.** (2002b). N-terminal *N*-myristoylation of
566 proteins: refinement of the sequence motif and its taxon-specific differences. *Journal of Molecular*
567 *Biology* **317**, 523-540.
- 568 **Meister, S., Plouffe, D. M., Kuhen, K. L., Bonamy, G. M. C., Wu, T., Barnes, S. W., Bopp, S. E.,**
569 **Borboa, R., Bright, A. T., Che, J., Cohen, S., Dharia, N. V., Gagaring, K., Gettayacamin, M.,**
570 **Gordon, P., Groessl, T., Kato, N., Lee, M. C. S., McNamara, C. W., Fidock, D. A., Nagle, A.,**
571 **Nam, T.-g., Richmond, W., Roland, J., Rottmann, M., Zhou, B., Froissard, P., Glynne, R. J.,**
572 **Mazier, D., Sattabongkot, J., Schultz, P. G., Tuntland, T., Walker, J. R., Zhou, Y., Chatterjee,**
573 **A., Diagana, T. T. and Winzeler, E. A.** (2011). Imaging of *Plasmodium* Liver Stages to Drive Next-
574 Generation Antimalarial Drug Discovery. *Science* **334**, 1372-1377.
- 575 **Miller, Louis H. and Su, X.** (2011). Artemisinin: Discovery from the Chinese Herbal Garden. *Cell*
576 **146**, 855-858.
- 577 **Mills, E., Price, H. P., Johner, A., Emerson, J. E. and Smith, D. F.** (2007). Kinetoplastid PPEF
578 phosphatases: dual acylated proteins expressed in the endomembrane system of *Leishmania*.
579 *Molecular and Biochemical Parasitology* **152**, 22-34.
- 580 **Moskes, C., Burghaus, P. A., Wernli, B., Sauder, U., Durrenberger, M. and Kappes, B.** (2004).
581 Export of *Plasmodium falciparum* calcium-dependent protein kinase 1 to the parasitophorous vacuole
582 is dependent on three N-terminal membrane anchor motifs. *Molecular Microbiology* **54**, 676-691.
- 583 **Murray, C. J. L., Rosenfeld, L. C., Lim, S. S., Andrews, K. G., Foreman, K. J., Haring, D.,**
584 **Fullman, N., Naghavi, M., Lozano, R. and Lopez, A. D.** (2012). Global malaria mortality between
585 1980 and 2010: a systematic analysis. *Lancet* **379**, 413-431.
- 586 **Ntwasa, M., Egerton, M. and Gay, N. J.** (1997). Sequence and expression of *Drosophila* myristoyl-
587 CoA: protein *N*-myristoyl transferase: evidence for proteolytic processing and membrane localisation.
588 *Journal of Cell Science* **110 (Pt 2)**, 149-156.

- 589 **O'Neill, P. M., Amewu, R. K., Nixon, G. L., Bousejra ElGarah, F., Mungthin, M., Chadwick, J.,**
590 **Shone, A. E., Vivas, L., Lander, H., Barton, V., Muangnoicharoen, S., Bray, P. G., Davies, J.,**
591 **Park, B. K., Wittlin, S., Brun, R., Preschel, M., Zhang, K. and Ward, S. A. (2010). Identification**
592 **of a 1,2,4,5-Tetraoxane Antimalarial Drug-Development Candidate (RKA□182) with Superior**
593 **Properties to the Semisynthetic Artemisinins. *Angewandte Chemie International Edition* **49**, 5693-**
594 **5697.**
- 595 **O'Neill, P. M. and Posner, G. H. (2004). A Medicinal Chemistry Perspective on Artemisinin and**
596 **Related Endoperoxides. *Journal of Medicinal Chemistry* **47**, 2945-2964.**
- 597 **Ohtsuka, T., Aoki, I (2003). *N*-Myristoyltransferase inhibitors as potential antifungal drugs. *Drugs of***
598 ***the Future* **28**, 143-152.**
- 599 **Olego-Fernandez, S., Vaughan, S., Shaw, M. K., Gull, K. and Ginger, M. L. (2009). Cell**
600 **morphogenesis of *Trypanosoma brucei* requires the paralogous, differentially expressed calpain-**
601 **related proteins CAP5.5 and CAP5.5V. *Protist* **160**, 576-590.**
- 602 **Painter, H. J., Morrissey, J. M., Mather, M. W. and Vaidya, A. B. (2007). Specific role of**
603 **mitochondrial electron transport in blood-stage *Plasmodium falciparum*. *Nature* **446**, 88-91.**
- 604 **Panethymitaki, C., Bowyer, P. W., Price, H. P., Leatherbarrow, R. J., Brown, K. A. and Smith,**
605 **D. F. (2006). Characterization and selective inhibition of myristoyl-CoA:protein *N*-**
606 **myristoyltransferase from *Trypanosoma brucei* and *Leishmania major*. *Biochemical Journal* **396**,**
607 **277-285.**
- 608 **Phyo, A. P., Nkhoma, S., Stepniewska, K., Ashley, E. A., Nair, S., McGready, R., ler Moo, C.,**
609 **Al-Saai, S., Dondorp, A. M., Lwin, K. M., Singhasivanon, P., Day, N. P., White, N. J., Anderson,**
610 **T. J. and Nosten, F. (2012). Emergence of artemisinin-resistant malaria on the western border of**
611 **Thailand: a longitudinal study. *Lancet* **379**, 1960-1966.**
- 612 **Pino, P., Sebastian, S., Kim, EunBin A., Bush, E., Brochet, M., Volkmann, K., Kozlowski, E.,**
613 **Llinás, M., Billker, O. and Soldati-Favre, D. (2012). A Tetracycline-Repressible Transactivator**
614 **System to Study Essential Genes in Malaria Parasites. *Cell Host & Microbe* **12**, 824-834.**
- 615 **Plouffe, D., Brinker, A., McNamara, C., Henson, K., Kato, N., Kuhen, K., Nagle, A., Adrián, F.,**
616 **Matzen, J. T., Anderson, P., Nam, T.-g., Gray, N. S., Chatterjee, A., Janes, J., Yan, S. F.,**

- 617 **Trager, R., Caldwell, J. S., Schultz, P. G., Zhou, Y. and Winzeler, E. A.** (2008). *In silico* activity
618 profiling reveals the mechanism of action of antimalarials discovered in a high-throughput screen.
619 *Proceedings of the National Academy of Sciences* **105**, 9059-9064.
- 620 **Price, H. P., Guther, M. L., Ferguson, M. A. and Smith, D. F.** (2010). Myristoyl-CoA:protein *N*-
621 myristoyltransferase depletion in trypanosomes causes avirulence and endocytic defects. *Molecular*
622 *and Biochemical Parasitology* **169**, 55-58.
- 623 **Price, H. P., Hodgkinson, M. R., Wright, M. H., Tate, E. W., Smith, B. A., Carrington, M.,**
624 **Stark, M. and Smith, D. F.** (2012). A role for the vesicle-associated tubulin binding protein ARL6
625 (BBS3) in flagellum extension in *Trypanosoma brucei*. *Biochimica et Biophysica Acta* **1823**, 1178-
626 1191.
- 627 **Price, H. P., Menon, M. R., Panethymitaki, C., Goulding, D., McKean, P. G. and Smith, D. F.**
628 (2003). Myristoyl-CoA:protein *N*-myristoyltransferase, an essential enzyme and potential drug target
629 in kinetoplastid parasites. *Journal of Biological Chemistry* **278**, 7206-7214.
- 630 **Price, H. P., Panethymitaki, C., Goulding, D. and Smith, D. F.** (2005). Functional analysis of
631 TbARL1, an *N*-myristoylated Golgi protein essential for viability in bloodstream trypanosomes.
632 *Journal of Cell Science* **118**, 831-841.
- 633 **Price, H. P., Stark, M. and Smith, D. F.** (2007). *Trypanosoma brucei* ARF1 plays a central role in
634 endocytosis and golgi-lysosome trafficking. *Molecular Biology of the Cell* **18**, 864-873.
- 635 **Price, R. N., Douglas, N. M. and Anstey, N. M.** (2009). New developments in *Plasmodium vivax*
636 malaria: severe disease and the rise of chloroquine resistance. *Current Opinion in Infectious Diseases*
637 **22**, 430-435
- 638 **Proto, W. R., Castanys-Munoz, E., Black, A., Tetley, L., Moss, C. X., Juliano, L., Coombs, G. H.**
639 **and Mottram, J. C.** (2011). *Trypanosoma brucei* metacaspase 4 is a pseudopeptidase and a virulence
640 factor. *Journal of Biological Chemistry* **286**, 39914-39925.
- 641 **Rackham, M. D., Brannigan, J. A., Moss, D. K., Yu, Z., Wilkinson, A. J., Holder, A. A., Tate, E.**
642 **W. and Leatherbarrow, R. J.** (2012). Discovery of Novel and Ligand-Efficient Inhibitors of
643 *Plasmodium falciparum* and *Plasmodium vivax* *N*-Myristoyltransferase. *Journal of Medicinal*
644 *Chemistry Article ASAP*, DOI: 10.1021/jm301474t.

- 645 **Rahlfs, S., Koncarevic, S., Iozef, R., Mailu, B. M., Savvides, S. N., Schirmer, R. H. and Becker,**
646 **K. (2009).** Myristoylated adenylate kinase-2 of *Plasmodium falciparum* forms a heterodimer with
647 myristoyltransferase. *Molecular and Biochemical Parasitology* **163**, 77-84.
- 648 **Rees-Channer, R. R., Martin, S. R., Green, J. L., Bowyer, P. W., Grainger, M., Molloy, J. E. and**
649 **Holder, A. A. (2006).** Dual acylation of the 45 kDa gliding-associated protein (GAP45) in
650 *Plasmodium falciparum* merozoites. *Molecular and Biochemical Parasitology* **149**, 113-116.
- 651 **Resh, M. D. (1994).** Myristylation and palmitoylation of Src family members: The fats of the matter.
652 *Cell* **76**, 411-413.
- 653 **Resh, M. D. (2006a).** Trafficking and signaling by fatty-acylated and prenylated proteins. *Nature*
654 *Chemical Biology* **2**, 584-590.
- 655 **Resh, M. D. (2006b).** Use of analogs and inhibitors to study the functional significance of protein
656 palmitoylation. *Methods* **40**, 191-197.
- 657 **Ridzuan, M. A., Moon, R. W., Knuepfer, E., Black, S., Holder, A. A. and Green, J. L. (2012).**
658 Subcellular location, phosphorylation and assembly into the motor complex of GAP45 during
659 *Plasmodium falciparum* schizont development. *PLoS One* **7**, e33845.
- 660 **Rodrigues, T., Prudencio, M., Moreira, R., Mota, M. M. and Lopes, F. (2012).** Targeting the liver
661 stage of malaria parasites: a yet unmet goal. *Journal of Medicinal Chemistry* **55**, 995-1012.
- 662 **Roth, A. F., Wan, J., Green, W. N., Yates, J. R. and Davis, N. G. (2006).** Proteomic identification
663 of palmitoylated proteins. *Methods* **40**, 135-142.
- 664 **Rottmann, M., McNamara, C., Yeung, B. K. S., Lee, M. C. S., Zou, B., Russell, B., Seitz, P.,**
665 **Plouffe, D. M., Dharia, N. V., Tan, J., Cohen, S. B., Spencer, K. R., Gonzalez-Paez, G. E.,**
666 **Lakshminarayana, S. B., Goh, A., Suwanarusk, R., Jegla, T., Schmitt, E. K., Beck, H.-P., Brun,**
667 **R., Nosten, F., Renia, L., Dartois, V., Keller, T. H., Fidock, D. A., Winzeler, E. A. and Diagana,**
668 **T. T. (2010).** Spiroindolones, a Potent Compound Class for the Treatment of Malaria. *Science* **329**,
669 1175-1180.
- 670 **Rudnick, D. A., McWherter, C. A., Rocque, W. J., Lennon, P. J., Getman, D. P. and Gordon, J.**
671 **I. (1991).** Kinetic and structural evidence for a sequential ordered Bi Bi mechanism of catalysis by

- 672 *Saccharomyces cerevisiae* myristoyl- CoA:protein *N*-myristoyltransferase. *Journal of Biological*
673 *Chemistry* **266**, 9732-9739.
- 674 **Russo, I., Oksman, A. and Goldberg, D. E.** (2009a). Fatty acid acylation regulates trafficking of the
675 unusual *Plasmodium falciparum* calpain to the nucleolus. *Molecular Microbiology* **72**, 229-245.
- 676 **Russo, I., Oksman, A., Vaupel, B. and Goldberg, D. E.** (2009b). A calpain unique to alveolates is
677 essential in *Plasmodium falciparum* and its knockdown reveals an involvement in pre-S-phase
678 development. *Proceedings of the National Academy of Sciences* **106**, 1554-1559.
- 679 **Sadlova, J., Price, H. P., Smith, B. A., Votypka, J., Volf, P. and Smith, D. F.** (2010). The stage-
680 regulated HASPB and SHERP proteins are essential for differentiation of the protozoan parasite
681 *Leishmania major* in its sand fly vector, *Phlebotomus papatasi*. *Cellular Microbiology* **12**, 1765-1779.
- 682 **Sahin, A., Espiau, B., Tetaud, E., Cuvillier, A., Lartigue, L., Ambit, A., Robinson, D. R. and**
683 **Merlin, G.** (2008). The *leishmania* ARL-1 and Golgi traffic. *PLoS One* **3**, e1620.
- 684 **Sebastian, S., Brochet, M., Collins, Mark O., Schwach, F., Jones, Matthew L., Goulding, D.,**
685 **Rayner, Julian C., Choudhary, Jyoti S. and Billker, O.** (2012). A *Plasmodium* Calcium-Dependent
686 Protein Kinase Controls Zygote Development and Transmission by Translationally Activating
687 Repressed mRNAs. *Cell Host & Microbe* **12**, 9-19.
- 688 **Shaw, B. D., Momany, C. and Momany, M.** (2002). *Aspergillus nidulans* swoF encodes an *N*-
689 myristoyl transferase. *Eukaryotic Cell* **1**, 241-248.
- 690 **Slack, R. D., Jacobine, A. M. and Posner, G. H.** (2012). Antimalarial peroxides: advances in drug
691 discovery and design. *MedChemComm* **3**, 281-297.
- 692 **Sleigh, S. H. and Barton, C. L.** (2010). Repurposing Strategies for Therapeutics. *Pharmaceutical*
693 *Medicine* **24**, 151-159.
- 694 **Struck, N. S., de Souza Dias, S., Langer, C., Marti, M., Pearce, J. A., Cowman, A. F. and**
695 **Gilberger, T. W.** (2005). Re-defining the Golgi complex in *Plasmodium falciparum* using the novel
696 Golgi marker PfGRASP. *Journal of Cell Science* **118**, 5603-5613.
- 697 **Sturm, A., Amino, R., van de Sand, C., Regen, T., Retzlaff, S., Rennenberg, A., Krueger, A.,**
698 **Pollok, J.-M., Menard, R. and Heussler, V. T.** (2006). Manipulation of Host Hepatocytes by the
699 Malaria Parasite for Delivery into Liver Sinusoids. *Science* **313**, 1287-1290.

700 **Tewari, R., Straschil, U., Bateman, A., Bohme, U., Cherevach, I., Gong, P., Pain, A. and Billker,**
701 **O.** (2010). The systematic functional analysis of *Plasmodium* protein kinases identifies essential
702 regulators of mosquito transmission. *Cell Host & Microbe* **8**, 377-387.

703 **Towler, D. A., Adams, S. P., Eubanks, S. R., Towery, D. S., Jackson-Machelski, E., Glaser, L.**
704 **and Gordon, J. I.** (1987). Purification and characterization of yeast myristoyl CoA:protein *N*-
705 myristoyltransferase. *Proceedings of the National Academy of Sciences* **84**, 2708-2712.

706 **Tull, D., Heng, J., Gooley, P. R., Naderer, T. and McConville, M. J.** (2012). Acylation-dependent
707 and-independent membrane targeting and distinct functions of small myristoylated proteins (SMPs) in
708 *Leishmania major*. *International Journal of Parasitology* **42**, 239-247.

709 **Tull, D., Vince, J. E., Callaghan, J. M., Naderer, T., Spurck, T., McFadden, G. I., Currie, G.,**
710 **Ferguson, K., Bacic, A. and McConville, M. J.** (2004). SMP-1, a member of a new family of small
711 myristoylated proteins in kinetoplastid parasites, is targeted to the flagellum membrane in
712 *Leishmania*. *Molecular Biology of the Cell* **15**, 4775-4786.

713 **Tyers, M. and Mann, M.** (2003). From genomics to proteomics. *Nature* **422**, 193-197.

714 **Udomsangpetch, R., Somsri, S., Panichakul, T., Chotivanich, K., Sirichaisinthop, J., Yang, Z.,**
715 **Cui, L. and Sattabongkot, J.** (2007). Short-term *in vitro* culture of field isolates of *Plasmodium*
716 *vivax* using umbilical cord blood. *Parasitology International* **56**, 65-69.

717 **Vennerstrom, J. L., Arbe-Barnes, S., Brun, R., Charman, S. A., Chiu, F. C. K., Chollet, J.,**
718 **Dong, Y., Dorn, A., Hunziker, D., Matile, H., McIntosh, K., Padmanilayam, M., Santo Tomas,**
719 **J., Scheurer, C., Scorneaux, B., Tang, Y., Urwyler, H., Wittlin, S. and Charman, W. N.** (2004).
720 Identification of an antimalarial synthetic trioxolane drug development candidate. *Nature* **430**, 900-
721 904.

722 **Weissbuch, I. and Leiserowitz, L.** (2008). Interplay Between Malaria, Crystalline Hemozoin
723 Formation, and Antimalarial Drug Action and Design. *Chemical Reviews* **108**, 4899-4914.

724 **Weston, S. A., Camble, R., Colls, J., Rosenbrock, G., Taylor, I., Egerton, M., Tucker, A. D.,**
725 **Tunnicliffe, A., Mistry, A., Mancina, F., de la Fortelle, E., Irwin, J., Bricogne, G. and Pauptit, R.**
726 **A.** (1998). Crystal structure of the anti-fungal target *N*-myristoyl transferase. *Nature Structural*
727 *Biology* **5**, 213-221.

- 728 **WHO** (2011). World Malaria Report 2011. World Health Organization: Geneva, Switzerland.
- 729 **Winkelmann, E. and Raether, W.** (1978). Chemotherapeutically Active Nitro-Compounds .4. 5-
730 Nitroimidazoles .3. *Arzneimittel-Forschung/Drug Research* **28-1**, 739-749.
- 731 **Wright, M. H., Heal, W. P., Mann, D. J. and Tate, E. W.** (2010). Protein myristoylation in health
732 and disease. *Journal of Chemical Biology* **3**, 19-35.
- 733 **Wyllie, S., Patterson, S., Stojanovski, L., Simeons, F. R., Norval, S., Kime, R., Read, K. D. and**
734 **Fairlamb, A. H.** (2012). The anti-trypanosome drug fexinidazole shows potential for treating visceral
735 leishmaniasis. *Sci Transl Med* **4**, 119re111.
- 736 **Yu, Z., Brannigan, J. A., Moss, D. K., Brzozowski, A. M., Wilkinson, A. J., Holder, A. A., Tate,**
737 **E. W. and Leatherbarrow, R. J.** (2012). Design and Synthesis of Inhibitors of *Plasmodium*
738 *falciparum* N-myristoyltransferase, a Promising Target for Anti-Malarial Drug Discovery. *Journal of*
739 *Medicinal Chemistry* **55**, 8879-8890.
- 740 **Zha, J., Weiler, S., Oh, K. J., Wei, M. C. and Korsmeyer, S. J.** (2000). Posttranslational N-
741 myristoylation of BID as a molecular switch for targeting mitochondria and apoptosis. *Science* **290**,
742 1761-1765.
- 743
- 744 **FIG. LEGENDS**
- 745 **Fig. 1:** Structures of clinically-relevant antimalarial drugs.
- 746 **Fig. 2:** Structures of clinically-relevant antileishmanial drugs.
- 747 **Fig. 3:** The transfer of myristate from myristoyl-CoA to the N-terminal glycine residue of a target peptide (grey circles) by
748 the enzyme N-myristoyltransferase (NMT).
- 749 **Fig. 4:** A) Ternary co-crystal structure of ScNMT (grey surface) with bound peptide substrate (green) and a non-
750 hydrolysable myristoyl-CoA analogue (cyan) bound in the active site. PDB ID: 1IID, Farazi *et al.* (Farazi *et al.* 2001b). B)
751 Myr-CoA or **NHM** from co-crystal structures with different NMTs; 4A95: NHM in PvNMT, green; 2WUU: NHM in
752 LdNMT, cyan; 2P6E: Myr-CoA in ScNMT, yellow; 1IIC: Myr-CoA in ScNMT, purple; 3IWE: Myr-CoA in HsNMT1,
753 orange; 1IYK: Myr-CoA in CaNMT, blue. Images generated using PyMOL (DeLano Scientific).
- 754 **Fig. 5:** A) Summary of the PfNMT inhibitor series obtained by a “Piggy-Back” approach from fungal NMT inhibitor RO-
755 09-4609. B) Binding mode of **Example 26** (Yu *et al.* 2012) bound to the *Plasmodium vivax* NMT active site. PDB
756 Accession Code: 4B14.

757 **Fig. 6:** Structures of hits obtained by high throughput screening against protozoal NMTs

For Peer Review

Haem Detoxification
Disruptors

Sesquiterpene Lactone
Derivatives

Pyrimidine Biosynthesis
Inhibitors

Mitochondrial Electron
Transport Disruptor

Apicoplast Targeting

Unknown Mechanism

For Peer Review

80x26mm (300 x 300 DPI)

or Peer Review

80x83mm (300 x 300 DPI)

A

B

80x94mm (300 x 300 DPI)

Pv Selective Hits

MRT00057965
PvNMT IC₅₀: 2.9 μM

Example 7 (Goncalves 2012b)
PvNMT IC₅₀: 4.7 μM
Hs1NMT IC₅₀: 19.4 μM

Pf Selective Hits

PF-03665853
PfNMT IC₅₀: 0.91 μM
LdNMT IC₅₀: > 80 μM

PF-02378143
PfNMT IC₅₀: 2.54 μM
LdNMT IC₅₀: 65.4 μM

PF-03431814
PfNMT IC₅₀: 0.48 μM
LdNMT IC₅₀: 72 μM

Ld Selective Hits

PF-00349412
PfNMT IC₅₀: 39.6 μM
LdNMT IC₅₀: 0.48 μM

PF-03402623
PfNMT IC₅₀: 71.9 μM
LdNMT IC₅₀: 0.093 μM

PF-03393842
PfNMT IC₅₀: 9.0 μM
LdNMT IC₅₀: 0.10 μM

166x162mm (300 x 300 DPI)