Title: Formation of the Orientale lunar multi-ring basin

Authors: Brandon C. Johnson^{1*†}, David M. Blair², Gareth S. Collins³, H. Jay Melosh⁴, Andrew M. Freed⁴, G. Jeffrey Taylor⁵, James W. Head⁶, Mark A. Wieczorek⁷, Jeffrey C. Andrews-Hanna⁸, Francis Nimmo⁹, James T. Keane¹⁰, Katarina Miljković^{1‡}, Jason M. Soderblom¹, Maria T. Zuber¹

Affiliations:

¹Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA.

²Massachusetts Institute of Technology Haystack Observatory, Route 40, Westford, MA 01886, USA.

⁴Impacts and Astromaterials Research Centre, Dept. Earth Science and Engineering, Imperial College London, London SW7 2AZ, UK

⁴Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, 550 Stadium Mall Drive, West Lafayette, IN 47907, USA.

⁵Hawai'i Institute of Geophysics and Planetology, University of Hawai'i, Honolulu, HI 96822, USA.

⁶Department of Earth, Environmental and Planetary Sciences, Brown University, 324 Brook Street, Providence, RI 02912, USA.

⁷Institut de Physique du Globe de Paris, Sorbonne Paris Cité, Université Paris Diderot, Case 7071, Lamarck A, 35 rue Hélène Brion, Paris Cedex 13 75205, France.

⁸Southwest Research Institute, Boulder, CO 80302, USA..

⁹Department of Earth and Planetary Sciences, University of California, Santa Cruz, California 95064, USA.

¹⁰Department of Planetary Science, Lunar and Planetary Laboratory, University of Arizona, 1629 E University Blvd., Tucson, AZ 85721, USA.

*Correspondence to: Brandon_Johnson@Brown.edu

† Current address: Department of Earth, Environmental and Planetary Sciences, Brown University, 324 Brook Street, Providence, RI 02912, USA.

‡ Current address: Department of Applied Geology, Curtin University, Perth, WA 6102, Australia.

Abstract: Multi-ring basins, large impact craters characterized by multiple concentric topographic rings, dominate the stratigraphy, tectonics, and crustal structure of the Moon. Using a hydrocode, we simulate the formation of Orientale, producing a subsurface structure consistent with high-resolution gravity data from the Gravity Recovery and Interior Laboratory (GRAIL) spacecraft. The impact produces a ~ 390 -km diameter transient crater that is not preserved due to subsequent gravitational collapse. Our simulations indicate that the flow of warm weak material at depth is crucial to the formation of the basin's outer rings, which are large normal faults that form at different times during the collapse stage. The key parameters controlling ring location and spacing are impactor diameter and lunar thermal gradients.

Main Text: Orientale, the youngest and best-preserved lunar multi-ring basin, exhibits three concentric topographic rings and an Inner Depression (*I*). The Inner Depression is a central topographic low associated with the zone of excavated crust that extends to a radial distance from the basin center, R , of 160 km and is bound by a scarp (2, 3). The Outer Rook ($R \approx 330$ km) and Cordillera ($R \approx 430$ km) rings are topographically consistent with fault scarps (3, 4). High-resolution gravity data from the Gravity Recovery and Interior Laboratory (GRAIL) spacecraft reveal that the Outer Rook and Cordillera rings are associated with offsets at the crust mantle interface and localized zones of crustal thinning (2). The Inner Rook ($R \approx 230$ km) is distinct with a topographic signature similar to the peak rings of smaller basins (3, 5) and is associated with a flattening of relief at the crust mantle boundary (2).

To understand the processes that produced the subsurface structure inferred from gravity (2), we model the formation of Orientale using the two dimensional version of iSALE, a multi-material, multi-rheology, finite difference, shock physics code (6-8). Because iSALE is a continuum

model, faults are manifest as localized regions of high strain rather than as discrete slip planes. Previous models of Orientale-scale impacts (9, 10) showed subtle strain localization in the crust around the crater, hinting at ring fault formation during crater collapse, but were unable to resolve fault offsets (i.e. the amount of slip along the fault) and topographic expression. Below we describe several model improvements that allow us to directly resolve the formation of Orientale's rings and faults with kilometers of offset, where previous basin formation models cannot (9-13). We include a dilatancy model (14), which describes how deformation increases the porosity of geological materials and contributes to shear localization (15). We use a damage model with an exponential dependence on strain (15), which results in more localized deformation in rocks that are already heavily fractured. For structures the size of Orientale the curvature of the Moon's surface is important (16), so we model impacts into a spherical Moon-like target with a realistic central gravity field (17).

We assume a vertical impact of an asteroid made of dunite at a typical lunar impact velocity of 15 km/s (18). Our axisymmetric models have a spatial resolution of 1 km. We vary the impactor diameter, pre-impact crustal thickness and thermal structure, while attempting to match ring locations as well as the crustal thickness, which is derived from GRAIL gravity measurements and topography from the Lunar Orbiter Laser Altimeter (LOLA) on the Lunar Reconnaissance Orbiter (19, 20). Because rock strength decreases as the melting temperature is approached (7), the assumed pre-impact thermal structure of the target body has the most substantial effect on the formation of large impact basins (9-12, 16, 21). Our best-fitting model uses a 64 km diameter impactor, a pre-impact crustal thickness of 52 km, and a linear thermal gradient of 14 K/km from a surface temperature of 300 K. The thermal structure transitions to an adiabat of ~ 0.04

K/km at a temperatures exceeding 1300 K (Fig. S1) and is consistent with the lunar thermal structure ~ 4 Gyr ago (17, 21).

Our model provides insight into the extent and temporal progression of excavation and crustal disruption, including the timing and formation of Orientale's rings (Fig. 1). The impact first produces a bowl-shaped crater with a maximum depth of ~ 180 km; this cavity is unstable and experiences gravitational collapse. Although much of the deepest material collapses back into the crater, the maximum depth of excavation is ~ 55 km (i.e. into the mantle), where we define excavated material as any material on a ballistic trajectory and above the pre-impact surface at the time the crater has its maximum volume. Of the $5.7 \times 10^6 \text{ km}^3$ of ejecta $4.2 \times 10^6 \text{ km}^3$ avoid collapsing into the crater and $1.2 \times 10^6 \text{ km}^3$ are deposited between the basin rim and a distance of one basin diameter ($R=430\text{-}860$ km). These values are consistent with constraints from GRAIL and LOLA (2).

The time when the crater reaches its maximal volume is also used to determine the width of the transient crater ($R \approx 195$ km), which extended to a radial distance between the present-day Inner Depression and the Inner Rook ring, and precedes the formation of any of the observed rings. While older studies have attempted to relate the rim of the transient crater to Orientale's rings (1, 22-25), the relative difference in their time of formation suggests that there is no reason to expect a correlation, and indeed no present day topographic features correspond to the transient crater (2). This implies that for large impact basins the transient crater and impactor size must be deduced not from topography, but from the gravity and crustal thickness signature using empirical relations from numerical models (21).

107
108 As the bowl-shaped transient crater collapses, a central uplift forms, composed of mantle
109 material (Fig. 1A). As the crater continues to collapse, the Outer Rook fault (Fig. 1B) forms and
110 ejecta are emplaced well outside the location where the Cordillera ring eventually forms after
111 further inward collapse (Fig 1C, Fig 2A). This aspect of the simulation is consistent with
112 observations suggesting that the Cordillera formed subsequent to ejecta emplacement (3). Our
113 model predicts that both normal faults cut through the crust and their dip angle, angle with
114 respect to the surface, gradually decreases in the mantle (Fig. 1c, Fig. 2A). As illustrated in Fig
115 2A, each fault has an offset of a few kilometers and has an average dip angle of 50–55° within
116 the crust. Although the fault offsets are still present at late times, as evidenced by the
117 displacement along tracer lines (Fig. 2C), numerical diffusion smooths the localized total plastic
118 strain associated with the faults (Fig. 1C, F).

119
120 The Outer Rook and Cordillera rings appear to be the scarps of large normal faults (e.g. 1, 3).
121 Our models indicate that these faults are the result of inward flow of warm, weak mantle material
122 during collapse of the transient crater. This flow of weaker underlying material (Fig. 1, Fig. S2),
123 pulls the cooler crust along with it, ultimately causing extensional faulting with large offsets far
124 from the transient crater rim. In the best-fitting model, the strongest crustal material prior to the
125 impact has a yield strength of 147 MPa, whilst the weakest mantle material has a yield strength
126 of 63 MPa (Fig. S1). The role of the weak underlying mantle material is consistent with the ring
127 tectonic theory of multi-ring basin formation (24).

After rising as high as 140 km above the surface, the central uplift collapses and produces a ~10 km high topographic bulge of crustal material at $R \approx 175$ km (Fig. 2B). The formation of this bulge drives the development of the Inner Rook ring (Fig. 1D, Fig. 2B). This outcome of the simulation is consistent with earlier suggestions that the formation of the Inner Rook ring is analogous to the formation of peak rings observed in smaller basins (5, 9). A secondary phase of inward collapse modifies the topographic expression of the Inner Rook (Fig. 1 D,E, Fig. 2 B,C), perhaps distinguishing it from peak rings of smaller craters (27). Although the topographic signature of the Inner Rook is inconsistent with a fault scarp (3), the Inner Rook is associated with a local flattening of relief at the crust mantle boundary (2), which is reproduced in our simulations (Fig 2C). Our modeling predicts that the Inner Rook is composed of unmelted, lightly shocked crustal material from 35-40 km depths that is pushed over previously emplaced ejecta (Fig. 2, Fig. S3, Fig. S4). This is consistent with observations indicating that localities composed of nearly pure crystalline plagioclase cluster around the Inner Rook (28) and suggests that the crust around Orientale is nearly pure crystalline plagioclase even at depths exceeding 35 km, consistent with lunar bulk composition estimates (29).

In the hours following the impact, relatively cool (~1000 K) crustal material flows inward toward the basin center and covers the denser impact melt pool, which is dominated by mantle material (Fig. 1F, Fig. S4). Combined with the fact that deeply excavated material collapses into the crater (Fig 1A), this explains how a basin-scale impact that penetrates to mantle depths could contain crustal material at the basin center as implied by crustal thickness models (2,20). Our models indicate that a thin deposit of crustal impact melt (Fig. S5) covers the interior surface of Orientale, consistent with the morphologic and spectral interpretation of the Maunder formation

(30). More oblique impacts, characterized by shallower excavation depths, would produce thicker crustal impact melt units (11).

Three hours after initial impact, the center of the basin is ~ 7 km deeper than currently observed (Fig. S6). A deep post-impact basin is integral to the process that produces positive gravity anomalies over most lunar basins (11, 32). Over the next few hundred million years, the basin center rises to its present position in response to the post-impact stress state (11, 32). Mechanical coupling of the lithosphere in the basin center with the lithosphere associated with the crustal collar (annulus of thickest crust following transient crater collapse) enables the initially sub-isostatic basin center to uplift becoming super-isostatic and producing the ~ 250 mGal (1mGal= 1 milliGalileo = 10^{-5} m s⁻²) free air gravity anomaly over the center of Orientale (11, 32, 33).

Although our models do not result in a scarp bounding the Inner Depression ($R \approx 160$ km), they suggest that this scarp forms after the Inner Rook. Any topographic feature formed interior of $R \approx 200$ km, including the transient crater rim, would be destroyed by collapse of the central uplift (Fig. 1). After crater collapse, the impact melt pool extends from the basin center out to the edge of the Inner Depression (Fig. S7). The Inner Depression is clearly associated with the mantle uplift (2). While we cannot rule out formation of the scarp bounding the Inner Depression within a few hours of the impact, our modeling supports the inference that the Inner Depression formed on a much longer time scale, tectonically as the basin adjusts in response to the post impact stress state (11, 34), aided by cooling and contraction of Orientale's massive melt pool (31, 35).

174 In our models, some mantle material splashes out onto the crust as the central uplift collapses
175 (Fig. 1 C-F, Fig. S4), but this may be due to the limitations of our axisymmetric model. The
176 southwestern quadrant of Orientale exhibits at least four large normal faults instead of two seen
177 in other regions (3). The initial crustal thickness varied over the broad area of the Orientale basin
178 and that, coupled with the effects of an oblique impact, may explain the marked difference
179 between the development of rings in the eastern and western sectors of the basin.

180
181 In addition to forming rings in approximately the correct locations (Fig. 1, Fig. S8B), models
182 with pre-impact crustal thicknesses that range from 48–52 km also reproduce the azimuthally
183 averaged crustal thickness profile derived from GRAIL gravity and LOLA topography (Fig. 3).
184 Our modeling shows that ring location and fault offsets are highly dependent on impactor size
185 and the pre-impact thermal gradient (Fig. S9, Fig. S10). In our best-fitting model, the region
186 where crust is thinned from the pre-impact value extends to $R_{\text{thin}} \approx 200$ km. The parameter R_{thin}
187 also provides an estimate for the radius of the zone of ejecta provenance or excavation cavity
188 (37). In our best-fitting model of Orientale these two metrics agree to within 5%. We favor the
189 model with pre-impact crustal thickness of 52 km because it produces a thicker cap of cool
190 crustal material at the center of the basin (Fig. 3) consistent with gravity observations.

191
192 Inverse models of GRAIL gravity data suggest that the Outer Rook and Cordillera are associated
193 with localized crustal thinning and offsets at the crust mantle interface (2). The crustal structure
194 also reveals offsets at the crust mantle interface that do not correlate with rings and may indicate
195 additional subsurface faults. The faults with small offsets forming on either side of the Cordillera
196 in our simulations may be consistent with these additional faults (Fig 2.). The amount of crustal

thinning and magnitude of crust-mantle relief, however, depends on the assumptions of these inverse models (2). Our models exhibit local minima in the crustal thickness (Fig. 3) and offsets at the crust mantle interface (Fig. 2) associated with the Outer Rook and Cordillera. These results are consistent with normal faults with offsets of a few kilometers cutting through the crust mantle interface. The simple geometry of a normal fault with a dip angle of 50 degrees and offset of 4 km cutting through 50-km-thick crust will create an ~40 km wide region where the crust is thinned by ~3 km. This idealized fault geometry is similar to the modeled Outer Rook ring (Fig. 3, red curve between 300-340 km from basin center). When extension occurs close to a fault (within ~40 km), the zone of crustal thinning is broader (Fig. 2, Fig. S8B, Fig. 3). Inverse models of the crust mantle interface are limited in resolution to a wavelength approximately equal to the crustal thickness. Thus, the predicted broader zone of extension may explain in part why the signature of the Cordillera is more robust than that of the Outer Rook (2). Similar local minima in crustal thickness as observed by GRAIL may reveal ring faults in highly degraded or mare filled basins.

References and Notes:

1. J. W. Head, Orientale multi-ringed basin interior and implications for the petrogenesis of lunar highland samples. *The Moon*. **11**, 327–356 (1974).
2. M. T. Zuber *et al.*, *Science*, this issue
3. A. L. Nahm, T. Öhman, D. A. Kring, Normal faulting origin for the Cordillera and Outer Rook Rings of Orientale Basin, the Moon. *J. Geophys. Res. Planets*. **118**, 190–205 (2013).
4. Y. N. Kattoum, J. C. Andrews-Hanna, Evidence for ring-faults around the Orientale basin on the Moon from gravity. *Icarus*. **226**, 694–707 (2013).
5. D. M. H. Baker *et al.*, The transition from complex crater to peak-ring basin on the Moon: New observations from the Lunar Orbiter Laser Altimeter (LOLA) instrument. *Icarus*. **214**, 377–393 (2011).

- 223 6. A. A. Amsden, H. M. Ruppel, C. W. Hirt, *SALE: A simplified ALE computer program for*
224 *fluid flow at speeds* (Los Alamos National Laboratories Report LA-8095, 1980).
- 225 7. G. S. Collins, H. J. Melosh, B. A. Ivanov, Modeling damage and deformation in impact
226 simulations. *Meteorit. Planet. Sci.* **39**, 217–231 (2004).
- 227 8. K. Wünnemann, G. S. Collins, H. J. Melosh, A strain-based porosity model for use in
228 hydrocode simulations of impacts and implications for transient crater growth in porous
229 targets. *Icarus*. **180**, 514–527 (2006).
- 230 9. R. W. K. Potter, D. A. Kring, G. S. Collins, W. S. Kiefer, P. J. McGovern, Numerical
231 modeling of the formation and structure of the Orientale impact basin. *J. Geophys. Res.*
232 *Planets*. **118**, 963–979 (2013).
- 233 10. R. W. K. Potter, Investigating the onset of multi-ring impact basin formation. *Icarus*. **261**,
234 91–99 (2015).
- 235 11. A. M. Freed *et al.*, The formation of lunar mascon basins from impact to contemporary
236 form. *J. Geophys. Res. Planets*. **119**, 2378–2397 (2014).
- 237 12. K. Miljkovic *et al.*, Asymmetric Distribution of Lunar Impact Basins Caused by
238 Variations in Target Properties. *Science*. **342**, 724–726 (2013).
- 239 13. M.-H. Zhu, K. Wünnemann, R. W. K. Potter, Numerical modeling of the ejecta
240 distribution and formation of the Orientale basin on the Moon. *J. Geophys. Res. Planets*.
241 **120**, 2118–2134 (2015).
- 242 14. G. S. Collins, Numerical simulations of impact crater formation with dilatancy. *J.*
243 *Geophys. Res. Planets*. **119**, 2600–2619 (2014).
- 244 15. L. G. J. Montési, A unified description of localization for application to large-scale
245 tectonics. *J. Geophys. Res.* **107**, 2045 (2002).
- 246 16. B. A. Ivanov, H. J. Melosh, E. Pierazzo, Basin-forming impacts: Reconnaissance
247 modeling. *Geol. Soc. Spec. Pap.*, 29–49 (2010).
- 248 17. More detailed descriptions of these models and methods are available as supplementary
249 materials on *Science Online*.
- 250 18. M. Le Feuvre, M. A. Wieczorek, Nonuniform cratering of the Moon and a revised crater
251 chronology of the inner Solar System. *Icarus*, **214**, 1–20 (2011).
- 252 19. D. E. Smith *et al.*, Summary of the results from the Lunar Orbiter Laser Altimeter after
253 seven years in orbit. *Icarus*, doi: 10.1016/j.icarus.2016.06.006 (2016).
- 254 20. M. A. Wieczorek *et al.*, The Crust of the Moon as Seen by GRAIL. *Science*. **339**, 671–675
255 (2013).

- 256 21. R. W. K. Potter, D. A. Kring, G. S. Collins, W. S. Kiefer, P. J. McGovern, Estimating
257 transient crater size using the crustal annular bulge: Insights from numerical modeling of
258 lunar basin-scale impacts. *Geophys. Res. Lett.*, **39**, L18203 (2012).
- 259 22. M. Laneuville, M. A. Wieczorek, D. Breuer, N. Tosi, Asymmetric thermal evolution of
260 the Moon. *J. Geophys. Res. Planets.* **118**, 1435–1452 (2013).
- 261 23. R. B. Baldwin, The tsunami model of the origin of ring structures concentric with large
262 lunar craters. *Phys. Earth Planet. In.* **6**, 327–339 (1972).
- 263 24. H. J. Melosh, W. B. McKinnon, The mechanics of ringed basin formation. *Geophys. Res.*
264 *Lett.* **5**, 985–988 (1978).
- 265 25. J. W. Head, Transition from complex craters to multi-ringed basins on terrestrial planetary
266 bodies: Scale-dependent role of the expanding melt cavity and progressive interaction
267 with the displaced zone. *Geophys. Res. Lett.* **37**, L02203–2009gl041790 (2010).
- 268 26. C. A. Hodges, D. E. Wilhelms, Formation of lunar basin rings. *Icarus.* **34**, 294–323
269 (1978).
- 270 27. G. S. Collins, H. J. Melosh, J. V. Morgan, M. R. Warner, Hydrocode Simulations of
271 Chicxulub Crater Collapse and Peak-Ring Formation. *Icarus.* **157**, 24–33 (2002).
- 272 28. L. C. Cheek, K. L. Donaldson Hanna, C. M. Pieters, J. W. Head, J. L. Whitten, The
273 distribution and purity of anorthosite across the Orientale basin: New perspectives from
274 Moon Mineralogy Mapper data. *J. Geophys. Res. Planets.* **118**, 1805–1820 (2013).
- 275 29. G. J. Taylor, M. A. Wieczorek, Lunar bulk chemical composition: a post-Gravity
276 Recovery and Interior Laboratory reassessment. *Philosophical Transactions of the Royal*
277 *Society A: Mathematical, Physical and Engineering Sciences.* **372**, 20130242–20130242
278 (2014).
- 279 30. P. D. Spudis, D. J. P. Martin, G. Kramer, Geology and composition of the Orientale Basin
280 impact melt sheet. *J. Geophys. Res. Planets.* **119**, 19–29 (2014).
- 281 31. W. M. Vaughan, J. W. Head, L. Wilson, P. C. Hess, Geology and petrology of enormous
282 volumes of impact melt on the Moon: A case study of the Orientale basin impact melt sea.
283 *Icarus.* **223**, 749–765 (2013).
- 284 32. H. J. Melosh *et al.*, The Origin of Lunar Mascon Basins. *Science.* **340**, 1552–1555 (2013).
- 285 33. M. T. Zuber *et al.*, Gravity Field of the Moon from the Gravity Recovery and Interior
286 Laboratory (GRAIL) Mission. *Science.* **339**, 668–671 (2013).
- 287 34. P. K. Byrne, et al. Deep-seated thrust faults bound the Mare Crisium lunar mascon. *Earth*
288 *Planet. Sci. Lett.* **427**, 183–190 (2015).
- 289 35. S. R. Bratt, S. C. Solomon, J. W. Head, The evolution of impact basins: Cooling,

- 290 subsidence, and thermal stress. *J. Geophys. Res.* **90**, 12415–12433 (1985).
- 291 36. D. Elbeshausen, K. Wünnemann, G. S. Collins, Scaling of oblique impacts in frictional
292 targets: Implications for crater size and formation mechanisms. *Icarus*. **204**, 716–731
293 (2009).
- 294 37. M. A. Wieczorek, R. J. Phillips, Lunar multiring basins and the cratering process. *Icarus*
295 **139**, 246–259 (1999).
- 296 38. E. Pierazzo, A. M. Vickery, H. J. Melosh, A reevaluation of impact melt production.
297 *Icarus*. **127**, 408–423 (1997).
- 298 39. W. Benz, A. Cameron, H. J. Melosh, The origin of the Moon and the single-impact
299 hypothesis III. *Icarus*. **81**, 113–131 (1989).
- 300 40. R. W. K. Potter, G. S. Collins, W. S. Kiefer, P. J. McGovern, D. A. Kring, Constraining
301 the size of the South Pole-Aitken basin impact. *Icarus*. **220**, 730–743 (2012).
- 302 41. T. M. Davison, G. S. Collins, F. J. Ciesla, Numerical modelling of heating in porous
303 planetesimal collisions. *Icarus*. **208**, 468–481 (2010).
- 304 42. S. L. Thompson, ANEOS analytic equations of state for shock physics codes input
305 manual. *Sandia Report SAND89-2951 UC-404* (1990).
- 306 43. T. J. Bowling *et al.*, Antipodal terrains created by the Rheasilvia basin forming impact on
307 asteroid 4 Vesta. *J. Geophys. Res. Planets* (2013), doi:10.1002/jgre.20123.
- 308 44. G. R. Johnson, W. H. Cook, A constitutive model and data for metals subjected to large
309 strains, high strain rates and high temperatures. *7th International Symposium on Ballistics*
310 (1983).
- 311 45. K. Wünnemann, G. S. Collins, G. R. Osinski, Numerical modelling of impact melt
312 production in porous rocks. *Earth Planet. Sci. Lett.* **269**, 530–539 (2008).
- 313 46. H. J. Melosh, Impact cratering: A geologic process. *Oxford University Press* (1989).
- 314 47. L. E. Senft, S. T. Stewart, Dynamic fault weakening and the formation of large impact
315 craters. *Earth Planet. Sci. Lett.* **287**, 471–482 (2009).
- 316 48. H. C. F. C. Hay, G. S. Collins, T. M. Davison, Complex Crater Collapse: A Comparison
317 of the Block and Melosh Models of Acoustic Fluidization. *45th Lunar Planet Sci. Conf.*
318 **45**, 1938 (2014).
- 319 49. D. A. Crawford, P. H. Schultz, A Model of Localized Shear Heating with Implications for
320 the Morphology and Paleomagnetism of Complex Craters. *Large Meteorite Impacts and*
321 *Planetary Evolution V*, #3047 (2013).
- 322 50. T. Kenkmann, G. S. Collins, K. Wünnemann, in *Impact Cratering Processes and*

323 *Products*, G. R. Osinski, E. Pierazzo, Pier, Eds. (2012), pp. 60–75.

- 324 51. W. R. Buck, A. N. B. Poliakov, Abyssal hills formed by stretching oceanic lithosphere.
325 *Nature*. **392**, 272–275 (1998).

- 326 52. J. C. Jaeger, N. G. W. Cook, R. Zimmerman, Fundamentals of Rock Mechanics. *John*
327 *Wiley & Sons* (2009).

- 328 53. Potter, R.W.K., Kring, D.A., and Collins, G.S., Scaling of basin-sized impacts and the
329 influence of target temperature, in Osinski, G.R., and Kring, D.A., eds., Large Meteorite
330 Impacts and Planetary Evolution V: Geological Society of America Special Paper 518,
331 p.99-113, (2015) doi:10.1130/2015.2518(06).

- 332 54. S. T. Stewart, Impact basin formation: The mantle excavation paradox resolved. *42nd*
333 *Lunar Planet Sci. Conf.*, #1633 (2011).

- 334 55. P. Del Gaudio *et al.*, Frictional melting of peridotite and seismic slip. *J. Geophys. Res.*
335 **114**, B06306 (2009).

336 **Acknowledgments:** We thank two anonymous reviewers for their thoughtful reviews. We
337 gratefully acknowledge the developers of iSALE (www.isale-code.de/projects/iSALE),
338 including Kai Wünnemann, Dirk Elbeshausen, and Boris Ivanov. We thank all those
339 involved with the development and operations of the GRAIL spacecraft and the
340 collection and reduction of the GRAIL data. This work was funded by the GRAIL
341 mission, which is supported by the NASA Discovery Program and is performed under
342 contract to the Massachusetts Institute of Technology and the Jet Propulsion Laboratory,
343 California Institute of Technology. At present, iSALE is not fully open source. It is distributed
344 on a case-by-case basis to academic users in the impact community, strictly for non-commercial
345 use. Scientists interested in using or developing iSALE should see [http:// www.isale-](http://www.isale-code.de/redmine/projects/isale/wiki/Terms_of_use)
346 [code.de/redmine/projects/isale/wiki/Terms_of_use](http://www.isale-code.de/redmine/projects/isale/wiki/Terms_of_use) for a description of application requirements.
347 Model output is available upon request from Brandon_Johnson@Brown.edu

348

Fig. 1. Time series for the best-fitting model of Orientale crater collapse and ring formation. Material positions are colored according to their total plastic strain. The origin marks the point of impact. The thin black curves mark material interfaces (e.g. crust-mantle interface). An animation of this figure is available (17).

Fig. 2. A vertically exaggerated time series of the formation of Orientale basin, highlighting the ring faults, formation of the Inner Rook, and material provenance. Thin lines connect tracer particles, which track the motion of a parcel of material, that were initially at equal depths. Before the impact, these lines are parallel with a constant spacing of 2 km. Gray curves represent material from pre-impact depths of 0-10 km, dark blue 10-20 km, magenta 20-30 km, red 30-40 km, blue 40-50 km. The black curves represent mantle material. For clarity, lines are not drawn if the next nearest tracer is more than 50 km away. The upper thick blue and lower thick red curves mark model topography and crust mantle interface, respectively. The thick vertical gray lines mark the distance from the basin center where the present-day rings of Orientale are observed at the surface. Elevation and distance are with respect to a sphere so that unperturbed material appears as a series of horizontal lines. This figure shows results for the best-fitting model.

Fig. 3. Post-impact crustal thickness. The model thickness of the crust is plotted from the best-fitting model (red) and for a model with a pre-impact crustal thickness of 48 km (blue). Azimuthally averaged observed profiles of GRAIL-derived crustal thickness (2) are shown for comparison (global average crustal thickness, $t_{av}=43$ km (black) and $t_{av}=34$ km (grey)). Error bars are 1-sigma deviations from the mean.

Supplementary Materials:

Materials and Methods

Figures S1-S10

Tables S1-S2

References (37-52)