ERRATUM

Erratum to: Risk factors for progression in children and young adults with IgA nephropathy: an analysis of 261 cases from the VALIGA European cohort

Rosanna Coppo¹ · Danilo Lofaro² · Roberta R. Camilla¹ · Shubha Bellur³ · Daniel Cattran⁴ · H. Terence Cook⁵ · Ian S. D. Roberts³ · Licia Peruzzi¹ · Alessandro Amore¹ · Francesco Emma⁶ · Laura Fuiano⁶ · Ulla Berg⁷ · Rezan Topaloglu⁸ · Yelda Bilginer⁸ · Loreto Gesualdo⁹ · Rosaria Polci¹⁰ · Malgorzata Mizerska-Wasiak¹¹ · Yasar Caliskan¹² · Sigrid Lundberg¹³ · Giovanni Cancarini¹⁴ · Colin Geddes¹⁵ · Jack Wetzels¹⁶ · Andrzej Wiecek¹⁷ · Magdalena Durlak¹⁸ · Stefano Cusinato¹⁹ · Cristiana Rollino²⁰ · Milena Maggio²¹ · Manuel Praga²² · Hilde K. Smerud²³ · Vladimir Tesar²⁴ · Dita Maixnerova²⁴ · Jonathan Barratt²⁵ · Teresa Papalia² · Renzo Bonofiglio² · Gianna Mazzucco²⁶ · Costantinos Giannakakis²⁷ · Magnus Soderberg⁷ · Diclehan Orhan⁸ · Anna Maria Di Palma⁹ · Jadwiga Maldyk¹¹ · Yasemin Ozluk¹² · Birgitta Sudelin¹³ · Regina Tardanico¹⁴ · David Kipgen¹⁵ · Eric Steenbergen¹⁶ · Henryk Karkoszka¹⁷ · Agnieszka Perkowska-Ptasinska¹⁸ · Franco Ferrario²⁸ · Eduardo Gutierrez²² · Eva Honsova²⁴

Published online: 4 October 2016
© IPNA 2016

Erratum to: *Pediatr Nephrol* DOI: 10.1007/s00467-016-3469-3

1. The name of the second author, Danilo Lofaro, was rendered incorrectly in the original publication but has now been corrected.
2. Table 3 in the original publication contains an error introduced in typesetting. The corrected table is reproduced here.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00467-016-3469-3>.

✉ Rosanna Coppo
rosanna.coppo@unito.it

¹ Fondazione Ricerca Molinette, Nephrology, Dialysis and Transplantation, Regina Margherita Hospital, Turin, Italy

² Annunziata Hospital, Cosenza, Italy

³ Oxford University Hospitals, Oxford, UK

⁴ Toronto General Hospital, University Health Network, Toronto, Ontario, Canada

⁵ Imperial College, Hammersmith Hospital, London, UK

⁶ Bambino Gesù Hospital, Rome, Italy

⁷ Karolinska University Hospital, Huddinge, Sweden

⁸ Hacettepe University, Ankara, Turkey

⁹ University of Bari and Foggia, Bari, Italy

¹⁰ Belcolle Hospital, Viterbo, Italy

¹¹ University of Warsaw, Warsaw, Poland

¹² University of Istanbul, Istanbul, Turkey

¹³ Karolinska Institutet, Stockholm, Sweden

¹⁴ Spedali Civili University Hospital, Brescia, Italy

¹⁵ Western Infirmary Glasgow, Glasgow, UK

¹⁶ Radboud University, Nijmegen, The Netherlands

¹⁷ Silesian University, Katowice, Poland

¹⁸ Warsaw Medical University, Warsaw, Poland

Table 3 Correlations between pathological features and outcomes in Cohort 1 and sub-Cohort 2 patients

Pathological features included in the MEST-Oxford Classification ^a	Cohort 1 (<i>n</i> = 261)				sub-Cohort 2 (<i>n</i> = 174)			
	Rate of renal function decline		Survival from combined event		Rate of renal function decline		Survival from combined event	
	Univariate (ml/min/1.73 m ² /year)	Multivariate β (SD) ^b	Univariate HR (95 % CI)	Multivariate HR (95 % CI) ^c	Univariate (ml/min/1.73 m ² /year)	Multivariate β (SD) ^b	Univariate HR (95 % CI)	Multivariate HR (95 % CI) ^c
Mesangial proliferation								
M0	−1.42 ± 7.97		1	1	−0.85 ± 7.34		1	1
M1	−2.71 ± 7.32	0.37 (0.96)	3.88 (1.69–8.88)	2.80 (1.21–6.50)	−1.46 ± 8.3	1.40 (1.32)	2.71 (0.79–9.35)	1.60 (0.35–7.27)
<i>p</i>	0.243	0.70	0.001	0.016	0.659	0.293	0.114	0.544
Segmental glomerulosclerosis								
S0	−0.89 ± 5.38		1	1	−0.47 ± 5.07		1	1
S1	−2.60 ± 9.58	−0.29 (0.83)	3.51 (1.30–9.47)	3.48 (1.24–9.76)	−1.68 ± 9.99	−0.23 (1.08)	2.70 (0.79–9.27)	2.45 (0.61–9.88)
<i>p</i>	0.076	0.73	0.013	0.018	0.297	0.831	0.114	0.016
Tubular atrophy/interstitial fibrosis								
T0	−1.59 ± 7.53		1	1	−0.84 ± 7.01		1	1
T1-2	−3.83 ± 10.99	−0.85 (1.75)	5.31 (2.18–12.95)	3.49 (1.39–8.76)	−3.06 ± 13.67	1.81 (2.49)	2.86 (0.61–13.3)	0.55 (0.08–3.88)
<i>p</i>	0.242	0.627	<0.001	0.008	0.349	0.468	0.18	0.545

CI, Confidence interval; HR, hazards ratio; eGFR, Estimated glomerular filtration rate

^a M1, Mesangial score >0.5; S1, any segmental sclerosis; T1-2, tubular atrophy and interstitial fibrosis of >25 %. For more detail, see Inclusion criteria, clinical dataset and definitions section and Table 1 footnote

^b Multivariate linear models are adjusted for age, gender, initial eGFR, follow-up blood pressure and proteinuria

^c Multivariate Cox regression models are adjusted for follow-up blood pressure and proteinuria

¹⁹ Borgomanero Hospital, Borgomanero, Italy

²⁰ San Giovanni Bosco Hospital, Turin, Italy

²¹ Hospital Maggiore di Lodi, Lodi, Italy

²² Hospital 12 de Octubre, Madrid, Spain

²³ University of Uppsala, Uppsala, Sweden

²⁴ General University Hospital, Prague, Czech Republic

²⁵ Leicester General Hospital, Leicester, UK

²⁶ University of Turin, Turin, Italy

²⁷ University of Rome, Rome, Italy

²⁸ Monza Hospital, Monza, Italy