Tractable consideration set structures for assortment optimization and network revenue management

Arne K. Strauss¹ and Kalyan Talluri²

¹Warwick Business School, University of Warwick, Coventry CV4 7AL, United Kingdom.

arne.strauss@wbs.ac.uk

²Imperial College Business School, London SW7 2AZ, United Kingdom.

kalyan.talluri@imperial.ac.uk

7 October 2016

Abstract

Discrete-choice models are widely used to model consumer purchase behavior in assortment optimization and revenue management. In many applications, each customer segment is associated with a consideration set that represents the set of products that customers in this segment consider for purchase. The firm has to make a decision on what assortment to offer at each point in time without the ability to identify the customer's segment. A linear program called the Choice-based Deterministic Linear Program (*CDLP*) has been proposed to determine these offer sets. Unfortunately, its size grows exponentially in the number of products and it is NP-hard to solve when the consideration sets of the segments overlap. The Segment-based Deterministic Concave Program with some additional consistency equalities (*SDCP+*) is an approximation of *CDLP* that provides an upper bound on *CDLP*'s optimal objective value. *SDCP+* can be solved in a fraction of the time required to solve *CDLP* and often achieves the same optimal objective value. This raises the question under what conditions can one guarantee equivalence of *CDLP* and *SDCP+*. In this paper, we obtain a structural result to this end, namely that if the segment consideration sets overlap with a certain tree structure or if they are fully nested, *CDLP* can be equivalently replaced with *SDCP+*. We give a number of examples from the literature where

this tree structure arises naturally in modeling customer behavior.

Keywords: discrete choice models, assortment optimization, network revenue management, consideration sets

Received: May 2015; *accepted:* December 2016 by Bill Cooper after two revisions.

1 Introduction

This paper is concerned with three closely related problems: The *Assortment Optimization* problem is to determine an assortment of products to maximize the expected revenue from a heterogeneous population of customers. This population is modeled as being made up of multiple latent customer segments. *Revenue management (RM)* is dynamic assortment optimization for products that share a common resource. In *Network Revenue Management (NRM)*, products may use inventory from multiple resources (for example, a hotel stay for three nights uses three days of inventory; airline itineraries involving multiple flights use seats on the connecting flight legs). Network interactions arise as the decision to offer a product depends on the future revenues attainable from the sale of the other products that share resources on the network.

Underlying all three problems is a model of how consumers choose a product to purchase. Talluri and van Ryzin (2004) introduced RM based on a discrete-choice model of customer purchases. Discrete-choice models represent purchase probability as a function of available products and customer characteristics. Such modeling was later applied to NRM by Gallego et al. (2004) and Liu and van Ryzin (2008) who formulated a Choice-based Deterministic Linear Program (*CDLP*), the main object¹ of study in this paper. The linear program *CDLP* has an exponential number of variables and is difficult to solve except for a few restricted choice models.

In our model, the customer population consists of multiple segments. Each segment is associated with a subset of products called the consideration set, along with segment-specific parameters of the choice model. Assuming that all products are available, the consideration set represents the set of products that customers in that segment would consider for purchase. Consideration sets play a crucial role in the solvability of the assortment optimization problem: If the segments' consideration sets overlap, the assortment optimization

¹We present our results in the context of NRM as this problem generalizes both RM as well as assortment optimization. Subsequent work on this problem can be found in Bodea et al. (2009), Kunnumkal and Topaloglu (2010), Zhang and Adelman (2009), Meissner and Strauss (2012), Méndez-Díaz et al. (2014) and Kunnumkal (2014).

problem is NP-hard when there are just two segments, even for simple choice models such as the Multinomial-Logit (MNL) (Bront et al., 2009).

Motivated by this intractability of *CDLP*, Talluri (2014) develops a weaker formulation called Segment-based Deterministic Concave Program (*SDCP*), weaker in the sense that its optimal objective value gives an upper bound on the optimal objective value of *CDLP*. The idea is to solve a collection of small subproblems, each corresponding to a segment, with some constraints that loosely link them together. *SDCP* is generally poor in approximating *CDLP* when segment consideration sets overlap, i.e. its optimal objective function value is significantly higher than that of *CDLP*, and it also performs poorly in revenue simulations. To improve this situation, Meissner et al. (2013) propose an extension of the *SDCP* formulation called *SDCP+* (defined in §2.4) that obtains a significantly tighter relaxation of *CDLP* for the case of overlapping consideration sets. In their numerical experiments, *SDCP+* achieves the *CDLP* optimal objective function value in many instances despite being faster to solve by an order of magnitude.

The main contribution of this paper is that we identify two consideration set structures for which *CDLP* is equivalent to *SDCP+*, i.e. a solution to *CDLP* can be obtained by instead solving the simpler problem *SDCP+* with significantly fewer decision variables in a fraction of computation time. We give a number of applications from the literature where such consideration set structures naturally occur in the modeling of customer behavior. Our results depend only on the consideration set structure and not on the structure of the network of resources and, importantly, apply for any general discrete-choice model.

The remainder of the paper is organized as follows: In §2 we introduce the notation, the demand model, the basic dynamic program and the two approximations of the dynamic program, namely the *CDLP* and the *SDCP+*. In §3 we present the main structural results. In §4 we illustrate some applications from the literature where the desired structure of the consideration set is naturally present. In §5 we summarize our conclusions.

2 Models

We introduce the notation in §2.1. In §2.2, we state a dynamic programming formulation of the choice-based NRM problem. Its intractability motivates the formulation of approximations: in §2.3 we define the Choice-

based Deterministic Linear Program (*CDLP*), and in §2.4 the Enhanced Segment-Based Deterministic Concave Program (*SDCP+*).

2.1 Notation

A product is a specification of a price (usually with restrictions such as advance purchase requirements) and a set of resources that the product uses. For instance, a product for a network airline would be the combination of fare class (price and restrictions) and the itinerary (the flight legs of the itinerary); in a hotel network, a product is a multi-night stay for a particular room type at a certain price point.

We define a discrete-time booking horizon that consists of T intervals, indexed by t . The sale process begins at time 0 and all resources perish instantaneously at time T . We make the standard assumption that the time intervals are small enough so that the probability of more than one customer arriving in a time period is negligible.

The underlying network has m resources (indexed by i) and n products (indexed by j), and we refer to the set of all resources as I and the set of all products as J . The resources used by j are represented by a resource-product incidence matrix A , with $a_{ij} = 1$ if product j uses resource i , and $a_{ij} = 0$ otherwise. Columns of A are the 0-1 incidence vectors $\mathbf{A}_j$. We denote the vector of capacities at time t as $\mathbf{c}_t$, so the initial set of capacities at time 0 is $\mathbf{c}_0$.

We assume that there are $\mathcal{L} := \{1, \dots, L\}$ customer segments, each with distinct purchase behavior. In each period, a customer arrives with probability λ and belongs to segment l with probability p_l . We denote $\lambda_l = p_l \lambda$ and assume $\sum_{l \in \mathcal{L}} p_l = 1$, so $\lambda = \sum_{l \in \mathcal{L}} \lambda_l$. We assume time-homogenous arrivals (homogenous in rates and segment mix), but the model and all solution methods in this paper can be clearly extended to the case where rates and mix change by period. Customers in segment l have a *consideration set* $C_l \subseteq J$ of products that they consider to purchase (see Shocker et al. (1991) for a survey on the consideration-set modeling literature).

In each period the firm offers a subset S of its products for sale, called the *offer set*. Given an offer set S , an arriving customer purchases a product j (at the price r_j) in the set S or decides not to purchase any (no-purchase). The no-purchase option is indexed by 0 and is always present for the customer.

A segment- l customer's choice probabilities are not affected by the availability of products $j \in J \setminus C_l$. A segment- l customer purchases $j \in S$ if $j \in S \cap C_l$ with probability $P_j^l(S)$, $S \subseteq J$. These functions are either given by an oracle or by a functional form such as in the MNL model where $P_j^l(S) = v_{lj} / (\sum_{k \in C_l \cap S} v_{lk})$ for a set of "weights" $\mathbf{v}_l \in \mathbb{R}^{|C_l|}$ that capture the attractiveness of the products for each segment l .

Whenever we specify probabilities for a segment l for a given offer set S , we just write it with respect to $S_l = C_l \cap S$ (note that $P_j^l(S) = P_j^l(S_l)$). So when the firm offers set S , it sells $j \in S$ with probability $P_j(S) = \sum_{l \in \mathcal{L}} p_l P_j^l(S_l)$ and makes no sale with probability $P_0(S) = 1 - \sum_{j \in S} P_j(S)$.

We define the vector $\mathbf{P}^l(S) = [P_1^l(S_l), \dots, P_n^l(S_l)]$ (recall the no-purchase option is indexed by 0, so it is not included in this vector). We define the vector $\mathbf{P}(S) = [P_1(S), \dots, P_n(S)]$. Notice that $\mathbf{P}(S) = \sum_{l \in \mathcal{L}} p_l \mathbf{P}^l(S)$. We define the vectors $\mathbf{Q}^l(S) = A\mathbf{P}^l(S)$ and $\mathbf{Q}(S) = A\mathbf{P}(S)$ to denote the expected resource consumption for an offer set S by segment l . Likewise, the expected revenue function for segment l is $R^l(S) = \sum_{j \in S_l} r_j P_j^l(S_l)$ and the expected revenue from a given arrival, $R(S) = \sum_{j \in S} r_j P_j(S)$.

In our notation and demand model we broadly follow Bront et al. (2009) and Liu and van Ryzin (2008).

2.2 Dynamic Program

We describe the stochastic dynamic program to determine the optimal offer set at each point in time. While computationally intractable, it gives a conceptual reference point of the value we are trying to approximate with tractable methods.

Let $V_t(\mathbf{c}_t)$ denote the maximum expected revenue that can be earned over the remaining time horizon $[t, T]$, given remaining capacity $\mathbf{c}_t$ in period t . Let $J(\mathbf{c}_t)$ denote the set of products that *can* be offered given remaining available capacity, i.e., $J(\mathbf{c}_t) := \{j \in J | \mathbf{A}_j \leq \mathbf{c}_t\}$. Then $V_t(\mathbf{c}_t)$ satisfies the Bellman equation

$$V_t(\mathbf{c}_t) = \max_{S \subseteq J(\mathbf{c}_t)} \left\{ \sum_{j \in S} \lambda P_j(S) (r_j + V_{t+1}(\mathbf{c}_t - \mathbf{A}_j)) + (\lambda P_0(S) + 1 - \lambda) V_{t+1}(\mathbf{c}_t) \right\}, \quad \forall t, \forall \mathbf{c}_t, \quad (1)$$

with the boundary condition $V_T(\mathbf{c}_T) = 0$ for all $\mathbf{c}_T$. Let $V^{DP} := V_0(\mathbf{c}_0)$ denote the optimal value of this dynamic program from 0 to T , for the given initial capacity vector $\mathbf{c}_0$. Solving the dynamic program (1) is intractable because the state space explodes even for small problems. Therefore, we are forced to look at

approximations to the dynamic program (1).

2.3 Choice Deterministic Linear Program (*CDLP*)

The Choice-based Deterministic Linear Program (*CDLP*) approximation defined in Gallego et al. (2004) and Liu and van Ryzin (2008) has 2^n decision variables w_S (recall that n denotes the number of products). The decision variables can be interpreted as the amount of time set S is offered:

$$\begin{aligned}
 \max \quad & \sum_{S \subseteq J} \lambda R(S) w_S \\
 \text{s.t.} \quad & \sum_{S \subseteq J} \lambda w_S Q(S) \leq c_0 \\
 (CDLP) \quad & \sum_{S \subseteq J} w_S = T \\
 & 0 \leq w_S, \quad \forall S \subseteq J.
 \end{aligned} \tag{2}$$

That is, we maximize the total expected revenue, subject to the constraint that the total expected capacity consumption on each resource i must be less than or equal to the initially available capacity c_{0i} . The second constraint (2) says that we offer product sets over T time units.

Liu and van Ryzin (2008) show that the optimal objective value of *CDLP* is an upper bound on V^{DP} . They also show that the problem can be solved efficiently by column generation for the MNL model with non-overlapping segment consideration sets. Bront et al. (2009) and Rusmevichientong et al. (2014) investigate this further and show that column generation is NP-hard if the consideration sets for the segments overlap for the MNL choice model with two segments.

2.4 Enhanced Segment-Based Deterministic Concave Program (*SDCP+*)

Talluri (2014) proposed an upper bound on *CDLP* called the Segment-based Deterministic Concave Program (*SDCP*). *SDCP* optimizes the offer set for each segment separately. *SDCP* and *CDLP* have the same objective values when the consideration sets for the different segments are disjoint. We do not elaborate on the *SDCP* formulation as it is not considered further in this paper, but it corresponds to the formulation (*SDCP+*) given below without the constraints (4).

In applications, the segments' consideration sets can overlap in a variety of ways and, as the choice probabilities depend on the offer set, they do not have any structure that we can exploit. We call a set of constraints *valid* for a linear programming approximation of the dynamic program (1) if adding the constraints preserves the property that its optimal objective value still forms an upper bound on V^{DP} . Meissner et al. (2013) develop a set of valid inequalities for *SDCP* called product cuts—the constraints (4) below—that tighten the *SDCP* bound. We call the formulation *SDCP+* in light of the additional constraints (4). Let S_{lm} represent subsets of $C_l \cap C_m$, i.e., subsets in the intersection of the consideration sets of segments l and m . *SDCP+* is:

$$\begin{aligned}
& \max \quad \sum_{l=1}^L \sum_{S_l \subseteq C_l} \lambda_l R^l(S_l) w_{S_l}^l \\
& \text{s.t.} \\
(SDCP+) \quad & \sum_{l=1}^L \mathbf{y}_l \leq \mathbf{c}_0 \\
& \sum_{S_l \subseteq C_l} \lambda_l \mathbf{Q}^l(S_l) w_{S_l}^l \leq \mathbf{y}_l, \quad \forall l \in \mathcal{L} \\
& \sum_{S_l \subseteq C_l} w_{S_l}^l = T, \quad \forall l \in \mathcal{L} \\
& \sum_{\{S_l \subseteq C_l | S_l \supseteq S_{lm}\}} w_{S_l}^l - \sum_{\{S_m \subseteq C_m | S_m \supseteq S_{lm}\}} w_{S_m}^m = 0, \\
& \quad \forall S_{lm} \subseteq C_l \cap C_m, \forall \{l, m\} \subset \mathcal{L} : C_l \cap C_m \neq \emptyset \\
& w_{S_l}^l \geq 0, \quad \forall S_l \subseteq C_l, \forall l \in \mathcal{L}, \\
& \mathbf{y}_l \geq \mathbf{0}, \quad \forall l \in \mathcal{L}.
\end{aligned} \tag{3}$$

$$\tag{4}$$

The vector $\mathbf{y}_l$ represents capacity allocation to segment l subject to total available capacity $\mathbf{c}_0$. We maximize total expected revenue from each segment l subject to several constraints. The first represents that the capacity allocations are limited by the overall available network capacity. The second set enforces that each segment can only consume at most as many resources as have been allocated to it. The third ensures that we offer product sets (possibly the empty set) over the full time horizon. The intuition behind the product cuts (4) is the following: *SDCP+* can be seen as a collection of segment-level implementations of *CDLP* linked via the constraint (3) and tightened via the product cuts (4). For any set $S_{lm} \subset C_l \cap C_m$, the length of time that set S_{lm} is offered to segment l (possibly alongside other products) must be equal to the length of time that it is being offered to segment m (again possibly alongside other products). The numerical

experiments of Meissner et al. (2013) show that generating just a few of these constraints can be sufficient to obtain close approximations to the optimal *CDLP* objective function value.

3 Analysis of *CDLP* and *SDCP+*

We wish to understand when the optimal objective value of *SDCP+* is the same as *CDLP*. To this end, we first develop a simple example to illustrate that there can be a strict gap between the optimal objective values of *CDLP* and *SDCP+*, even if all product cuts are satisfied. The underlying reason for that gap is that there is no solution to *CDLP* that can be projected onto the segment consideration sets so as to coincide with the segment-level optimal solution. When are these two formulations equivalent then? We explore this issue in the remainder of this section following the example.

Example 1 Our example has five products and three segments and their respective consideration sets are shown in Figure 1. For this example we show that there is a gap between the optimal objective values of *SDCP+* and *CDLP* even if we generate all constraints of type (4). Assume $T = 1$, capacity $c = 1$, revenue $r_j = 1$ for all products $j \in J = \{1, 2, 3, 4, 5\}$, $\lambda_l = 1/3$ for all segments $l \in \{A, B, C\}$, and the purchase probabilities defined as follows: $P_j^A(\{1, 2\}) := 0.5$ for $j = 1, 2$, $P_j^A(\{2, 5\}) := 0.5$ for $j = 2, 5$, $P_2^B(\{2\}) := 1$, $P_j^B(\{1, 2, 3\}) := 1/3$ for $j = 1, 2, 3$, $P_4^C(\{4\}) := 1$ and $P_j^C(\{3, 5\}) := 0.5$ for $j = 3, 5$, and 0 for all other sets.

Figure 1: Consideration sets for 3 segments, 5 products

We show that there is a feasible solution to *SDCP+* for this example with objective value 1. A feasible solution to *SDCP+* is given by $y_l = 1/3$ for all segments l , and $w_{\{1,2\}}^A = w_{\{2,5\}}^A = 0.5$, $w_{\{2\}}^B = w_{\{1,2,3\}}^B = 0.5$, $w_{\{4\}}^C = w_{\{3,5\}}^C = 0.5$, and $w_{S_l}^l = 0$ otherwise for all $l \in \{A, B, C\}$, $S_l \subset C_l$. This solution is feasible to *SDCP+* since $\lambda_l \sum_{S_l \subset C_l} Q^l(S_l) w_{S_l}^l = 1/3 = y_l$ for all segments l , and the product cuts (4) for all pairs of segments $\{l, m\}$ and sets $S_{lm} \subset C_l \cap C_m$, $S_{lm} \neq \emptyset$ are satisfied as reported in Table 1. They also hold for $S_{lm} = \emptyset$ since the solution satisfies $\sum_{S_l \subset C_l} w_{S_l}^l = 1$ for all segments l . The other constraints are likewise satisfied as can easily be checked. The objective value is 1 since $\lambda_l \sum_{S_l \subset C_l} R^l(S_l) w_{S_l}^l = 1/3$ for each

Figure 2: Intersection graph for the example in Figure 1.

$l \in \{A, B, C\}$.

Next, we show that there is no corresponding solution to *CDLP* with the same objective value. Note that *CDLP* has $2^5 = 32$ variables corresponding to subsets $S \subset J = \{1, 2, 3, 4, 5\}$. Under the single-leg example described above, we can enumerate all 32 subsets S and calculate the corresponding objective coefficient $\lambda R(S)$. We find that $\lambda R(S) \leq 2/3$ for all $S \subset J$, with equality reached for the sets $\{1, 2, 3\}$, $\{1, 2, 4\}$, $\{2, 3, 5\}$, $\{1, 2, 3, 4\}$ and $\{1, 2, 3, 5\}$. It follows that there can be no feasible solution to *CDLP* that has objective value greater than $2/3$ since the objective is a convex combination of these coefficients (note that $T = 1$).

$\{l, m\}$	S_{lm}	$S_l \supseteq S_{lm}$	$\sum_{S_l \supseteq S_{lm}} w_{S_l}^l$	$S_m \supseteq S_{lm}$	$\sum_{S_m \supseteq S_{lm}} w_{S_m}^m$
$\{A, B\}$	$\{1\}$	$\{1\}, \{1, 2\}, \{1, 5\}, \{1, 2, 5\}$	0.5	$\{1\}, \{1, 2\}, \{1, 3\}, \{1, 2, 3\}$	0.5
$\{A, B\}$	$\{2\}$	$\{2\}, \{1, 2\}, \{2, 5\}, \{1, 2, 5\}$	1	$\{2\}, \{1, 2\}, \{2, 3\}, \{1, 2, 3\}$	1
$\{A, B\}$	$\{1, 2\}$	$\{1, 2\}, \{1, 2, 5\}$	0.5	$\{1, 2\}, \{1, 2, 3\}$	0.5
$\{A, C\}$	$\{5\}$	$\{5\}, \{1, 5\}, \{2, 5\}, \{1, 2, 5\}$	0.5	$\{5\}, \{3, 5\}, \{4, 5\}, \{3, 4, 5\}$	0.5
$\{B, C\}$	$\{3\}$	$\{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$	0.5	$\{3\}, \{3, 4\}, \{3, 5\}, \{3, 4, 5\}$	0.5

Table 1: Evaluation of all product cuts (4) for the example given in Figure 1.

Moving on from the example, we seek to obtain a structural result on when *CDLP* and *SDCP+* are equivalent. Since the overlap of the consideration sets plays a critical role in Example 1, let us represent the overlap structure in a graph. Specifically, we define a bipartite *intersection graph* as follows: There are two types of nodes, one type called segment node, the other is called intersection node. Each node of the former type corresponds to a segment, each of the latter represents a set of the form $C_k \cap C_l$ for some segment pair (k, l) . If there are two pairs of segments (m, n) and (k, l) with $C_m \cap C_n = C_k \cap C_l$, then $S = C_m \cap C_n$ is represented by a single intersection node. Edges from segment node k connect to all the sets of the form $C_k \cap C_l \neq \emptyset$ for any $l \neq k$. In graph theory, a connected graph without cycles is called a tree, and a disjoint union of trees is called a forest (Bondy and Murty, 1976).

The intersection graph of the example of Figure 1 has a cycle, as can be seen from Figure 2. This turns

out to be the critical feature: If the segment consideration sets do not have a cycle and are arranged say in the form of a tree (or, in general, a forest), then the product cuts are sufficient to ensure equivalence between $CDLP$ and $SDCP+$, as stated in Proposition 1 below. Before establishing this result, we provide some intuition for it: The intersection tree tells us which segments are directly or indirectly connected to each other, in the sense that a solution w^l for some segment l is dependent on the solution w^k for any segment node k that is reachable from segment node l . Example 1 illustrates this point: It is not possible to arrange the segment-level solutions in a way such that they are consistent. By consistent, we mean that there is a feasible solution to $CDLP$ that can be projected onto the segments to obtain the $SDCP+$ solution. For instance, if we would arrange the $SDCP+$ solution so that the sets $S_1^A = \{1, 2\}$, $S_1^B = \{1, 2, 3\}$, and $S_1^C = \{3, 5\}$ are offered in parallel (recall that they are all offered for the same duration), then we cannot find a single set $S \subset J$ whose projection onto the segment consideration sets results in S_1^A , S_1^B and S_1^C , respectively. To see this, note that S_1^A requires us *not* to offer product 5, whereas S_1^C *does* require us to offer product 5. In fact, there is no possibility to arrange the segment-level solutions so that all sets that would be offered in parallel are consistent, and this shows that the segment-level solutions of Example 1 do not have a corresponding $CDLP$ solution.

The product cuts ensure that any offer set in the intersection of any two segments' consideration sets is being offered to both segments for the same time (possibly alongside offering other products). This is the case in Example 1: All sets are offered for the same duration. Suppose we have a tree-structured intersection graph with a segment node l connected to a segment node k via a single intersection node. For a given solution w^l , the product cuts allow us to arrange the segment-level solutions w^k in a way such that they represent the projections of a feasible global solution onto the respective segments. We can repeat this argument to construct a global $CDLP$ solution by moving along the tree. However, if there is a cycle, then this pairwise approach to construct a global solution does not work any more because there is no guarantee that the last segment-level solution (w^C in Example 1) in the sequence along the cycle is consistent with the first segment-level solution (w^A in Example 1).

Proposition 1. *If the intersection graph is a forest, then $CDLP$ and $SDCP+$ have the same optimal objective value (henceforth referred to by the shorthand notation $CDLP = SDCP+$).*

Proof

Figure 3: Merging procedure used in the proof of Proposition 1.

Any solution to $CDLP$ is a solution to $SDCP+$ as shown in Meissner et al. (2013), hence $CDLP \leq SDCP+$. It remains to show $CDLP \geq SDCP+$. We use a merging procedure in this proof. For clarity, we first explain it on a simple example with two segments, A and B , displayed in Figure 3 along with their corresponding consideration sets. Figure 3 shows a feasible solution to $SDCP+$. We wish to construct a feasible solution to $CDLP$ from this solution with the same objective value. For simplicity assume that the only sets with positive weights that contain U are S_2 in the consideration set of A and S_3 in the consideration set of B . Note that the product cuts imply the restriction $w_{S_2}^A = w_{S_3}^B$ since S_2 and S_3 are the only offer sets with positive weight in the segment-level solution for segments A and B respectively that contain the set U in the intersection of the consideration sets. Moreover, $w_{S_1}^A + w_{S_2}^A = T$ and $w_{S_3}^B + w_{S_4}^B = T$. This implies that $w_{S_1}^A = w_{S_4}^B$. So we construct weights for the $CDLP$ formulation as $w_{S_1 \cup S_4}^{CDLP} = w_{S_1}^A = w_{S_4}^B$ and $w_{S_2 \cup S_3}^{CDLP} = w_{S_2}^A = w_{S_3}^B$. This $CDLP$ solution satisfies $w_{S_1 \cup S_4}^{CDLP} + w_{S_2 \cup S_3}^{CDLP} = T$, as well as the capacity constraints, and has the same objective value as the $SDCP+$ solution.

In the proof of Proposition 1 (see appendix), we essentially repeat this argument for the more complicated case with L segments and arbitrary consideration sets using an induction argument (made possible by the tree structure).

□

In addition to the tree-structured intersection graphs, we identify another structure that guarantees equivalence of $CDLP$ and $SDCP+$. We show that nested consideration sets also guarantee $CDLP = SDCP+$ even though such consideration sets do not have the tree structure.

Proposition 2. *For a nested consideration set structure $C_1 \subseteq C_2 \subseteq \dots \subseteq C_L$, $CDLP = SDCP+$.*

Proof

See appendix.

□

Let us consider why we do not necessarily need a tree structure in the intersection graph: the main induction step in the proof of Proposition 1 is a merging procedure between a leaf node and the rest of the intersection graph. What we require is that at every step we should be able to find a leaf node. Once we identify the leaf node (segment), and remove it, the intersection graph of the remaining segments can be quite different from the original. Indeed, that is the reason for the tractability of the nested consideration structure in Proposition 2 even though the original intersection graph is not a tree. Hence, we can write a more general version of Proposition 1 as follows:

Proposition 3. *CDLP is equivalent to SDCP+ when the intersection graph has a sequence of segment-nodes, such that the first node in the sequence is a leaf node, and after removal of each leaf node and a re-drawing of the intersection graph with the remaining segments the next segment-node in the sequence is also a leaf node in the new intersection graph.*

4 Applications

In this section we present some applications from the literature where the consideration set structures that we have described appear naturally in the modeling. These applications are: RM of advance tickets and ticket options for sport events (§4.1), RM for primary care clinics (§4.2), dynamic pricing of home delivery time slots (§4.3), low cost airlines (§4.4), and retail (§4.5).

4.1 Revenue management of advanced tickets and options for sports tickets

Sport event ticket options have become so popular that there is a software company called TTR that specializes in selling Internet platforms to teams and events that wish to offer options. Balseiro et al. (2011) consider a scenario where advance tickets for the tournament are sold before it starts—hence the identities of the two teams playing in a tournament final are unknown at this time. However, fans of a specific team are only interested in attending the event if their team makes it to the final. To address this uncertainty, the authors propose team-specific call options under which a customer can pay a small non-refundable amount in advance for the right to attend the event if and only if the specified team makes it to the final and if he pays an additional amount once the finalists are known. Such options allow event organizers in principle

Figure 4: A star consideration structure (left) and the corresponding intersection graph (right) in sports RM.

to oversell capacity many times because only fans with advance tickets or with options for the two finalist teams will be able to attend the event. Fans are segmented by the teams they support; assuming there are L teams in the tournament in total, we therefore have L customer segments. A customer from segment l has the choice between buying an advance ticket A that would give access to the final regardless of who will be playing, and an option O^l for team l . In other words, a customer in segment l has a consideration set $\{A, O^l\}$. Thus we obtain a star consideration set structure (Figure 4).

Balseiro et al. (2011) use the *CDLP* to solve the problem, and show for their specific model that an equivalent, more compact formulation exists that actually is special case of *SDCP+*. The main result of our paper provides a more general explanation for the equivalence of *CDLP* and *SDCP+*, namely that the consideration set structure is a tree.

4.2 Revenue management for primary care clinics

Gupta and Wang (2008) present an application of revenue management under patient choice of primary care providers and appointment time-slots. Specifically, the problem is to manage physicians' consultation time slot availabilities over a finite booking horizon so as to maximize revenues. Each physician has a panel of patients for whom he is the designated primary-care provider; these patient groups correspond to customer segments with preferences for particular physicians. Patients of any segment can choose between all available combinations of all appointment time-slots and all physicians, hence the consideration sets are all identical. Same-day patients form another segment and are assumed to be willing to accept any available slot with any physician on the workday. In this application, the product is a combination of physician-time combination. All patient segments consider all products, and therefore the intersection tree has a star structure as in the previous example. Gupta and Wang (2008) proposed various heuristics to tackle the problem; our main

result tells us that we can use the tractable *SDCP+* formulation in lieu of *CDLP* as an alternative solution approach.

4.3 Dynamic pricing of home delivery time-slots

Another application related to appointment scheduling is the work of Asdemir et al. (2009) who look at the question of how to dynamically price delivery time-slots for attended home delivery over a finite booking horizon using dynamic programming. Different prices can be quoted for different delivery time-slots at any given point in time. There are different customer segments in a given area based on a choice model that reflects their preferences for specific time-slots as well as price sensitivity. Their model considers all geographic areas as independent of each other. All segments in a given area are assumed to consider all available delivery time-slots. If the time slot prices have to be the same across all segments within a given geographical area (which would be reasonable so as to avoid customer dissatisfaction due to perceived unfairness of this group-based discrimination), we again have a star-shaped tree as the intersection graph. Similar to the previous application, the geographic area and time-slot combination is the product.

As solution method, Asdemir et al. (2009) propose a dynamic program that has an exponentially growing state space in the number of delivery time-slots. Alternatively, one could again use *SDCP+* (equivalent to *CDLP* owing to the tree structure) to obtain an approximate policy that can be calculated even for large problem instances where dynamic programming becomes intractable.

4.4 Nested consideration set structure (low-cost airline model)

Consider a fully nested consideration set structure where the L consideration sets are nested as $C_1 \subseteq C_2 \subseteq \dots \subseteq C_L$. This models buy-up/buy-down amongst unrestricted products where complete dilution is possible. This type of structure is encountered on single-leg flights or in a retail context where customers remove products from consideration based on certain cutoff values for the products' attributes or qualities, and these can be ranked linearly. The latter example was proposed by Feldman and Topaloglu (2015) to motivate work on assortment optimization under MNL with nested consideration sets. Proposition 2 shows that this is tractable, following the structure defined in Proposition 3.

Figure 5: Intersection graph for the retail data set in Kamakura and Russell (1989) (left), transformed after merging segment 5 and 3 and after reducing consideration sets to those with purchase probability greater than 10% (middle), and after removing leaf nodes as in Proposition 3 (right).

4.5 Retail

When defining segments and their consideration sets, there is often a certain degree of subjectivity and modeling flexibility. For instance, consider the segmentation study based on retail scanner data by Kamakura and Russell (1989). Initially, they discover nine customer segments with intersection graph as illustrated on the left of Figure 5. There are four loyal segments for the brands A, B, C and P, respectively, and five switching segments. They found that segment 5 and segment 3 could be merged without major impact on the preference structure. Furthermore, one could refine the sets of the switching segments by including only the products with purchase probabilities greater than 10% (the threshold mentioned in Kamakura and Russell (1989)). The resulting simplified consideration sets of the modified segments 1', 2', 3', 4' were $\{A\}$, $\{A,B\}$, $\{A,B,C\}$ and $\{B,C,P\}$, respectively. The corresponding intersection graph is depicted in the middle of Figure 5 along with the simple tree structure on the right that results from removing leaf nodes as in Proposition 3. This serves as an example of how a modeler could sensibly change the segmentation so as to obtain a tractable structure; effectively it comes down to balancing the loss in modeling accuracy with

tractability of the subsequent optimization.

5 Conclusions

Discrete-choice models are widely used to model consumer purchase behavior in assortment optimization and revenue management. The firm has to make a decision on what assortment to offer at each point in time without the ability to identify the customer's segment. In many applications, each customer segment is associated with a consideration set that represents the set of products that customers in this segment consider for purchase. The formulation *CDLP* has been proposed to determine these offer sets but its size grows exponentially in the number of products and it is computationally intractable for even modest-sized applications when segment consideration sets overlap. The formulation *SDCP+* runs much faster than *CDLP* and often obtains the same optimal objective function value. In this paper we show that *CDLP* and *SDCP+* are equivalent if the intersection graph of the segment consideration sets is a tree or if the consideration sets are nested. We give a number of examples from the literature that naturally exhibit these structures.

References

- Kursad Asdemir, Varghese S. Jacob, and Ramayya Krishnan. Dynamic pricing of multiple home delivery options. *European Journal of Operational Research*, 196(1):246–257, 2009.
- S. Balseiro, C. Gocmen, G. Gallego, and R. Phillips. Revenue management of consumer options for sporting events. Tech. rep., IEOR, Columbia University, New York, February 2011.
- T. Bodea, M. Ferguson, and L. Garrow. Choice-based revenue management: Data from a major hotel chain. *Manufacturing & Service Operations Management*, 11(2):356–361, 2009.
- J.A. Bondy and U.S.R. Murty. *Graph Theory with Applications*. Elsevier Science Publishing Co., 1976.
- J. J. M. Bront, I. Méndez-Díaz, and G. Vulcano. A column generation algorithm for choice-based network revenue management. *Operations Research*, 57(3):769–784, 2009.
- J. Feldman and H. Topaloglu. Assortment optimization under the multinomial logit model with nested consideration sets. Working Paper, Cornell University, July 2015.

- G. Gallego, G. Iyengar, R. Phillips, and A. Dubey. Managing flexible products on a network. Technical Report TR-2004-01, Dept of Industrial Engineering, Columbia University, NY, NY, 2004.
- D. Gupta and L. Wang. Revenue management for a primary-care clinic in the presence of patient choice. *Operations Research*, 56(3):576–592, 2008.
- W. A. Kamakura and G. J. Russell. A probabilistic choice model for market segmentation and elasticity structure. *Journal of Marketing Research*, 26:379–390, 1989.
- S. Kunnumkal. Randomization approaches for network revenue management with customer choice behavior. *Production and Operations Management*, 23(9):1617–1633, 2014.
- S. Kunnumkal and H. Topaloglu. A new dynamic programming decomposition method for the network revenue management problem with customer choice behavior. *Production and Operations Management*, 19(5):575–590, 2010.
- Q. Liu and G. van Ryzin. On the choice-based linear programming model for network revenue management. *Manufacturing and Service Operations Management*, 10(2):288–310, Spring 2008.
- J. Meissner and A. K. Strauss. Network revenue management with inventory-sensitive bid prices and customer choice. *European Journal of Operational Research*, 216(2):459–468, 2012.
- J. Meissner, A. K. Strauss, and K. Talluri. An enhanced concave program relaxation for choice network revenue management. *Production and Operations Management*, 22(1):71–87, 2013.
- I. Méndez-Díaz, J. Miranda Bront, G. Vulcano, and P. Zabala. A branch-and-cut algorithm for the latent-class logit assortment problem. *Discrete Applied Mathematics*, 164(1):246–263, 2014.
- P. Rusmevichientong, D. Shmoys, C. Tong, and H. Topaloglu. Assortment optimization under the multinomial logit model with random choice parameters. *Production and Operations Management*, 23(11):2023–2039, September 2014.
- A.D. Shocker, M. Ben-Akiva, B. Boccara, and P. Nedungadi. Consideration set influences on consumer decision-making and choice: Issues, models, and suggestions. *Marketing Letters*, 2(3):181–197, 1991.
- K. T. Talluri and G. J. van Ryzin. Revenue management under a general discrete choice model of consumer behavior. *Management Science*, 50(1):15–33, January 2004.

K.T. Talluri. New formulations for choice network revenue management. *INFORMS Journal of Computing*, 26(2):401–413, 2014.

D. Zhang and D. Adelman. An approximate dynamic programming approach to network revenue management with customer choice. *Transportation Science*, 43(3):381–394, 2009.

Appendix

Proof of Proposition 1

Proof

Any solution to $CDLP$ is a solution to $SDCP+$ as shown in Meissner et al. (2013), hence $CDLP \leq SDCP+$. It remains to show $CDLP \geq SDCP+$.

Consider the case of a single segment $L = 1$, and a given feasible solution $(w_{S_L}^L, y_L)_{S_L}$ to $SDCP+$. Then $w_S^{CDLP} := w_S^L$ for all $S \subset J$ is a feasible solution to $CDLP$ with the same objective value.

Next, we consider $L > 1$. Without loss of generality, the discussion will focus on an intersection graph that is a finite tree rather than a forest since the same arguments can be made for each tree that makes up the forest. Assuming that it is a tree, there must be at least two leaves, i.e. nodes with degree 1. By definition, intersection nodes have at least degree 2, so there exists a segment node that is a leaf. Without loss of generality, let this node correspond to the consideration set of segment L , and let $\overline{SDCP+}$ represent the problem $SDCP+$ with the segment L removed. Consider a feasible solution (w, y) to $SDCP+$, where (w, y) is shorthand notation for $w_{S_l}^l$ for $S_l \subseteq C_l$, for all segments $l \in \mathcal{L} := \{1, 2, \dots, L\}$, and y_{il} for all resources i and $l \in \mathcal{L}$.

This solution induces a feasible solution $(\bar{w}, \bar{y})$ to $\overline{SDCP+}$ by defining $\bar{w}_{S_l}^l := w_{S_l}^l$ for all $S_l \subseteq C_l$, for all $l \in \bar{\mathcal{L}} := \mathcal{L} \setminus \{L\}$, and $\bar{y}_l := y_l$ for all $l \in \bar{\mathcal{L}}$. The solution $(\bar{w}, \bar{y})$ produces an objective value equal to that of $SDCP+$ less $\sum_{S_L \subseteq C_L} \lambda_L R^L(S_L) w_{S_L}^L$. By the induction assumption, there exists a feasible solution $\bar{w}_S^{CDLP}$ for all $S \subseteq \bar{J} := \bigcup_{l=1}^{L-1} C_l$ to $CDLP$ with the same objective value, and $\bar{w}^{CDLP}$ induces $(\bar{w}, \bar{y})$ meaning that $\bar{w}_{S_l}^l = \sum_{S \subseteq \bar{J} | S \cap C_l = S_l} \bar{w}_S^{CDLP}$ for all $l \in \bar{\mathcal{L}}$, $S_l \subseteq C_l$ for $l \in \bar{\mathcal{L}}$, and $\bar{y}_l = \sum_{S_l \subseteq C_l} \lambda_l Q^l(S_l) \bar{w}_{S_l}^l$ for all $l \in \bar{\mathcal{L}}$.

Now we construct a feasible solution w_S^{CDLP} for all $S \subset J$ to $CDLP$ that induces (w, y) for $SDCP+$ with the same objective value. Since L is a leaf of the intersection tree, all interactions with other segments

are via a set S^{int} that is associated with the intersection node to which L is connected. Let us denote all segments that are connected to this intersection node by $\mathcal{L}^{\text{int}}$.

Consider a set $U \subseteq S^{\text{int}}$ that is *maximal* for segment L with respect to S^{int} , that is there is no set $S_L \subseteq C_L$ such that $U \subsetneq S_L \cap S^{\text{int}}$ and positive support $w_{S_L}^L > 0$. Note that for a feasible solution to $SDCP+$, the product cuts ensure that if a set is maximal for L with respect to S^{int} , it is maximal for all segments $l \in \mathcal{L}^{\text{int}}$ with respect to S^{int} . Moreover, from the definition of maximal

$$\sum_{\{S_l \subseteq C_l \mid S_l \cap S^{\text{int}} \supseteq U\}} w_{S_l}^l = \sum_{\{S_l \subseteq C_l \mid S_l \cap S^{\text{int}} = U\}} w_{S_l}^l, \forall l \in \mathcal{L}^{\text{int}}.$$

We select an arbitrary *maximal* set $U \subseteq S^{\text{int}}$ and segment $l \in \mathcal{L}^{\text{int}}$. The following argument shows that the total weight $\tau(U)$ that we offer sets that intersect with S^{int} exactly in U is the same in solutions w^L and $\bar{w}^{CDLP}$:

$$\tau(U) = \sum_{S_L \subseteq C_L \mid S_L \cap S^{\text{int}} = U} w_{S_L}^L \quad (5)$$

$$= \sum_{S_l \subseteq C_l \mid S_l \cap S^{\text{int}} = U} w_{S_l}^l \quad (6)$$

$$= \sum_{S_l \subseteq C_l \mid S_l \cap S^{\text{int}} = U} \bar{w}_{S_l}^l \quad (7)$$

$$= \sum_{S_l \subseteq C_l \mid S_l \cap S^{\text{int}} = U} \sum_{S \subseteq \bar{J} \mid S \cap C_l = S_l} \bar{w}_S^{CDLP} \quad (8)$$

$$= \sum_{S \subseteq \bar{J} \mid S \cap S^{\text{int}} = U} \bar{w}_S^{CDLP}.$$

The first equality holds by definition, the second due to maximality and the product cuts being satisfied by the solution w to $SDCP+$, the third since $w_{S_l}^l = \bar{w}_{S_l}^l$, the fourth because $\bar{w}^{CDLP}$ induces $\bar{w}$, and the final one as a result of a reformulation.

As a consequence, we can merge the solution $w_{S_l}^L$ with $\bar{w}^{CDLP}$ over total weight $\tau(U)$ to obtain w^{CDLP} for all sets that intersect with S^{int} only in the fixed set U . We illustrate the process in Figure 6 by drawing two parallel bars of equal length representing the weight $\tau(U)$, each bar with intervals corresponding to the support of the solutions w^L and $\bar{w}^{CDLP}$ (the order of the sets does not matter). Merging the sets as depicted ensures that the constructed solution w^{CDLP} induces w^L as well as w^l for $l \in \bar{\mathcal{L}}$ (the latter due to the induction assumption on $\bar{w}^{CDLP}$).

Now remove all the solution components $w_{S_L}^L$ and $w_{S_l}^l$ with positive support for all $l \in \mathcal{L}^{\text{int}}$ with $S_l \cap S^{\text{int}} = U$ and $S_L \cap S^{\text{int}} = U$. After the removal, the product cut equations for the remaining solution remain valid because of the equalities (5–6). We repeat this merging process by taking a maximal $U \subseteq S^{\text{int}}$ at each stage till we conclude with $U = \emptyset$. At every stage, as U is a maximal set, all the sets that contained U , namely sets of the form $U \subsetneq S_l \cap S^{\text{int}}, l \in \mathcal{L}^{\text{int}}$ were maximal sets in previous stages and therefore accounted for by equalities (5–6) for the set S_l ; now combining it with the product cuts for the set U , we again obtain equalities (5–6).

The solution w^{CDLP} that emerges from this process is feasible to CDLP : it holds that $\sum_{U \subseteq S^{\text{int}}} \tau(U) = T$ (note that $U = \emptyset \subset S^{\text{int}}$), and therefore, by construction, $\sum_S w_S^{\text{CDLP}} = T$. That the capacity constraint of CDLP is satisfied follows from the induction assumption that $\bar{w}^{\text{CDLP}}$ induces $(\bar{w}, \bar{y})$, with $\bar{w} := w$ and $\bar{y} := y$, combined with the fact that (w, y) is feasible to SDCP+ and that we

constructed w^{CDLP} in a way such that we only added capacity consumption equal to that of segment L under solution w^L . So the combined solution also satisfies the induction for L segments.

The objective value of CDLP equals that of SDCP+ because in the merging process we only add products of $C_L \setminus S^{\text{int}}$ to the solution $\bar{w}^{\text{CDLP}}$, and since these products do not influence other segments as they are only in the consideration set of segment L , we only add the contribution of segment L to the objective without a change of the contribution of other segments.

□

Proof of Proposition 2

Proof

Any solution to CDLP induces a feasible solution in SDCP+ with the same value. To show equivalence,

Figure 6: Illustration of the procedure to merge solutions in the support of w^L and $\bar{w}^{\text{CDLP}}$ to obtain w^{CDLP} for a fixed set $U \subseteq S^{\text{int}}$.

we have only to construct a feasible solution to *CDLP* from a feasible solution to *SDCP+* with the same objective value. We refer to a subset of $S \subseteq C_l$ with a positive weight w_S^l in *SDCP+* as positive in l .

Assume without loss of generality that a solution to *SDCP+* has at least one set $S_1 \subseteq C_1$ with $w_{S_1}^1 > 0$. Note that segment node 1 is a leaf in the intersection graph due to the fully nested structure. Then there are sets $S_l \subseteq C_l$ such that $w_{S_1}^1 \geq w_{S_l}^l$ for all $l > 1$. Moreover, we can describe a sequence of maximal nested sets S_l for $l = 2, \dots, L$ containing S_1 that, due to the product cuts (4), have positive $w_{S_1}^1$ and these variables are non-increasing: namely, $S_1 \subseteq S_2 \subseteq \dots \subseteq S_L$ and $w_{S_1}^1 \geq \dots \geq w_{S_L}^L > 0$. Moreover, as the consideration sets are nested, the product cuts (4) imply that $S_{l-1} = S_l \cap C_{l-1}$, as no positive maximal set in $l-1$ contains S_{l-1} .

As each is maximal within its consideration set, these sets have the property that S_l is not contained in any set $S \subseteq C_l$ of segment l with positive weight $w_S^l > 0$. Therefore, we can create a solution to *CDLP* by giving the maximal set S_L a weight $w_{S_L}^L$ and subtracting $w_{S_L}^L$ from $w_{S_1}^1, \dots, w_{S_L}^L$. Now repeat this operation, at each step peeling off the maximal sequence of weights, to obtain a solution to *CDLP*. Note that this procedure terminates in a finite number of steps.

The two solutions have the same objective value: $\sum_S \lambda R(S) w_S = \sum_l \lambda_l \sum_{S_l} R^l(S_l) [\sum_{S: S \cap C_l = S_l} w_S] = \sum_l \lambda_l \sum_{S_l} R^l(S_l) w_{S_l}^l$ where $\sum_{S: S \cap C_l = S_l} w_S = w_{S_l}^l$ holds by construction for all $S_l \subset C_l$ for all l .

□