1
2 DR. JACOB PONTOPPIDAN THYSSEN (Orcid ID : 0000-0003-3770-1743)
3 DR. TARUNVEER S AHLUWALIA (Orcid ID : 0000-0002-7464-3354)
4 DR. NATALIA BALLARDINI (Orcid ID : 0000-0001-6759-346X)
5 MR. PETER BAGER (Orcid ID : 0000-0002-3577-8459)
6 DR. LIESBETH DUIJTS (Orcid ID : 0000-0001-6731-9452)
7 MR. NIELS J. ELBERT (Orcid ID : 0000-0001-5936-4050)
8 DR. JOACHIM HEINRICH (Orcid ID : 0000-0002-9620-1629)
9 DR. SUSANNE LAU (Orcid ID : 0000-0002-5189-4265)
10 PROF. PATRICK G HOLT (Orcid ID : 0000-0003-1193-0935)
11 PROF. HANS BISGAARD (Orcid ID : 0000-0003-4131-7592)

12
13
14 Article type : Letter to the Editor

15
16
17 **Interaction between filaggrin mutations and neonatal cat exposure in atopic dermatitis**
18 **Letter to the editor**

19
20 Jacob P. Thyssen¹, Tarunveer S. Ahluwalia², Lavinia Paternoster³, Natalia Ballardini^{4,5}, Anna Bergström⁴,
21 Erik Melén^{4,5}, Bo Chawes², Jakob Stokholm², Jonathan O'B Hourihane⁶, Donnchadh M. O'Sullivan⁶, Peter
22 Bager⁷, Mads Maelbye^{7,8,9}, Mariona Bustamante^{10,11,12}, Maties Torrent^{12,13,14}, Ana Esplugues^{12,15,16}, Liesbeth
23 Duijts^{17,18}, Chen Hu^{17,19,20}, Niels J. Elbert^{19,20}, Suzanne G.M.A. Pasmans²⁰, Tamar E.C. Nijsten^{17,18,20},
24 Andrea von Berg²¹, Marie Standl²², Tamara Schikowski²³, Gunda Herberth²⁴, Joachim Heinrich^{22,25,26},
25 Young-Ae Lee²⁷, Ingo Marenholz²⁷, Susanne Lau²⁸, John A. Curtin²⁹, Angela Simpson²⁹, Adnan
26 Custovic³⁰, Craig E. Pennell³¹, Carol A. Wang³¹, Patrick G. Holt³², Hans Bisgaard^{2*}, Klaus Bønnelykke^{2*}

27
28 * These authors contributed equally to the study

- 29
30 1) Department of Dermatology and Allergy, Herlev and Gentofte Hospital, University of Copenhagen,
31 Hellerup, Denmark.
32 2) COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte
33 Hospital, University of Copenhagen, Copenhagen, Denmark.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.14162](#)

This article is protected by copyright. All rights reserved

- 3) MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, UK.
- 4) Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden
- 5) Sachs' Children and Youth Hospital, Södersjukhuset, Stockholm, Sweden
- 6) Paediatrics and Child Health and INFANT Centre, University College Cork, Ireland.
- 7) Statens Serum Institut, Copenhagen, Denmark.
- 8) Department of Clinical Medicine, University of Copenhagen.
- 9) Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA
- 10) ISGlobal, Barcelona Institute for Global Health, Barcelona, Spain
- 11) Universitat Pompeu Fabra (UPF), Barcelona, Spain
- 12) CIBER de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain.
- 13) Ib-salut, Area de Salut de Menorca, Spain
- 14) Institut d'Investigació Sanitària Illes Balears (IdISBa), Spain
- 15) FISABIO– Universitat Jaume I –Universitat de València Joint Research Unit of Epidemiology and Environmental Health. Avenida de Catalunya 21, 46020, Valencia, Spain
- 16) Nursing Department, Faculty of Nursing and Chiropody, Universitat de València, Av. Blasco Ibáñez, 13, 46010 Valencia, Spain
- 17) Department of Pediatrics, Division of Respiratory Medicine and Allergology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands.
- 18) Department of Pediatrics, division of Neonatology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands.
- 19) The Generation R Study Group, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands.
- 20) Department of Dermatology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands. Research Institute, Department of Pediatrics, Marien-Hospital Wesel, Wesel, Germany
- 21) Research Institute, Department of Pediatrics, Marien-Hospital Wesel, Wesel, Germany
- 22) Institute of Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany
- 23) IUF-Leibniz Research Institute for Environmental Medicine, Düsseldorf, Germany
- 24) Department of Environmental Immunology/Core Facility Studies, Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany
- 25) Institute and Clinic for Occupational, Social and Environmental Medicine, University Hospital, Ludwig Maximilian University (LMU), Munich, Germany
- 26) Allergy and Lung Health Unit, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Australia.

27) Max-Delbrück-Center (MDC) for Molecular Medicine, Berlin, Germany; Clinic for Pediatric Allergy, Experimental and Clinical Research Center of MDC and Charité Universitätsmedizin Berlin, Berlin, Germany

28) Department of Pediatrics, Division of Pneumology, Immunology Charité, Universitätsmedizin Berlin, Berlin, Germany

29) Division of Infection Immunity and Respiratory Medicine, School of Biological Sciences, The University of Manchester, Manchester Academic Health Science Centre, and Manchester University NHS Foundation Trust

30) Section of Paediatrics, Department of Medicine, Imperial College London, London, United Kingdom

31) School of Medicine and Public Health, Faculty of Medicine and Health, University of Newcastle, Callaghan, New South Wales, Australia

32) Telethon Kids Institute, University of Western Australia, Nedlands, Australia

Total word count: 978

Number of figures/tables: 2

Number of supplementary figure/tables: 3

Number of references: 9

Running head: Interaction between *FLG* mutations and neonatal cat exposure in AD

Funding sources: Please see supplementary information.

Conflicts of interests: The authors have no relevant conflict of interest to disclose.

Prior presentations: The content has not been published previously and is not otherwise submitted for publication.

1 Correspondence:

2
3 Professor Klaus Bønnelykke, MD, PhD
4 COPSAC, Copenhagen Prospective Studies on Asthma in Childhood
5 Herlev and Gentofte Hospital
6 Ledreborg Alle 34
7 2820 Gentofte
8 Denmark
9 E-mail: kb@copsac.com
10 Website: www.copsac.com
11

12 To the editor.

13 Atopic dermatitis (AD) is a prevalent inflammatory skin disease. Loss-of-function mutations in filaggrin
14 gene (*FLG*) represent the strongest genetic risk factors for AD, being strongly associated with early disease
15 onset and persistence into adulthood. The epidermis of individuals with mutations in *FLG* is fundamentally
16 different from normal skin being characterized by increased penetration of allergens.

17

18 Recent birth cohort studies showed a significant interaction between cat ownership at birth and mutations in
19 *FLG* (R501X, 2282del4) on the development of early-onset AD.¹ This finding was replicated for the
20 2282del4 *FLG* mutation in a Dutch cohort study, and extended to further associate with risk of allergic
21 sensitization.² We performed analyses in multiple birth cohorts to examine the consistency and overall
22 strength of the previously observed interaction.

23

24

25 Consortium collaborators were invited to participate in the study, ³ and 13 birth cohorts provided data on
26 cat exposure, AD, and *FLG* mutations (Table 1 and Supplementary Table 1+2). All cohorts had information
27 on the most common mutations in *FLG*, R501X and 2282del4, and the majority also had information on
28 R2447X and S3247X (Table 1). Heterozygous, compound heterozygous and homozygous *FLG* mutation
29 carriers were pooled as mutation carriers. Cat ownership/exposure was based on questionnaires or
30 interviews. AD diagnoses were based on questionnaires in 10 cohorts, by physician examination in 2
31 cohorts (COPSAC2000, COPSAC2010), and a combination in 1 cohort (MAS). For further details, please
32 see supplementary information online.

33

34 The predetermined primary outcome was AD onset before one year of age (*AD_{early}*) based on previous
35 observations. ¹ Secondary outcomes included i) current AD at seven years of age or the year of assessment

closest to, but before, 7 years (' $AD_{current}$ '), and ii) a history of AD during the first 7 years of life, or last year of assessment (AD_{ever}). For further details, please see supplementary information online.

A total of 22,133 children were studied (Table 1 and supplementary Table 1). The median prevalence (range) of mutations in *FLG* was 9.4% (4.6-12.3), cat exposure 15% (7.9-29.6), AD_{early} , $AD_{current}$, and AD_{ever} , respectively, 18% (9.7-34.6), 13.9% (3.9-20), and 39.5% (20.4-67). There was no interaction between *FLG* mutations and cat exposure on the risk of the primary outcome ' AD_{early} ' (OR 1.10 (95% CI 0.86-1.43), $P=0.0$), (Figure 1 and Table 1). There was a statistically significant interaction for the secondary outcome of having AD at last time of examination or questioning at 7 years of age ($AD_{current}$), in the direction of increased risk of AD from cat exposure in children with *FLG* mutations (OR 1.36 (95% CI 1.02-1.82) $P=0.0086$), but this was not statistically significant after adjustment for multiple testing. The *FLG*-stratified analyses showed a trend towards cat exposure being a risk factor in children with *FLG* mutations and a protective factor in children without *FLG* mutations (Figure 1 and Supplementary Table 3). No interaction was found for the other secondary outcome ' AD_{ever} ' (OR 1.06 (95% CI 0.82-1.37), $P=0.63$)

We found no interaction between cat exposure in infancy and mutations in *FLG* on 'early-onset AD' or 'AD ever'. A nominally significant interaction in the expected direction was found for the secondary outcome 'current AD' at 7 years of age, but this did not survive adjustment for multiple testing.

A particular study strength is the large number of independent birth cohorts with prospective assessment of exposure and outcomes. Most cohorts had genotype information for the 4 most common *FLG* mutations ensuring a high degree of correct classification. It is a limitation that AD diagnoses were based on questionnaire data in most cohorts, potentially reducing diagnostic specificity. Since AD is a chronic and relapsing disease, short episodes of other eczemas, e.g. due to irritant or allergic contact dermatitis, may be misinterpreted as AD by parents and caregivers, in particular in the first years of life where flexural accentuation is not yet occurring.⁴ Notably, the high prevalence of early AD in some cohorts could mask a true cat exposure-*FLG* mutation interaction. One may argue that AD measured at 7 years of age is expected to have a higher specificity due to flexural involvement.⁵ It is another limitation that cat exposure was only assessed around birth, and it is possible that later exposure to cat could have an unmeasured effect on AD. Similar, the extent of cat exposure might vary between studies and families. Other environmental factors were not included since covariate availability differed between the cohorts. Reverse causality cannot be excluded, as families who had experienced atopic disease might have avoided having pets to prevent allergic disease in their (next) child. However, one would expect families with *FLG* mutations, and thereby increased risk of eczema, to avoid cat ownership, which would tend towards an apparent protective effect of having a cat.

1
2 No association between cat ownership and AD was found in another meta-analysis of 13 studies (relative
3 risk 0.94 (95%CI 0.76-1.16)).⁶ When a compelling gene-environment interaction was observed between
4 *FLG* mutations and cat ownership on the risk of early-onset AD in birth cohorts, it raised the possibility
5 that preventive measures against pediatric AD could be identified by taking the genetic susceptibility into
6 account.^{1, 2} The COPSAC2000 study, which provided the basis for the previous report of interaction
7 between cat and *FLG* mutations,¹ benefited from close follow-up of children, and high AD diagnostic
8 accuracy, whereas most birth cohorts in the present meta-analysis used questionnaires, potentially
9 explaining the discrepancy between the studies.

10
11 No pathomechanism has been established for the proposed association between cat ownership and AD in
12 *FLG* mutation carriers. Possibly, very small cat allergens might penetrate into the viable layers of the
13 epidermis, where they can exert immune effects, possibly through IL-1 β promotion.^{7, 8} Studies
14 demonstrating increased risk of peanut allergy *FLG* mutation carriers, also suggest increased peanut
15 allergen skin penetration.⁹

16
17 In conclusion, this meta-analysis could not confirm an interaction between cat exposure in infancy and
18 *FLG* mutations on development of early-onset AD. Gene-environment interactions remain largely
19 unknown.

Cohort	ALSPAC	BAMSE	Baseline	COPSAC 2000	COPSAC 2010	DNBC	Generation R	GINplus	INMA*	LISA	MAS	MAAS	RAINE
Cohort inclusion criteria	Children from the general population	Children from the general population	Children from the general population	Children born from mothers with asthma	Children from the general population	Children from the general population	Children from the general population	Children from the general population	Children from the general population	Children from the general population	Children from the general population	Children from the general population	Children from the general population
Study year baseline	1991-1992	1994-1996	2008	2000	2010	1996-2001	2002-2006	1995-98	1997-2006**	1997-99	1990	1996-1997	1989-1991
Filaggrin mutations genotyped	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4, R2447X	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4	R501X, 2282del4, R2447X, S3247X	R501X, 2282del4	R501X, 2282del4, R2447X, S3247X	R501X; 2282del4, R3247X, R2447X	R501X, 2282del4, S3247X, rs138726443
Proportion with FLG mutations	11% (834/7743)	7.2% (138/1906)	10.8% (146/1344)	12.3% (49/396)	10.3% (72/700)	9.7% (91/935)	9.4% (268/2849)	7.0% (104/1490)	4.6% (28/606)	7.0% (69/987)	9.7% (79/813)	10.1% (87/864)	9.3% (140/1500)
Basis for atopic dermatitis diagnosis	Questionnaire	Questionnaire	Questionnaire and clinical diagnosis	Clinical diagnosis	Clinical diagnosis	Questionnaire	Questionnaire	Questionnaire	Questionnaire	Questionnaire	Questionnaire and clinical diagnosis	Questionnaire	Questionnaire
AD 'early onset (≤1y)'	18% (1368/7743)	16.9% (323/1906)	22.7% (292/1282)	25.3% (100/396)	11.1% (78/700)	14.7% (137/935)	21.6% (545/2521)	11.3% (167/1477)	31.7% (192/606)	9.7% (94/973)	14.14% (115/813)	34.6% (160/462)	22.8% (341/1498)
AD 'current'	20% (1270/6402)	17.2% (327/1896)	15.2% (178/1168)	13.9% (55/396)	13.6% (95/700)	6.8% (65/963)	18.6% (496/2658)	5.93% (77/1298)	19.5% (118/604)	3.9% (32/825)	7.5% (58/773)	14.1% (96/681)	13.3% (184/1386)
AD 'ever (0-7 y)'	67% (4367/6501)	39.5% (743/1878)	26.3% (354/1344)	42.2% (175/396)	27.6% (193/700)	20.4% (196/963)	41.4% (1180/2849)	35.3% (447/1266)	49.6% (307/618)	32.6% (269/826)	36.2% (294/813)	60.7% (306/504)	39.7% (596/1500)
AD assessment	6, 18, 30, 42, 57, 69, 81 months	1, 2, 4 and 8 years	6, 12 and 24 months	1 month, and then	1 month, and then	6 and 18 month, and 7 years	6 months, and 1, 2, 3, 4 and 6 years	1, 2, 3, 4, 6 and 10 years	1, 2, and 4 years	6, 12, 18, 24 months and 4, 6	1, 3, 6, 12, 18, 24 month and then	1,3,5,8 years	1, 3, 5, 8 years

time-points				every 6 months.	every 6 months.					and 10 years	yearly		
Child age at cat exposure assessment	During pregnancy	At baseline (3 months) and/or 1 year follow-up.	6 months and 12 months	Birth	Birth	18 months	age < 1 year)	1 year	1 year	3 months and 1 year	3 months	Birth	1 year
Early life cat exposure %	29.6	10.6**	8.8	15	20	16.4-22.4 **	25.3	7.9	11.5	12.2	12.7	20.5	18

Table 1. Baseline characteristics of participants in birth cohorts.

*INMA sub cohorts: VAL, SAB, MEN
 ** please see supplementary Table 1 for further details.

Reference List

- (1) Bisgaard H, Simpson A, Palmer CN, Bonnelykke K, McLean I, Mukhopadhyay S, et al. Gene-environment interaction in the onset of eczema in infancy: filaggrin loss-of-function mutations enhanced by neonatal cat exposure. PLoS Med 2008 Jun 24;5(6):e131.
- (2) Schuttelaar ML, Kerkhof M, Jonkman MF, Koppelman GH, Brunekreef B, de Jongste JC, et al. Filaggrin mutations in the onset of eczema, sensitization, asthma, hay fever and the interaction with cat exposure. Allergy 2009 Dec;64(12):1758-65.
- (3) Middeldorp CM, Felix JF, Mahajan A, McCarthy MI. The Early Growth Genetics (EGG) and EARly Genetics and Lifecourse Epidemiology (EAGLE) consortia: design, results and future prospects. Eur J Epidemiol 2019 Mar 18.
- (4) Nakamura T, Haider S, Colicino S, Murray CS, Holloway J, Simpson A, et al. Different definitions of atopic dermatitis: impact on prevalence estimates and associated risk factors. Br J Dermatol 2019 Mar 1.
- (5) Paternoster L, Savenije OEM, Heron J, Evans DM, Vonk JM, Brunekreef B, et al. Identification of atopic dermatitis subgroups in children from 2 longitudinal birth cohorts. J Allergy Clin Immunol 2018 Mar;141(3):964-71.
- (6) Pelucchi C, Galeone C, Bach JF, La VC, Chatenoud L. Pet exposure and risk of atopic dermatitis at the pediatric age: a meta-analysis of birth cohort studies. J Allergy Clin Immunol 2013 Sep;132(3):616-22.
- (7) Davila I, Dominguez-Ortega J, Navarro-Pulido A, Alonso A, Antolin-Amerigo D, Gonzalez-Mancebo E, et al. Consensus document on dog and cat allergy. Allergy 2018 Jun;73(6):1206-22.
- (8) Rahrig S, Dettmann JM, Brauns B, Lorenz VN, Buhl T, Kezic S, et al. Transient epidermal barrier deficiency and lowered allergic threshold in filaggrin-hornerin (FlgHrnr(-/-)) double-deficient mice. Allergy 2019 Jul;74(7):1327-39.
- (9) Brown SJ, Asai Y, Cordell HJ, Campbell LE, Zhao Y, Liao H, et al. Loss-of-function variants in the filaggrin gene are a significant risk factor for peanut allergy. J Allergy Clin Immunol 2011 Mar;127(3):661-7.

Figure 1. Interaction between cat exposure and common *FLG* mutations in relation to a) Early onset atopic dermatitis, b) Current atopic dermatitis and c) atopic dermatitis in the first 7 years of life.

A.

all_14162_f1a.tiff

B.

all_14162_f1b.tiff

C.

Figure 1. Interaction with cat exposure and common FLG mutations in relation to A) early onset atopic dermatitis, B) current atopic dermatitis, and C) atopic dermatitis in the first 7 years of life.

all_14162_f1c.tiff