Title: Axonal marker neurofilament light (NfL) predicts long-term outcomes and progressive neurodegeneration after traumatic brain injury

Authors: Neil S.N. Graham^{†1,2}, Karl A. Zimmerman^{†1,2}, Federico Moro³, Amanda Heslegrave^{4,5}, Samia Abed Maillard⁶, Adriano Bernini⁶, John-Paul Miroz⁶, Cornelius K Donat¹, Maria Yanez Lopez⁷, Niall Bourke^{1,2}, Amy E Jolly^{1,2}, Emma-Jane Mallas^{1,2}, Eyal Soreq^{1,2}, Mark H Wilson^{8,9}, Gavin Fatania¹⁰, Dylan Roi¹⁰, Maneesh C Patel¹⁰, Elena Garbero¹¹, Giovanni Nattino¹¹, Camelia Baci¹², Enrico Fainardi¹³, Arturo Chiericato¹², Primož Gradisek¹⁴, Sandra Magnoni¹⁵, Mauro Oddo^{6,16}, Henrik Zetterberg^{17,18}, Guido Bertolini¹¹, David J. Sharp^{*1,2,19}

Affiliations:

¹Department of Brain Sciences, Imperial College London; London W12 0NN, UK.

²UK DRI Centre for Care Research and Technology, Imperial College London W12 0BZ; London, UK.

³Laboratory of Acute Brain Injury and Therapeutic Strategies, Department of Neuroscience, Istituto di Ricerche Farmacologiche Mario Negri IRCCS; 24126 Bergamo, Italy.

⁴Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square; London WC1N 3BG, UK.

⁵UK Dementia Research Institute at UCL; London WC1N 3BG, UK.

⁶Neuroscience Critical Care Research Group, Department of Intensive Care Medicine, CHUV Lausanne University Hospital and University of Lausanne; 1011 Lausanne, Switzerland.

⁷Centre for the Developing Brain, School of Biomedical Engineering and Imaging Sciences, King's College London; London SE1 7EH, UK.

⁸Department of Neurosurgery, Imperial College Healthcare NHS Trust; London W6 8RF, UK.

⁹Department of Surgery & Cancer, Faculty of Medicine, Imperial College London; London SW7 2AZ, UK

¹⁰Department of Imaging, Imperial College Healthcare NHS Trust; London W6 8RF, UK

¹¹Laboratory of Clinical Epidemiology, Istituto di Ricerche Farmacologiche Mario Negri IRCCS; 24126 Bergamo, Italy.

¹²Neuroranimazione, ASST Grande Ospedale Metropolitano Niguarda; 20162 Milano, Italy

¹³Department of Experimental and Clinical Sciences, Careggi University Hospital, University of Firenze; 50139 Florence, Italy.

¹⁴Clinical Department of Anaesthesiology and Intensive Therapy, University Medical Center; 1000 Ljubljana, Slovenia

¹⁵Department of Anesthesia and Intensive Care, Santa Chiara Hospital; 38122 Trento, Italy

¹⁶Medical Direction, CHUV Lausanne University Hospital; 1011 Lausanne, Switzerland

¹⁷Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, the Sahlgrenska Academy at the University of Gothenburg; 431 41 Mölndal, Sweden

¹⁸Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital; 431 41 Mölndal, Sweden

¹⁹Centre for Injury Studies, Imperial College London SW7 2AZ; London, UK

* Corresponding author: david.sharp@imperial.ac.uk

†contributed equally

One Sentence Summary:

Axonal injury after TBI can be reliably quantified using plasma NfL, which predicts long-term functional outcomes and progressive neurodegeneration.

Abstract:

Axonal injury is a key determinant of long-term outcomes after traumatic brain injury (TBI) but has been difficult to measure clinically. Fluid biomarker assays can now sensitively quantify neuronal proteins in blood. Axonal components such as neurofilament light (NfL) potentially provide a diagnostic measure of injury. In the multi-centre BIO-AX-TBI study of moderate-severe TBI we investigated relationships between fluid biomarkers, advanced neuroimaging and clinical outcomes. Cerebral microdialysis was used to assess biomarker concentrations in brain extracellular fluid aligned with plasma measurement. An experimental injury model was used to validate biomarkers against histopathology. Plasma NfL increased after TBI, peaking at ten days to six weeks but remaining abnormal at one year. Concentrations were around ten times higher early after TBI than in controls (patients with extracranial injuries). NfL concentrations correlated with diffusion MRI measures of axonal injury and predicted white matter neurodegeneration. Plasma TAU predicted early grey matter atrophy. NfL was the strongest predictor of functional outcomes at one year. Cerebral microdialysis showed that NfL concentrations in plasma and brain extracellular fluid were highly correlated. An experimental injury model confirmed a dose-response relationship of histopathologically defined axonal injury to plasma NfL. In conclusion, plasma NfL provides a sensitive and clinically meaningful measure of axonal injury produced by TBI. This reflects the extent of underlying damage, validated using advanced MRI, cerebral microdialysis and an experimental model. The results support the incorporation of NfL sampling subacutely after injury into clinical practice to assist with the diagnosis of axonal injury and to improve prognostication.

Main Text:

INTRODUCTION

Long-term outcomes of traumatic brain injury (TBI) are frequently poor (*1*), with around half of patients affected by residual disability (*2, 3*). Traumatic axonal injury (TAI) is a key determinant of these clinical outcomes (*4, 5*). However, validated methods to quantify the extent of axonal injury are lacking in a clinical context. Several proteins are potential biomarkers of axonal injury after TBI. These include cytoskeletal proteins neurofilament light (NfL) and microtubule associated protein TAU (TAU), as well as the more the more widely expressed neuronal ubiquitinating enzyme ubiquitin carboxyl-terminal hydrolase L1 (UCH-L1). Proteins whose expression is predominantly astroglial, such as calcium binding protein S100 B (S100B), or glial fibrillary acidic protein (GFAP) may also provide a surrogate readout of axonal damage by quantifying the inflammatory response and non-neuronal tissue damage (*6*).

Neurofilament light has shown diagnostic, monitoring and prognostic utility across several neurological diseases (*7*). Advanced immunoassays such as single molecule array (Simoa) allow ultra-low concentrations in venous blood to be accurately measured, with concentrations reflecting cerebrospinal fluid concentrations (*8-10*). After moderate-severe TBI, NfL is acutely elevated in plasma, sometimes remaining increased years after injury (*8, 11*). Plasma concentrations continue to rise post-injury, in contrast to other blood biomarkers which typically peak within days (*10, 12*). Initial work shows that blood concentrations of NfL can predict late clinical outcomes, although it is uncertain how much prognostic information is provided when compared to other biomarkers such as GFAP, UCH-L1, TAU, and S100B (*8, 10, 13, 14*). The optimal time to sample NfL after TBI is unclear as the dynamics of its fluctuation over time have not been characterised in detail.

The source of plasma NfL after TBI is also unclear, which limits its diagnostic value. NfL is a type IV intermediate filament that is found within axons. It is present in high quantities within long myelinated axons in the cerebral deep white matter that are susceptible to TAI (*15, 16*). Hence, plasma concentrations of NfL may reflect its release from white matter damaged following head injury. One way to assess this possibility is to directly measure concentrations from extracellular fluid collected from damaged white matter. This has not been done previously but can be achieved using invasive cerebral microdialysis with catheters permeable to neuronal breakdown products. High concentrations of TAU in brain extracellular fluid early post-injury have been demonstrated using this method, with concentrations correlating with imaging evidence of axonal injury (*17*). Preliminary microdialysis work with NfL has shown increased concentrations in peri-contusional

and apparently normal-appearing white matter regions that correlate with extracellular TAU concentrations (18). However, NfL concentrations in microdialysate and peripheral venous samples have not been compared. Establishing that plasma and extracellular NfL correlates helps to clarify its origin and the simultaneous quantification of a range of brain injury proteins (for example NfL, TAU, UCH-L1 and GFAP) in microdialysate will help understand whether it provides unique information.

Diffusion tensor imaging (DTI) provides another way to assess the relationship between plasma NfL and axonal injury within the brain. Fractional anisotropy is widely used as a measure of axonal injury that: (a) correlates with histopathological evidence of TAI (19); (b) provides prognostic information about long-term clinical outcomes; and (c) predicts the pattern of progressive white neurodegeneration indicated by brain atrophy (5, 20). Hence, if plasma NfL originates from damaged axons, concentrations should correlate with diffusion imaging measures of TAI in the brain and might also predict the extent of post-traumatic neurodegeneration as quantified by white matter atrophy (8).

Plasma NfL could also originate outside the brain and it remains unclear how specific increased concentrations are to TBI. Mutations in the NEFL gene, which encodes NfL, cause the hereditary peripheral neuropathy Charcot Marie Tooth disease and blood NfL concentrations can be elevated in both chronic (21) and acute peripheral neuropathies (22). This is important as TBI often occurs alongside peripheral nerve injury, which can potentially increase NfL. This has been an issue for other potential blood biomarkers in TBI. For example, S100B has substantial expression in adipose tissue as well as astrocytes, confusing the interpretation of high concentrations when extensive soft tissue injury is present (23). Hence, comparing NfL concentrations in groups of patients with TBI and non-TBI trauma is necessary for a complete understanding of the specificity of NfL as a measure of traumatic axonal injury in the central nervous system.

Experimental injury models of TBI can be used to test the relationship between blood biomarkers and controlled injuries, providing ground truth validation. The controlled cortical impact (CCI) model provides a reproducible means to control the extent and distribution of injury, and to examine the associated histopathological effects (15, 24, 25). High field magnetic resonance imaging can be used to provide diffusion imaging measures aligned to human studies, allowing histopathological effects of injury to be compared with neuroimaging measures of axonal injury and blood concentrations of NfL. Previous experimental work on blood NfL in TBI is limited but

suggests a relationship between plasma NfL concentrations and neurofilament-immunoreactive axonal swellings indicative of injury (26, 27). A dose-response relationship of NfL to injury has not previously been directly established.

The multicentre prospective BIO-AX-TBI study (NCT03534154) and aligned experimental injury model were conducted to: (a) investigate the optimal timing of axonal injury fluid biomarker assessment after single moderate-severe TBI; (b) assess how these relate to radiological measures of axonal injury and clinical outcomes; and (c) compare plasma biomarker concentrations against brain extracellular fluid in humans and ground truth histopathology in an experimental injury model (Fig. 1) (28). We tested the following specific hypotheses in relation to NfL: (a) concentrations will increase after TBI and this will be detectable in plasma; (b) specific changes in NfL will be seen after TBI when compared to patients with extracranial injuries but no TBI; (c) plasma NfL will correlate with concentrations in extracellular fluid on cerebral microdialysis; (d) plasma NfL will correlate with diffusion MRI measures of axonal injury, specifically reduced FA; (e) plasma NfL concentrations will relate to histopathological evidence of axonal injury in the CCI model; (f) plasma NfL will predict white matter atrophy over time; and (g) plasma NfL will predict clinical outcomes at six and twelve months post-injury.

RESULTS

Demographics, clinical characteristics and outcomes of patients and healthy controls

One hundred ninety-seven patients were recruited after moderate-severe TBI from eight major trauma centres (Table S1). Median age at injury was 47 years (IQR 30) with 77% (152) male (Fig. 2A,B). 74% (146) underwent MRI, with a median age of 44 (IQR 30) of whom 80% (116) were male. Injuries were high energy in 56% (110), comprising falls from more than three metres or collision at more than 30 km/h (Table S2). 91% (180) of injuries were accidental, with 47% (93) road-traffic related. Acute loss of consciousness was definite or suspected in 75% (147) of patients and 69% (136) had definite or suspected retrograde amnesia. The cohort included a range of severities within the moderate-severe classification, with pre-hospital Glasgow Coma Scale (GCS) 3 to 8 in 45% (89) and a median lowest GCS on admission of 7.5 (IQR 8) (Fig. 2C).

Acute CT (Fig. 2E) identified parenchymal contusions in 66% (130), subarachnoid haemorrhage in 60% (118), subdural haematoma in 57% (113) and extradural haematomas in 22% (44). Diffuse axonal injury was reported on CT in 18% (35). Raised intracranial pressure was present in 29% (58) of patients and was persistent/refractory in 16% (31). Neurosurgical evacuation or decompression were performed in 34% (66) of patients.

Subacute MRI was performed, between ten days to six weeks post-injury in 71% (140) of patients (Fig. 2F), a time-window chosen to ensure stabilisation of dynamic diffusion changes. Parenchymal contusions were present in 85% (119), with subarachnoid haemorrhage in 21% (30), subdural haematomas in 68% (95), extradural haematomas in 4% (6), oedema in 27% (38) and microhaemorrhages in 69% (97). Group-wise lesion maps were hand-drawn for each scan to be accounted-for in the imaging analysis (fig. S1).

During the study period, 7% (13) of patients died, at a median of 11 days post-injury. Functional outcomes were assessed over a year (Fig. 2G). All patients had a known outcome at six or twelve months post injury: specifically, the GOSE was available in 94% (185) at 6 months and 82% (161) at 12 months. At six months, 29% (58) of patients made a good recovery, defined as GOSE score 7 or 8. Six individuals were in a vegetative state (GOSE 2) at 6 months, and were unchanged at 12 months. By 12 months post-injury 32% (62) of patients made a good recovery.

To assess the blood biomarker specificity, 25 patients hospitalised with non-TBI traumatic (NTT) extracranial injuries were recruited (for injury details see Table S3). These patients, who had no head injury history, had a median age at injury of 41 (IQR 29) of whom 92% (23) were male.

Patients with NTT had fluid biomarker assessment twice in the first ten days post-injury, aligned to the TBI group. The groups did not differ significantly in age or sex ($P>0.05$).

Healthy volunteers were assessed across all the sites, comprising 128 individuals with a median age of 39 (IQR 27) of whom 78% (61) were male. 95% (121) underwent MRI and 75% (96) had blood biomarker assessment. This group was slightly younger ($W = 10784$, $P = 0.028$) and more female ($X^2 9.10$, $df = 1$, $P = 0.003$) than the TBI patients. However, there was no significant age difference between patients and healthy controls who underwent MRI ($P>0.05$). To contextualize MRI atrophy rate assessments after TBI within normal life-course change, 30 healthy controls with a median age of 40 (IQR 27), of whom 70% (21) were male, were imaged twice. Neither sex nor age differed significantly between the main and longitudinal healthy control groups. ($P>0.05$, Table S1).

Blood biomarker profiles after traumatic brain injury

Injury markers including plasma NfL were serially assessed on two occasions within ten days of injury, subacutely at ten days to six weeks, six and twelve months (Fig. 3). Plasma NfL was markedly elevated post injury (Fig. 3A,B), increasing over time and peaking subacutely, around 20 days post-injury (median peak NfL 507.3 pg/ml, IQR 728.2). NfL remained elevated at six and twelve months post-injury compared with healthy controls (both $P<0.001$, $W=387/431$, respectively, Fig. 3C). 85% (45/53) of TBI patients had abnormally high NfL at six months (defined as > 2 standard deviations from the healthy control mean). 52% (13/25) remained elevated at twelve months post-injury (Fig. 3C, Table S4). Age at injury was not a significant determinant of NfL concentration at 6 or 12 months using linear regression (fig. S2).

TBI patients also had elevations of plasma total TAU, serum S100B, plasma UCH-L1 and GFAP compared with healthy controls (all $P<0.05$). Concentrations peaked within the first day of injury (median peak for TAU 7.0pg/ml; S100B 215.1pg/ml; UCH-L1 97.6pg/ml; GFAP 12,590.5 pg/ml). These biomarkers normalised following the subacute visit except for plasma GFAP, which remained chronically raised at twelve months ($W=691$, $P=0.041$; Table S4). 34% (18/53) of patients had raised plasma GFAP at six, as did 32% (8/25) at twelve months. Age at injury modestly contributed to GFAP concentrations at six (adjusted $R^2 0.27$, $P < 0.001$) but not twelve months post injury. Patients with raised plasma GFAP at six months were significantly older than those with normal concentrations ($W=134$, $P=0.001$).

Similar plasma concentrations of NfL and TAU were seen in patients with and without visible pathology on CT. There were no significant differences ($P>0.05$) in peak NfL or TAU when comparing Marshall grade I patients ($n=23$, no lesions on scan) with Marshall grades II-VI ($n=168$, encompassing all diffuse / mass lesions, Table S5). In contrast, higher peak concentrations of UCH-L1 ($W=910$, $P=0.033$), S100B ($W=776$, $P=0.0027$) and GFAP ($W=779$, $P=0.005$) were seen in patients with Marshall grades II-VI versus grade I (Table S5). Similarly, there were no significant differences ($P>0.05$) in peak NfL between patients with lesions on MRI ($n=125$) compared to those without lesions ($n=18$) (Table S5). Higher concentrations of TAU ($W=1518$, $P<0.001$), UCH-L1 ($W=1428$, $P<0.001$), S100B ($W=1469$, $P<0.001$) and GFAP ($W=1484$, $P<0.001$) were observed in patients with lesions on MRI (Table S5). Where lesions were present on MRI, peak blood concentrations of NfL, TAU, UCH-L1, S100B and GFAP all correlated with lesion volume (all $P<0.001$, $\rho=0.50, 0.52, 0.52, 0.53, 0.45$ respectively, Table S5).

Neurosurgery was performed in 38% ($n=75$) of patients, motivating an assessment of whether interventions caused elevated peak fluid biomarker concentrations. Peak biomarker values were compared between the surgery and nonsurgery groups in the first 6 weeks following injury. As expected, the surgical group had significantly higher peak NfL after surgery ($W=2920$), TAU ($W=2960$), UCH-L1 ($W=3141$), GFAP ($W=3125$) and S100B concentrations ($W=3111$, all P values < 0.002 , Table S6). This neurosurgical group had more severe injuries than the non-neurosurgical group. Hence the biomarker difference might not be the result of neurosurgery. To investigate this issue, a sensitivity analysis was performed taking into account the white matter FA, in patients for which these data were available ($n=131$). Of these, 37% ($n=49$) underwent surgery. For each biomarker, linear regression was performed between peak plasma concentrations and FA. Group means on the residuals from this regression were then compared for each biomarker, between patients with and without surgery, as above. Accounting for severity in this way, we continued to detect a significant neurosurgery-associated increase in peak NfL concentration ($P=0.042$, $W=1581$) but not the other fluid biomarkers assessed (Table S7).

Patients with no history of TBI ($n=25$) showed lower concentrations of all blood biomarkers compared to the TBI group (all P values <0.001). Non-TBI patients did show increased plasma NfL and UCH-L1 compared to healthy controls, but there were no differences in TAU, GFAP or S100B ($P>0.05$, Fig. 3, Table S4). The difference in age between groups (see Fig. 2A, previously) did not affect the results (see Supplementary Materials - Sensitivity Analysis, Table S8).

NfL, TAU and UCH-L1 concentrations are highly correlated between plasma and brain microdialysate

Eighteen patients underwent cerebral microdialysis and simultaneous plasma sampling within the first ten days of injury (median age 34 years, 78% male, Table S9). Microdialysis was initiated a median of 21 hours post-injury (IQR 25) at the time of craniotomy/cranioplasty, and continued for a median of 6 days in total (IQR 4.75). This allowed blood concentrations to be compared with brain extracellular fluid sampled from white matter potentially affected by traumatic axonal injury. Frontal white matter without evidence of focal brain injury was sampled, from the right hemisphere in 83% (15) and left 17% (3). Indicative catheter placement is shown in Fig. 4A. The microdialysis group had a mortality rate of 28% (5) within a month of injury. They were well matched to the broader TBI group in respect of demographics and admission GCS but had significantly higher rates of refractory raised intracranial pressure ($P<0.001$; $\chi^2=21.1$, Table S1).

NfL, total TAU, UCH-L1 and GFAP concentrations were significantly (all $P<0.001$) higher in brain extracellular fluid than plasma (Fig. 4B, Table S10. Median NfL, TAU and UCH-L1 concentrations were around 100 times higher in microdialysate than plasma, whereas GFAP was around four times higher (Table S10). Linear mixed effects modelling accounting for time since injury and patient variability (individual patient data shown in fig. S2, fig. S3) showed a strong relationship between blood and brain extracellular fluid concentrations for NfL ($P=0.001$, $t=3.3$, $AIC=-128.4$), total TAU ($P=0.003$, $t=2.9$, $AIC=57.8$) and UCH-L1 ($P<0.001$, $t=4.8$, $AIC=19.7$) but not GFAP ($P=0.76$).

Imaging evidence of traumatic axonal injury

Diffusion MRI was investigated to assess evidence of TAI. Individual z-score maps of fractional anisotropy (FA) were generated at each scanning site by comparing patients with local controls. We examined the whole white matter skeleton and corpus callosum. Both showed significant reductions in FA at the subacute, six and twelve month time points (all $P<0.001$, Fig. 5A, Table S11. Significant reductions ($P<0.05$) in FA were also present in a large number of white matter tracts at all three time points (Fig. 5B). FA declined over time between the acute and six month assessments in a large number of white matter tracts including the whole white matter and corpus callosum regions ($n=49$, $V=975$, $p<0.001$)(Fig. 5C,D). Further reductions were present from 6-12

months but were not significant ($n=36$, $V=209$, $P=0.052$). Focal lesions were excluded from the calculation of FA in all analyses (Supplementary methods) (29, 30).

Post-traumatic neurodegeneration

Brain volume changes over time were investigated to assess neurodegeneration triggered by TBI (Fig. 5E). The Jacobian determinant (JD) measure provides a sensitive index of atrophy rates (31). Grey matter JD was reduced, indicating progressive atrophy, in patients during the first six months post-injury ($W=469$, $P<0.001$) with accompanying CSF expansion ($W=1214$, $P=0.002$). In contrast, white matter JD was only reduced six to twelve months post-injury ($W=309$, $P=0.001$) with accompanying CSF expansion ($W=769$, $P=0.005$) but no abnormality of grey matter JD versus healthy controls (Fig. 5E). Linear mixed-effects modelling showed a significant ($P<0.05$) interaction between tissue class and time ($F(2)=8.4$, $P<0.001$). This was the result of lower grey matter JD in the first six months ($V=183$, $P=0.01$), but no significant differences in white matter JD between subacute to six month, and six to twelve month intervals longitudinally, within patients ($P>0.05$, Fig. 5E). Specific analyses of the hippocampal and thalamic regions revealed progressive volume loss between the subacute and six month time points ($W=1287$, $P<0.001$; $W=1151$, $P=0.014$, respectively) but no increased atrophy was observed between six and twelve months compared to healthy controls in these regions of interest (Table S11). Voxelwise whole-brain analyses showed significant longitudinal atrophy of right internal capsule and posterior thalamic radiation from six to twelve months post-injury (map of regions with significantly $P<0.05$ higher atrophy rates than controls shown in Fig. 5F). No other significant ($P<0.05$) group differences were present voxelwise between patients and controls at the first interscan interval, or for other tissue classes at the second interscan interval.

We used linear regression to test the relationship of peak biomarker concentrations in blood within the first six weeks of injury to early grey matter atrophy, and late white matter atrophy. Peak concentrations of all blood biomarkers predicted grey matter atrophy rates in the first six months post-injury (all $P<0.004$, adj. R^2 0.20 to 0.34, Table S12, Fig. 6A), which was strongest for TAU (adj. R^2 0.34, $P<0.001$). In contrast, only peak plasma NfL predicted late white matter atrophy in the six to twelve month interval (adj. R^2 0.14, $P=0.037$). We have previously reported that axonal injury in the chronic phase after TBI measured by FA predicts later neurodegeneration measured by JD (20). Here, we found a similar voxelwise relationship between acute axonal injury and later

white matter neurodegeneration. At each voxel within the white matter skeleton, the subacute measure of axonal integrity, FA, was correlated with the JD atrophy rate between 6 and 12 months. Significant ($P < 0.05$, TFCE-corrected) correlations were present voxelwise within central regions including the corpus callosum, right corona radiata, and left cerebral peduncle (see spatial significance map, Fig. 6B).

Biomarkers and functional outcomes

Long-term functional outcome was measured using the Glasgow Coma Scale Extended (GOSE). This was binarised using a score of six or higher to indicate favourable outcomes. Peak plasma NfL concentration in the first six weeks post-injury, during the subacute time period, had the highest pseudo R^2 value in predicting the GOSE at six and twelve months post injury (McFadden's R^2 , 0.24, 0.48, respectively) (Fig. 6C, Table S12). Z-scored FA in the corpus callosum had the second highest pseudo R^2 , for both six and twelve month GOSE scores. Weaker but significant ($P < 0.05$) other predictors of outcome included peak plasma UCH-L1, plasma GFAP and serum S100B, but not plasma TAU (Table S12).

Plasma NfL correlates with diffusion MRI measures of axonal injury

Next we tested whether peak axonal injury blood biomarker concentrations correlated with diffusion measures of axonal injury. Plasma NfL showed a moderate negative correlation with FA ($\rho = -0.44$, $P < 0.001$) with plasma TAU more weakly yet significantly correlated ($\rho = -0.34$, $P < 0.001$) (Fig. 7A). Factor analysis of the peak blood biomarker concentrations and the corpus callosum FA was performed to assess the latent variable structure underlying these observations in a data-driven way (Fig. 7B). Three factors explained 71% of the variance in the data. Peak NfL and subacute FA in the corpus callosum both loaded heavily on the same factor, suggesting a close association of these measures (loading strengths 0.56 and -0.99, respectively). The other two factors were loaded variably by the blood biomarkers, but with little contribution of corpus callosum FA and lower NfL loading.

Diffusion tensor MRI was also performed in 12 of 18 patients who had undergone microdialysis (mean 25 days after injury; SD 12, Table S13). Across the group, FA was reduced relative to controls in the whole white matter ($W = 339$, $P < 0.001$, Table S13). More specifically, zFA from a

spherical region of interest around the catheter tip insertion site was also reduced ($V=0$, $P=0.003$, Table S13). Individual variability in catheter site FA and microdialysate biomarker concentrations did not show a significant correlation using Spearman's rank test, across NfL, TAU, UCH-L1 or GFAP (all $P>0.05$).

Validation of axonal injury markers in an experimental injury model

A controlled cortical impact (CCI) model was used to validate plasma NfL against neuroimaging and histopathological evidence of axonal injury. Other fluid biomarkers were not assessed in the rodent work. Thirty-four rats underwent either sham-operation ($n=4$), 1mm CCI ($n=15$, 'mild injury') or 2mm CCI ($n=15$, 'moderate-severe injury'). Ten naïve animals served as histological controls. Plasma NfL increased over two weeks following 1mm or 2mm injury. Mixed effects modelling with time and injury as fixed effects showed a significant interaction between time and injury severity ($F(4,90)=14.9$, $P<0.001$, Fig 8A). This was the result of similar plasma NfL concentrations at baseline, but higher plasma NfL at all post-injury timepoints in 1mm and 2mm CCI animals (all $P<0.001$, Fig. 8A, Table S14). Plasma NfL concentrations in sham/naïve animals did not change over time (Table S15). Cerebrospinal fluid was also sampled 15 days after CCI. CSF NfL was increased ($F(2,27)=24.2$, $P<0.001$) with higher concentrations in the 2mm group than the naïve/sham ($t=6.9$, $P<0.001$) and 1mm CCI animals ($t=5.0$, $P<0.001$, Fig. 8B, Table S14). Peak plasma NfL correlated with CSF NfL ($P=0.003$, $\rho=0.52$) (fig. S3).

White matter integrity was assessed in the corpus callosum (CC) using DTI at 14 days post-injury (Fig. 8C). Mean FA across the segments was significantly reduced ($P<0.05$) after injury in three 1mm CCI animals and eight 2mm CCI animals (Fig. 8D). Comparing zFA across CC segments between 1mm and 2mm CCI animals, a main effect of injury severity ($F(1,100)=28.3$, $P<0.001$) and segment ($F(4, 100) = 7.435$, $P<0.001$) was present, driven by lower FA in segment 4 for 2mm impacts ($P=0.001$, Fig. 8D). Plasma NfL concentrations were not significantly correlated with zFA ($P>0.05$, fig S3). Histopathological examination was performed at 15 days post-injury. Corpus callosum thickness appeared to be reduced (Fig. 8E), with a significant interaction between severity and segment of the corpus callosum ($F(8,80)=4.5$, $P<0.001$, Fig. 8F). Following 2mm impact, Luxol fast blue-positive tissue was markedly thinned proximally to the injury site (segments 2-4; all $P<0.02$, Table S15), but unchanged following 1mm or sham

injury (Fig. 8E,F). Mean CC thickness was positively correlated with zFA ($r=0.67$, $P=0.008$) and negatively correlated with peak plasma NfL ($\rho=-0.61$, $P=0.02$, fig. S3).

Axonal integrity was assessed using fluorescent neurofilament staining. Disorganised swollen axons and axonal bundles were visible after 2 mm, but not 1mm or sham injury, with axonal spheroid bulbs in CC segments proximal to injury (Fig. 8G). There was a significant main effect of injury severity ($F(2,80)=40.57$, $P<0.001$) and anatomical region ($F(4,80)=4.2$, $P=0.004$), with a reduction in NF staining in segments 2-5 in 2mm-CCI animals (all $P<0.04$, Table S15). Neurofilament stain intensity was negatively correlated with peak plasma NfL ($\rho=-0.48$, $P=0.042$) but not significantly correlated with zFA ($P>0.05$, fig. S4).

DISCUSSION

We present the results of BIO-AX-TBI, a multi-centre prospective study of moderate-severe TBI. Our goal was to investigate neuroimaging and fluid biomarkers of axonal injury, a key type of brain injury that remains difficult to diagnose using standard clinical tools. We focus on the assessment of neurofilament light (NfL), a promising marker of axonal damage. Our results show that plasma NfL concentrations are elevated in plasma after TBI. Variability in NfL plasma concentration is not explained by CT findings, as high concentrations are seen in patients with no visible pathology on CT. However, NfL correlates with the extent of axonal damage seen on diffusion MRI and predicts subsequent white matter neurodegeneration and late functional outcomes. We directly measure NfL from damaged white matter using invasive cerebral microdialysis. This shows that plasma NfL measurement reflects concentrations in damaged white matter within the brain. Microdialysate concentrations were around 100 times higher than plasma, with brain extracellular fluid and plasma concentrations strongly correlated. NfL also correlated with histopathologically-defined axonal injury within the white matter produced by an experimental injury model. TAU, GFAP, UCH-L1 and S100B were also elevated after TBI in plasma and brain extracellular fluid post-injury, and plasma TAU concentrations predicted cortical atrophy in the first six months after injury. Notably, concentrations of NfL peaked ten days to six weeks after TBI, in contrast to the acute peak seen for all other blood biomarkers assessed. The results show that plasma NfL is a sensitive and clinically-relevant measure of axonal injury, which might provide important prognostic information when measured subacutely post-TBI.

Neurofilament light has previously been studied in a range of neurological conditions characterized by axonal degeneration (7). NfL constitutes a core part of the axonal backbone in conjunction with higher molecular weight neurofilaments (medium and heavy), and cytoplasmic proteins including α -internexin and peripherin (32, 33). High expression is seen within the long myelinated axons of the cerebral deep white matter (16), which are susceptible to TAI that arises from shearing forces of rotational acceleration in head injury (15). Plasma and cerebrospinal fluid NfL increase after various types of TBI and previous studies have indicated the potential of NfL for predicting outcomes after single moderate/severe injury (8, 10, 13, 14). NfL concentrations are known to increase even after mild TBI (34, 35). Concentrations have previously been shown to be highest early after injury but to remain elevated in some individuals in the long term. NfL is elevated in a range of proteinopathies (7). Hence, NfL concentrations potentially provide information about the extent of post-traumatic neurodegeneration long after injury (8, 11, 36, 37).

Our work provides direct evidence that plasma NfL might be an informative indicator of TAI. Most importantly, we show that microdialysate taken directly from damaged white matter contains very high amounts of NfL. We used an innovative approach to microdialysis, with commercially available 3% Dextran 500 perfusion fluid, facilitating multi-centre microdialysis sampling. Ultrasensitive Simoa analysis of microdialysate and plasma were carried out on simultaneously acquired samples. Acute elevations of TAU, UCH-L1 and GFAP were also seen in microdialysate. Brain extracellular fluid concentrations of NfL, as well as TAU and UCH-L1, were around 100 times higher than plasma. Blood and brain concentrations of NfL, TAU and UCH-L1 were each highly correlated, indicating that plasma samples provide an accurate reflection post-traumatic changes in damaged white matter.

To our knowledge, GFAP has never previously been quantified in cerebral microdialysate. The difference in GFAP concentrations between plasma and microdialysate were much smaller than for other markers, reflecting relatively elevated plasma GFAP in the context of high extracellular fluid concentrations. In addition, there was no relationship between plasma and extracellular fluid GFAP concentrations, in contrast to the other biomarkers studied. The reasons for discrepant findings for GFAP are unclear, but the high plasma GFAP concentrations might reflect traumatic damage to astroglial endfeet within the blood brain barrier (BBB), which contain GFAP that would be expected to be released directly into the blood with BBB trauma (38). Hence, GFAP could provide a sensitive marker of BBB disruption after head injury, but further work is necessary to understand the relationship between GFAP, astroglial injury and BBB permeability after TBI.

The time-course of increased NfL in the acute and subacute period after TBI was distinct from other biomarkers. Plasma NfL peaked subacutely between ten days and six weeks post-injury. This contrasted with TAU, UCH-L1 and glial markers GFAP and S100B, which all peaked in the first 24 hours. This could be due to the longer half-life of NfL, which is estimated from mouse modelling to be around 3 weeks (39). Alternatively, NfL might also continue to be released from degenerating axons, a process that is known to progress for years after injury (40, 41). The delayed peak of NfL has important implications for optimally timing clinical measurement and measuring NfL 2-4 weeks after brain injury might be optimal for TAI diagnosis and prognostication. The differential time course across TAU and NfL may favour the development of a combined measure, incorporating the testing timepoint and a combination of early and later-peaking markers to optimally inform diagnosis and prognosis. Assessment of a further cohort with high frequency sampling, for example daily post-injury, would facilitate the development of such a tool.

Plasma NfL was closely related to diffusion MRI, which was used to quantify the extent of TAI. DTI provides a sensitive measure of axonal damage (15, 42), which can often be missed on CT or standard MRI (30). In keeping with previous work (5, 43), we observed widespread FA reductions after moderate/severe TBI. These abnormalities were seen in a large number of white matter tracts, including the corpus callosum where similar changes in FA were also seen in the experimental injury model. One previous study has shown that serum NfL at 30 days post-injury correlates with corpus callosum FA (8). We extend this work by showing that NfL measured subacutely correlates with diffusion MRI measures of TAI. This relationship was stronger for NfL than TAU, and a factor analysis provided further evidence that plasma NfL and DTI are closely related in quantifying underlying axonal injury. High concentrations of NfL were seen in patients irrespective of CT or MRI evidence of focal brain injury, showing that plasma NfL can be an informative indicator of traumatic axonal injury in patients without abnormalities on CT imaging. We found that higher concentrations of peak plasma NfL were associated with the occurrence of neurosurgery, even accounting for greater injury severity in the surgical group, measured using FA. Future work assessing plasma NfL in the context of elective surgeries unrelated to trauma will help to validate this finding and facilitate interpretation in TBI.

Traumatic axonal injury is a trigger for progressive neurodegeneration and NfL is a sensitive albeit non-specific marker of the proteinopathies present in a wide range of neurodegenerative pathologies (7, 44). Animal models show that increases in plasma NfL coincide with the appearance of TAU, amyloid beta and alpha synuclein pathology in the brain (44). We have previously shown that patterns of axonal injury measured by FA predict neurodegeneration in the chronic phase post-injury (20). NfL is used widely as a biomarker in neurodegenerative disease (7) and concentrations correlate with atrophy rates in Alzheimer's (45), and other diseases (46-48). Hence, we hypothesised that peak NfL early after TBI would predict late axonal degeneration because it would quantify the extent of early neurodegenerative insult. In our multi-center cohort, peak plasma NfL predicted late white matter neurodegeneration six to twelve months post-injury. Furthermore, diffusion abnormalities subacutely predicted spatial patterns of late white matter atrophy. Peak TAU plasma concentrations also predicted grey matter atrophy in the first six months after injury.

The results we report here, as well as our previous work (11), also show that plasma NfL can remain elevated in the chronic phase after TBI, which may reflect the presence of progressive neurodegenerative pathology. Increased plasma NfL correlates with microglial activation in white

matter damaged by traumatic axonal injury, suggesting that it provides a biomarker of the long-term post-traumatic pathology (11). Animal models also show that NfL can be used to track the effect of drugs that alter concentrations of TAU and amyloid pathology (44). Our previous work shows that minocycline given in the chronic phase after TBI produces transient increases in plasma NfL (11), which may be due to an inhibition of the restorative effects of chronically activated microglia (49, 50). Taken together these results suggest that plasma NfL is a promising marker for post-traumatic neurodegeneration that is characterised by progressive TAU and amyloid pathology (51). It may prove particularly useful for identifying patients with substantial post-traumatic neurodegeneration triggered by diffuse axonal injury and for the assessment of new treatments for post-traumatic neurodegeneration in early stage clinical trials (20, 51-54).

We went on to use animal modelling to confirm that isolated brain injury caused elevated plasma NfL in a dose-response manner. CCI modelling produced plasma and CSF NfL concentrations that increased with more severe injuries. These increases were accompanied by histopathological evidence of axonal injury and accompanying diffusion abnormalities within the corpus callosum seen in our human work (reduced FA). The animal work was conducted over two weeks post-injury, with imaging and CSF sampling at around 14 days. This timing paralleled our acute/subacute human studies, facilitating direct comparison. Two weeks after CCI there was evidence of axonal injury within the corpus callosum, with disorganised swollen axons, axonal bundles and axonal spheroid bulbs. We have previously reported related results in detail and shown how computational modelling of biomechanical strain produced at the time of injury predicts these histopathological changes and the accompanying diffusion abnormalities (15). In this study, NfL concentrations in CSF and plasma correlated with extent of axonal damage quantitatively, indicated by reduced white matter thickness and reduced neurofilament immunofluorescence staining in the callosum. Concentrations in plasma peaked around day 3 in the experimental model. This is in keeping with the human peak at ten days to six weeks, given the circa 10 times more rapid turnover of protein in rat (55).

Increased plasma NfL concentrations were seen in patients with trauma without TBI, although concentrations were 5-15 times higher in the TBI group than the non-TBI trauma group with a larger difference developing with increasing time since injury. This is a previously unappreciated finding. Whereas previous work has shown that additional extracranial injuries alongside TBI did not change the incremental value of NfL in predicting CT changes in TBI, biomarker changes in non-TBI patients had not specifically been assessed (56). The increase in NfL in non-TBI patients

might indicate that TBI had occurred, which was not apparent on our clinical assessment. However, we used stringent criteria to exclude even mild TBI from the non-TBI group. Vertebral fractures were present in some non-TBI patients and although there was no evidence of spinal cord injury in these cases, it is possible that subtle spinal cord injury may have been an NfL source (57). It is also possible that peripheral nerve injuries contributed to the NfL rise in our non-TBI group. We excluded patients with any clinically apparent peripheral sensory or motor abnormalities, but sub-clinical nerve damage could have resulted in a modest increase in NfL (21). Studies including patients with clearly defined peripheral nerve trauma, such as limb amputation, would help to further clarify this issue.

Limitations

There were several potential limitations of the study. One concern is the potential influence of focal post-traumatic lesions on the calculation of atrophy rates and diffusion MRI measures. To mitigate this, each scan was assessed for focal lesions, with custom ‘masks’ created to exclude these areas, with FA and JD measures only sampled from non-lesioned voxels. A further possible concern is inter-site variability, though we carefully aligned data collection and analysis, with scanner differences specifically considered. We imaged healthy age and sex-matched controls and used these to generate a normalised FA map for each patient per-site voxelwise. Volumetric longitudinal healthy control data were acquired at a single centre, compared with patient data which was collected in three different sites. As the atrophy rate calculation uses an individual’s scan pair, each patient effectively acts as their own control, so we do not expect scanner differences to influence atrophy rates. All blood and microdialysate samples were centrally analysed using the same test kits and reagent batches to further minimise cross-site variability and facilitate comparisons.

In summary, we report the prospective BIO-AX-TBI study of advanced fluid and imaging markers of axonal injury after moderate-severe TBI. Combining human data with experimental modelling, we show that NfL is released from white matter proportionately to injury severity, and that it can be accurately assessed in blood. Plasma NfL correlated with FA on DTI MRI, an established neuroimaging biomarker of axonal injury, and when assessed subacutely, predicts chronic white matter atrophy and functional outcomes at six and twelve months post injury in our cohort. Total TAU predicted grey matter atrophy rates after injury, providing complementary information to NfL. We conclude that plasma NfL measured subacutely after TBI might reflect axonal injury

within the brain. These longitudinal dynamics lend themselves to clinical assessment of individuals with persistent symptoms after injury who consult clinicians in the outpatient setting, and where there may be uncertainty about whether any relevant axonal damage was produced at the time of injury.

MATERIALS AND METHODS

Study design

BIO-AX-TBI is a prospective multi-centre cohort study of patients after TBI (28). The aim of the study was to characterize the time course of NfL in addition to the axonal marker tau and associated proteins UCH-L1, GFAP and S100B, and to relate these to MRI measures of axonal injury and outcomes in adult moderate-severe TBI patients. We examined how measures were correlated, and whether early plasma biomarker levels contribute to the prediction of functional outcomes. In a sub-group of patients, we also combined microdialysis, neuroimaging and plasma sampling of axonal proteins to investigate the axonal origin of the plasma biomarkers and gain understanding of the mechanisms of axonal injury progression. A controlled cortical impact injury model was performed, aligned to BIO-AX-TBI, comprising analyses of neurofilament light, neuroimaging and histopathology to provide ground-truth validation. Imaging assessment and plasma biomarkers were available longitudinally in a subset of individuals with TBI. The sample size was based on the number (140 patients, 12 controls per study site) of participants needed to reliably test the contribution of DTI to prognostic modelling (type 1 error=0.05, power=0.95). Patient data from the BIO-AX-TBI study was analysed once the required number of patients with DTI and acute/subacute blood biomarkers had been followed-up at 6 months (Fig.1A, N=146 in the final analysis group) (28). All analyses were conducted blinded to group allocation or patient ID.

Prospective cohort study overview

Patients with moderate-severe TBI, defined by the Mayo classification (58), were recruited from eight trauma centres. The Mayo classification uses several measures to determine injury severity, making the tool more resilient to uncertainty or missing data in certain domains. Any injury with CT or MRI evidence of intraparenchymal injury eg. contusion or penetrating injury, extradural, subdural haematoma or subarachnoid haemorrhage is classified as moderate-severe. In addition, any patients meeting clinical criteria such as death due to injury, post traumatic amnesia lasting more than 24 hours, loss of consciousness lasting more than 30 minutes or lowest GCS < 14. The classification does not sub-divide the moderate-severe group and only a single criterion needs to be met for an injury to be within the overarching moderate/severe category.

Patients unable to provide consent underwent an assent process and were re-consented if they were subsequently regained capacity. Inclusion criteria were age between 18 to 80 at time of injury and a diagnosis of moderate-severe TBI. Exclusion criteria were pre-existing neurological disease, previous TBI requiring hospitalisation, relevant drug or alcohol abuse, pregnancy and for the MRI component, the presence of typical contraindications such as ferrous implants.

Clinical and demographic information were collected electronically using TBI common data elements. Blood biomarkers were assessed twice in the first ten days following injury. Patients were followed-up at a subacute timepoint ten days to six weeks post-injury. Follow-up visits included blood, MRI and clinical assessment (Fig. 1B). Additional visits were performed at six and twelve months post-injury with a functional assessment including the GOSE (59). In a subset of sites (Fig. 1B), six and twelve month assessments included blood and neuroimaging. Cerebral microdialysis was performed acutely at two sites, in patients with GCS of 3-8, evidence of trauma on CT and undergoing craniotomy or craniectomy facilitating catheter insertion. In these individuals, plasma biomarker assessment was performed up to twice per day acutely.

Healthy controls were recruited at each study site according to the same exclusion criteria. Controls underwent a single MRI and blood biomarker assessment. A separate group of longitudinally-scanned controls were included, comprising healthy volunteers who had participated in more than one study on the same scanner system with an aligned acquisition protocol (52).

Non-TBI patients were recruited at one site (St Mary's Hospital, London) and underwent blood biomarker assessment on two occasions in the first ten days following injury. Inclusion criteria were the extracranial injuries requiring hospitalisation. As with the main patient group, exclusions included premorbid substantial neurological disease in addition to any suspicion of TBI (for example symptomatic possible injuries), or a diagnosis of spinal cord injury, specific to this group. Participants were recruited over the following time windows: patients with TBI from November 2017 to July 2020, healthy volunteers from April 2018 to July 2019 and patients with NTT from January 2019 to September 2019.

Fluid Biomarker Analysis

Plasma samples were analysed for NfL, TAU, GFAP and UCH-L1 using a Simoa-HD1 platform. Serum S100B was measured using a Millipore ELISA kit (Supplementary Materials). Microdialysis samples were analysed in a similar manner on a Simoa-HD1 platform (see Supplementary Materials).

Statistical analysis

Standard t-tests were used for comparisons of normally distributed continuous data, and the non-parametric Wilcoxon test for non-normally distributed continuous variables. Paired t-tests, repeated measures ANOVA designs or linear mixed effects models, were used for longitudinal within-subject data, as appropriate. Owing to the data distribution, linear regressions on fluid biomarker concentrations were performed on $\log_{10}$ transformed data. Bonferroni multiple comparison corrections were performed to correct for the number of timepoints and biomarkers/groups assessed, other than for NfL where a clear a-priori hypothesis existed. Hence, NfL comparisons were corrected for the groups/timepoints only. For biomarker analyses, mean results of samples measured in duplicate by a technician using the same batch of reagents blinded to clinical information were used. Peak biomarker concentrations within an individual were defined as the highest concentration over 6 weeks post-injury. In exploratory analysis of other biomarker concentrations and within the experimental injury model, a correction for the number of biomarkers was conducted. Factor analyses (R core 'factanal' function, maximum likelihood method, using varimax rotation to explain the maximum possible variance) were performed using the peak concentrations of each fluid biomarker and the corpus callosum zFA subacutely, identifying a latent variable structure across three factors. In the model variance was spread evenly across the factors. Chi squared tests of sufficiency for models with one, or two factors only showed that these would be inadequate. Unless otherwise stated, all boxplots all show median, hinges at 25th and 75th centiles, and whiskers extending to the largest value less than or equal to 1.5 times the interquartile range from the hinge. To assess the relationships between biomarker concentrations in blood and cerebral microdialysate, a linear mixed effects model was used requiring complete data sampled in plasma/brain at aligned timepoints. Two values were identified missing in two patients (a GFAP result and UCH-L1 result). Using the markers surrounding the concentration and the time since injury as features vectors, they were imputed independently using a sklearn distance-based k nearest neighbours imputer (k=2). To account for temporal misalignment between brain and blood markers, we first up-sampled the brain markers using a

time-weighted interpolation model. We then removed any missing paired observations in either the beginning or the end of the matching sequence. Finally, redundant brain observations were omitted to match the sampling frequency in the blood. In the model for each biomarker, subject was a fixed effect, with main effects of brain concentration, and an interaction of log-transformed brain concentration and time since injury, to predict log-transformed plasma concentration.

Neuroimaging statistical analyses - diffusion

Voxelwise analysis of zFA was cluster corrected using TFCE, with multiple-comparisons correction using a family-wise error rate of $P < 0.05$. Longitudinal and cross sectional analysis of diffusion metrics was conducted using the general linear model with non-parametric permutation testing (10,000) in FSL Randomise (60), with age and gender included as a nuisance covariates in cross sectional analyses, and individualised lesion masking. In patients with microdialysis, the catheter tip was identified on CT imaging. CT scans were registered to DTITK space and a 10mm radius spherical region of interest was generated. The spherical region of interest was overlaid on the white matter skeleton and FA sampled within the overlapping voxels.

Neuroimaging statistical analyses - Structural

Voxelwise analyses of longitudinal volumetric imaging data were conducted using non-parametric permutation testing (10,000) with age and sex as nuisance covariates, with individualised lesion masking. Given the strong prior hypothesis of the direction of change (atrophy post-TBI), we report a one-sided t test for this analysis. Voxelwise correlation between JD and DTI metrics were conducted after registration of DTI images to MNI152 space, using non-parametric permutation testing ($n=10,000$).

Experimental injury model statistical analyses

A one-way ANOVA was used to test differences in plasma NfL concentration at baseline and CSF NfL concentrations. A mixed-effects model, with time and injury as fixed effects, were used to test differences in plasma NfL concentrations across different animal injuries and time since injury. CSF NfL concentrations was not different between naïve animals and shams, and hence these groups were combined as controls. DTI changes were analysed using a mixed effects model with injury and segment as fixed effects. White matter integrity, as shown by LFB staining, and

neurofilament expression pattern and density, as indicated by fluorescence intensity, was assessed using a two-way ANOVA with segment and impact severity as factors.

Supplementary Materials

Supplementary Methods

Sensitivity Analysis

Table S1. Participant demographics

Table S2. Clinical characteristics of TBI patients.

Table S3. Characteristics of non-TBI trauma patients

Table S4. Blood biomarkers after TBI, extracranial injuries and in healthy controls

Table S5. Blood biomarker concentrations: lesion related analysis.

Table S6. Blood biomarker concentrations: neurosurgery analysis.

Table S7. Blood biomarker concentrations: neurosurgery analysis accounting for white matter FA.

Table S8. Blood biomarker concentrations: age-related sensitivity analysis.

Table S9. Cerebral microdialysis: patient characteristics and biomarker trends

Table S10. Biomarker concentrations in microdialysate and plasma

Table S11. MRI measures of white matter integrity and progressive atrophy

Table S12. Relationship of blood biomarkers and long-term outcomes after TBI.

Table S13. Cerebral Microdialysis: DTI imaging measures

Table S14. Blood and imaging markers in an experimental injury model

Table S15. Plasma biomarkers after experimental injury: mixed effects model results.

Table S16. Imaging acquisition parameters and scanner details

Fig. S1. Spatial pattern of MRI-defined focal lesions in patients TBI.

Fig. S2. No significant relationship between age at injury and plasma NfL concentration at six and twelve months post-TBI

Fig. S3. Individual fluid biomarker trends in cerebral microdialysate and blood

Fig. S4. Experimental injury model fluid biomarker, neuroimaging and histology relationships.

Fig. S5. Neuroimaging acquisition and analysis pipelines.

Fig. S6. Experimental injury model methods

Contributor list. The BIOAXTBI Group

Supplementary References

References and notes:

Acknowledgments:

We would like to thank the patients and their families for their involvement in this research, in addition to our funders for supporting the project.

Funding:

ERA-NET NEURON Cofund (MR/R004528/1), part of the European Research Projects on External Insults to the Nervous System call, within the Horizon 2020 funding framework, provided the core funds for the project (PIs: DJS, HZ, MO, GB, SM)
Alzheimer's Research UK Clinical Research Fellowship (NG)
UK Dementia Research Institute (DRI) Care Research and Technology Centre (NG/DJS/KZ)
National Institute of Health Research (NIHR) Professorship (NIHR-RP-011-048) (DJS)
The NIHR Clinical Research Facility and Biomedical Research Centre (BRC) at Imperial College Healthcare NHS Trust (DJS)
Medical Research Council Clinician Scientist Fellowship (DJS)
Wallenberg Scholarship (HZ)
Swedish Research Council (#2018-02532) (HZ)
The European Research Council (#681712) (HZ)
Swedish State Support for Clinical Research (#ALFGBG-720931) (HZ)
Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862) (HZ)
The European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860197 (MIRIADE) (HZ)
The UK Dementia Research Institute at UCL (HZ)
Swiss National Science Foundation (#31NE30_173675, #32003B_188501) (MO)

Author contributions:

Writing – original draft: NSNG, KAZ, CKD, DJS
Data curation and centralisation: EG and GB
Imaging data curation: NJB
Experimental injury work: CKD and MYL
Core data analysis: NSNG and KAZ.
Neuroimaging review: MCP, DR and GF
Design and performance of cerebral microdialysis: SAB, AB, JM, MO, PG and SM
Fluid biomarker analyses: AH and HZ
Microdialysis modelling: ES
Writing – reviewing and editing: NSNG, KAZ, FM, AH, SAM, AB, JM, CKD, MY, NB, AEJ, EM, ES, MHW, GF, DR, MCP, EG, GN, CB, EF, AC, PG, SM, MO, HZ, GB, DJS

Competing interests: HZ has served at scientific advisory boards for Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics and CogRx, has given lectures in symposia sponsored by Fujirebio, Alzecure and Biogen, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures

Incubator Program (outside submitted work). MO received research support and speaker fees from Neuroptics, USA, unrelated to the present work, and is member of the Scientific Advisory Board of Neuroptics.

Data and materials availability: All the data are present in the main text or in the supplementary material file. Raw, unprocessed data are available from the corresponding author upon reasonable request and following the completion of a material transfer agreement (MTA) with Imperial College London.

References

1. A. I. R. Maas, D. K. Menon, P. D. Adelson, N. Andelic, M. J. Bell, A. Belli, P. Bragge, A. Brazinova, A. Buki, R. M. Chesnut, G. Citerio, M. Coburn, D. J. Cooper, A. T. Crowder, E. Czeiter, M. Czosnyka, R. Diaz-Arrastia, J. P. Dreier, A. C. Duhaime, A. Ercole, T. A. van Essen, V. L. Feigin, G. Gao, J. Giacino, L. E. Gonzalez-Lara, R. L. Gruen, D. Gupta, J. A. Hartings, S. Hill, J. Y. Jiang, N. Ketharanathan, E. J. O. Kompanje, L. Lanyon, S. Laureys, F. Lecky, H. Levin, H. F. Lingsma, M. Maegele, M. Majdan, G. Manley, J. Marsteller, L. Mascia, C. McFadyen, S. Mondello, V. Newcombe, A. Palotie, P. M. Parizel, W. Peul, J. Piercy, S. Polinder, L. Puybasset, T. E. Rasmussen, R. Rossaint, P. Smielewski, J. Soderberg, S. J. Stanworth, M. B. Stein, N. von Steinbuchel, W. Stewart, E. W. Steyerberg, N. Stocchetti, A. Synnot, B. Te Ao, O. Tenovuo, A. Theadom, D. Tibboel, W. Videtta, K. K. W. Wang, W. H. Williams, L. Wilson, K. Yaffe, InTBIR Participants and Investigators, Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol* **16**, 987-1048 (2017).
2. J. Ponsford, K. Draper, M. Schönberger, Functional outcome 10 years after traumatic brain injury: its relationship with demographic, injury severity, and cognitive and emotional status. *J Int Neuropsychol Soc* **14**, 233-242 (2008).
3. L. Whitnall, T. M. McMillan, G. D. Murray, G. M. Teasdale, Disability in young people and adults after head injury: 5-7 year follow up of a prospective cohort study. *J Neurol Neurosurg Psychiatry* **77**, 640-645 (2006).
4. D. H. Smith, D. F. Meaney, W. H. Shull, Diffuse axonal injury in head trauma. *J Head Trauma Rehabil* **18**, 307-316 (2003).
5. A. Sidaros, A. W. Engberg, K. Sidaros, M. G. Liptrot, M. Herning, P. Petersen, O. B. Paulson, T. L. Jernigan, E. Rostrup, Diffusion tensor imaging during recovery from severe traumatic brain injury and relation to clinical outcome: a longitudinal study. *Brain* **131**, 559-572 (2008).
6. H. Zetterberg, D. H. Smith, K. Blennow, Biomarkers of mild traumatic brain injury in cerebrospinal fluid and blood. *Nat Rev Neurol* **9**, 201-210 (2013).
7. L. Gaetani, K. Blennow, P. Calabresi, M. Di Filippo, L. Parnetti, H. Zetterberg, Neurofilament light chain as a biomarker in neurological disorders. *Journal of Neurology, Neurosurgery & Psychiatry* **90**, 870-881 (2019).
8. P. Shahim, A. Politis, A. van der Merwe, B. Moore, Y.-Y. Chou, D. L. Pham, J. A. Butman, R. Diaz-Arrastia, J. M. Gill, D. L. Brody, H. Zetterberg, K. Blennow, L. Chan, Neurofilament light as a biomarker in traumatic brain injury. *Neurology* **95**, e610-e622 (2020).
9. M. Gisslén, R. W. Price, U. Andreasson, N. Norgren, S. Nilsson, L. Hagberg, D. Fuchs, S. Spudich, K. Blennow, H. Zetterberg, Plasma Concentration of the Neurofilament Light Protein (NFL) is a Biomarker of CNS Injury in HIV Infection: A Cross-Sectional Study. *EBioMedicine* **3**, 135-140 (2016).
10. P. Shahim, M. Gren, V. Liman, U. Andreasson, N. Norgren, Y. Tegner, N. Mattsson, N. Andreasen, M. Ost, H. Zetterberg, B. Nellgard, K. Blennow, Serum neurofilament light protein predicts clinical outcome in traumatic brain injury. *Scientific reports* **6**, 36791 (2016).
11. G. Scott, H. Zetterberg, A. Jolly, J. H. Cole, S. De Simoni, P. O. Jenkins, C. Feeney, D. R. Owen, A. Lingford-Hughes, O. Howes, M. C. Patel, A. P. Goldstone, R. N. Gunn, K. Blennow, P. M. Matthews, D. J. Sharp, Minocycline reduces chronic microglial activation after brain trauma but increases neurodegeneration. *Brain* **141**, 459-471 (2018).

12. E. P. Thelin, F. A. Zeiler, A. Ercole, S. Mondello, A. Buki, B. M. Bellander, A. Helmy, D. K. Menon, D. W. Nelson, Serial Sampling of Serum Protein Biomarkers for Monitoring Human Traumatic Brain Injury Dynamics: A Systematic Review. *Front Neurol* **8**, 300 (2017).
13. E. Thelin, F. Al Nimer, A. Frostell, H. Zetterberg, K. Blennow, H. Nyström, M. Svensson, B. M. Bellander, F. Piehl, D. W. Nelson, A Serum Protein Biomarker Panel Improves Outcome Prediction in Human Traumatic Brain Injury. *J Neurotrauma* **36**, 2850-2862 (2019).
14. F. Al Nimer, E. Thelin, H. Nyström, A. M. Dring, A. Svenningsson, F. Piehl, D. W. Nelson, B.-M. Bellander, Comparative Assessment of the Prognostic Value of Biomarkers in Traumatic Brain Injury Reveals an Independent Role for Serum Levels of Neurofilament Light. *PLOS ONE* **10**, e0132177 (2015).
15. C. K. Donat, M. Yanez-Lopez, M. Sastre, N. Baxan, M. Goldfinger, R. Seamber, F. Müller, P. Davies, P. Hellyer, P. Siegkas, S. Gentleman, D. J. Sharp, M. Ghajari, From biomechanics to pathology: predicting axonal injury from patterns of strain after traumatic brain injury. *Brain* **144**, 70-91 (2021).
16. W. W. Schlaepfer, R. G. Lynch, Immunofluorescence studies of neurofilaments in the rat and human peripheral and central nervous system. *J Cell Biol* **74**, 241-250 (1977).
17. S. Magnoni, C. L. Mac Donald, T. J. Esparza, V. Conte, J. Sorrell, M. Macri, G. Bertani, R. Biffi, A. Costa, B. Sammons, A. Z. Snyder, J. S. Shimony, F. Triulzi, N. Stocchetti, D. L. Brody, Quantitative assessments of traumatic axonal injury in human brain: concordance of microdialysis and advanced MRI. *Brain* **138**, 2263-2277 (2015).
18. S. Magnoni, T. J. Esparza, V. Conte, M. Carbonara, G. Carrabba, D. M. Holtzman, G. J. Zipfel, N. Stocchetti, D. L. Brody, Tau elevations in the brain extracellular space correlate with reduced amyloid-beta levels and predict adverse clinical outcomes after severe traumatic brain injury. *Brain* **135**, 1268-1280 (2012).
19. C. L. Mac Donald, K. Dikranian, P. Bayly, D. Holtzman, D. Brody, Diffusion Tensor Imaging Reliably Detects Experimental Traumatic Axonal Injury and Indicates Approximate Time of Injury. *The Journal of Neuroscience* **27**, 11869 (2007).
20. N. S. N. Graham, A. Jolly, K. Zimmerman, N. J. Bourke, G. Scott, J. H. Cole, J. M. Schott, D. J. Sharp, Diffuse axonal injury predicts neurodegeneration after moderate-severe traumatic brain injury. *Brain*, (2020).
21. Å. Sandelius, H. Zetterberg, K. Blennow, R. Adiutori, A. Malaspina, M. Laura, M. M. Reilly, A. M. Rossor, Plasma neurofilament light chain concentration in the inherited peripheral neuropathies. *Neurology* **90**, e518-e524 (2018).
22. L. Martín-Aguilar, P. Camps-Renom, C. Lleixà, E. Pascual-Goñi, J. Díaz-Manera, R. Rojas-García, N. De Luna, E. Gallardo, E. Cortés-Vicente, L. Muñoz, D. Alcolea, A. Lleó, C. Casasnovas, C. Homedes, G. Gutiérrez-Gutiérrez, M. C. Jimeno-Montero, J. Berciano, M. J. Sedano-Tous, T. García-Sobrino, J. Pardo-Fernández, C. Márquez-Infante, I. Rojas-Marcos, I. Jericó-Pascual, E. Martínez-Hernández, G. Moris de la Tassa, C. Domínguez-González, I. Illa, L. Querol, Serum neurofilament light chain predicts long-term prognosis in Guillain-Barré syndrome patients. *Journal of Neurology, Neurosurgery & Psychiatry* **92**, 70-77 (2021).
23. R. E. Anderson, L. O. Hansson, O. Nilsson, R. Djalil-Merzoug, G. Settergren, High serum S100B levels for trauma patients without head injuries. *Neurosurgery* **48**, 1255-1258; discussion 1258-1260 (2001).
24. C. K. Donat, G. Scott, S. M. Gentleman, M. Sastre, Microglial Activation in Traumatic Brain Injury. *Front Aging Neurosci* **9**, 208 (2017).
25. C. Edward Dixon, G. L. Clifton, J. W. Lighthall, A. A. Yaghmai, R. L. Hayes, A controlled cortical impact model of traumatic brain injury in the rat. *Journal of Neuroscience Methods* **39**, 253-262 (1991).
26. W. H. Cheng, S. Stukas, K. M. Martens, D. R. Namjoshi, E. B. Button, A. Wilkinson, A. Bashir, J. Robert, P. A. Crompton, C. L. Wellington, Age at injury and genotype modify acute inflammatory and neurofilament-light responses to mild CHIMERA traumatic brain injury in wild-type and APP/PS1 mice. *Experimental Neurology* **301**, 26-38 (2018).
27. A. Bashir, Z. A. Abebe, K. A. McInnes, E. B. Button, I. Tatarnikov, W. H. Cheng, M. Haber, A. Wilkinson, C. Barron, R. Diaz-Arrastia, S. Stukas, P. A. Crompton, C. L. Wellington, Increased severity of the CHIMERA model induces acute vascular injury, sub-acute deficits in memory recall, and chronic white matter gliosis. *Experimental neurology*. **324**, 113116 (2020).
28. N. S. Graham, K. A. Zimmerman, G. Bertolini, S. Magnoni, M. Oddo, H. Zetterberg, F. Moro, D. Novelli, A. Heslegrave, A. Chierigato, E. Fainardi, J. M. Fleming, E. Garbero, S. Abed-Maillard, P. Gradisek, A. Bernini, D. J. Sharp, Multicentre longitudinal study of fluid and neuroimaging BIOMarkers of AXonal injury after traumatic brain injury: the BIO-AX-TBI study protocol. *BMJ Open* **10**, e042093 (2020).
29. L. Squarcina, A. Bertoldo, T. E. Ham, R. Heckemann, D. J. Sharp, A robust method for investigating thalamic white matter tracts after traumatic brain injury. *Neuroimage* **63**, 779-788 (2012).

30. A. E. Jolly, M. Balaet, A. Azor, D. Friedland, S. Sandrone, N. S. N. Graham, K. Zimmerman, D. J. Sharp, Detecting axonal injury in individual patients after Traumatic Brain Injury. *Brain* **144**, 92-113 (2021).
31. D. M. Cash, C. Frost, L. O. Iheme, D. Unay, M. Kandemir, J. Fripp, O. Salvado, P. Bourgeat, M. Reuter, B. Fischl, M. Lorenzi, G. B. Frisoni, X. Pennec, R. K. Pierson, J. L. Gunter, M. L. Senjem, C. R. Jack, Jr., N. Guizard, V. S. Fonov, D. L. Collins, M. Modat, M. J. Cardoso, K. K. Leung, H. Wang, S. R. Das, P. A. Yushkevich, I. B. Malone, N. C. Fox, J. M. Schott, S. Ourselin, Assessing atrophy measurement techniques in dementia: Results from the MIRIAD atrophy challenge. *Neuroimage* **123**, 149-164 (2015).
32. A. R. Gafson, N. R. Barthélemy, P. Bomont, R. O. Carare, H. D. Durham, J.-P. Julien, J. Kuhle, D. Leppert, R. A. Nixon, R. O. Weller, H. Zetterberg, P. M. Matthews, Neurofilaments: neurobiological foundations for biomarker applications. *Brain: A Journal of Neurology*, (2020).
33. P. N. Hoffman, D. W. Cleveland, J. W. Griffin, P. W. Landes, N. J. Cowan, D. L. Price, Neurofilament gene expression: a major determinant of axonal caliber. *Proc Natl Acad Sci U S A* **84**, 3472-3476 (1987).
34. H. Zetterberg, M. A. Hietala, M. Jonsson, N. Andreasen, E. Styrd, I. Karlsson, A. Edman, C. Popa, A. Rasulzada, L. O. Wahlund, P. D. Mehta, L. Rosengren, K. Blennow, A. Wallin, Neurochemical aftermath of amateur boxing. *Arch Neurol* **63**, 1277-1280 (2006).
35. S. Neselius, H. Brisby, A. Theodorsson, K. Blennow, H. Zetterberg, J. Marcusson, CSF-biomarkers in Olympic boxing: diagnosis and effects of repetitive head trauma. *PLoS One* **7**, e33606 (2012).
36. C. Bernick, G. Shan, H. Zetterberg, S. Banks, V. R. Mishra, L. Bekris, J. B. Leverenz, K. Blennow, Longitudinal change in regional brain volumes with exposure to repetitive head impacts. *Neurology* **94**, e232-e240 (2020).
37. P. Shahim, Y. Tegner, B. Gustafsson, M. Gren, J. Ärlig, M. Olsson, N. Lehto, Å. Engström, K. Höglund, E. Portelius, H. Zetterberg, K. Blennow, Neurochemical Aftermath of Repetitive Mild Traumatic Brain Injury. *JAMA Neurology* **73**, 1308-1315 (2016).
38. M. Simard, G. Arcuino, T. Takano, Q. S. Liu, M. Nedergaard, Signaling at the gliovascular interface. *J Neurosci* **23**, 9254-9262 (2003).
39. S. Millicamps, G. Gowing, O. Corti, J. Mallet, J.-P. Julien, Conditional NF-L Transgene Expression in Mice form In Vivo Analysis of Turnover and Transport Rate of Neurofilaments. *The Journal of Neuroscience* **27**, 4947-4956 (2007).
40. W. W. Schlaepfer, Calcium-induced degeneration of axoplasm in isolated segments of rat peripheral nerve. *Brain Res* **69**, 203-215 (1974).
41. M. E. Vargas, B. A. Barres, Why is Wallerian degeneration in the CNS so slow? *Annu Rev Neurosci* **30**, 153-179 (2007).
42. C. L. Mac Donald, K. Dikranian, S. K. Song, P. V. Bayly, D. M. Holtzman, D. L. Brody, Detection of traumatic axonal injury with diffusion tensor imaging in a mouse model of traumatic brain injury. *Exp Neurol* **205**, 116-131 (2007).
43. K. M. Kinnunen, R. Greenwood, J. H. Powell, R. Leech, P. C. Hawkins, V. Bonnelle, M. C. Patel, S. J. Counsell, D. J. Sharp, White matter damage and cognitive impairment after traumatic brain injury. *Brain* **134**, 449-463 (2011).
44. M. Bacioglu, L. F. Maia, O. Preische, J. Schelle, A. Apel, S. A. Kaeser, M. Schweighauser, T. Eninger, M. Lambert, A. Pilotto, D. R. Shimshek, U. Neumann, P. J. Kahle, M. Staufenbiel, M. Neumann, W. Maetzler, J. Kuhle, M. Jucker, Neurofilament Light Chain in Blood and CSF as Marker of Disease Progression in Mouse Models and in Neurodegenerative Diseases. *Neuron* **91**, 56-66 (2016).
45. N. Mattsson, U. Andreasson, H. Zetterberg, K. Blennow, Association of Plasma Neurofilament Light With Neurodegeneration in Patients With Alzheimer Disease. *JAMA Neurol* **74**, 557-566 (2017).
46. E. B. Johnson, L. M. Byrne, S. Gregory, F. B. Rodrigues, K. Blennow, A. Durr, B. R. Leavitt, R. A. Roos, H. Zetterberg, S. J. Tabrizi, R. I. Schill, E. J. Wild, Neurofilament light protein in blood predicts regional atrophy in Huntington disease. *Neurology* **90**, e717-e723 (2018).
47. J. D. Rohrer, I. O. Woollacott, K. M. Dick, E. Brotherhood, E. Gordon, A. Fellows, J. Toombs, R. Drueyeh, M. J. Cardoso, S. Ourselin, J. M. Nicholas, N. Norgren, S. Mead, U. Andreasson, K. Blennow, J. M. Schott, N. C. Fox, J. D. Warren, H. Zetterberg, Serum neurofilament light chain protein is a measure of disease intensity in frontotemporal dementia. *Neurology* **87**, 1329-1336 (2016).
48. C. Barro, P. Benkert, G. Disanto, C. Tsagkas, M. Amann, Y. Naegelin, D. Leppert, C. Gobbi, C. Granziera, Ö. Yaldizli, Z. Michalak, J. Wuerfel, L. Kappos, K. Parmar, J. Kuhle, Serum neurofilament as a predictor of disease worsening and brain and spinal cord atrophy in multiple sclerosis. *Brain* **141**, 2382-2391 (2018).
49. K. Nagamoto-Combs, D. W. McNeal, R. J. Morecraft, C. K. Combs, Prolonged microgliosis in the rhesus monkey central nervous system after traumatic brain injury. *J Neurotrauma* **24**, 1719-1742 (2007).

50. K. Nagamoto-Combs, R. J. Morecraft, W. G. Darling, C. K. Combs, Long-term gliosis and molecular changes in the cervical spinal cord of the rhesus monkey after traumatic brain injury. *J Neurotrauma* **27**, 565-585 (2010).
51. N. S. Graham, D. J. Sharp, Understanding neurodegeneration after traumatic brain injury: from mechanisms to clinical trials in dementia. *J Neurol Neurosurg Psychiatry*, (2019).
52. J. H. Cole, A. Jolly, S. de Simoni, N. Bourke, M. C. Patel, G. Scott, D. J. Sharp, Spatial patterns of progressive brain volume loss after moderate-severe traumatic brain injury. *Brain* **141**, 822-836 (2018).
53. A. F. Ramlackhansingh, D. J. Brooks, R. J. Greenwood, S. K. Bose, F. E. Turkheimer, K. M. Kinnunen, S. Gentleman, R. A. Heckemann, K. Gunanayagam, G. Gelosa, D. J. Sharp, Inflammation after trauma: microglial activation and traumatic brain injury. *Ann Neurol* **70**, 374-383 (2011).
54. V. E. Johnson, J. E. Stewart, F. D. Begbie, J. Q. Trojanowski, D. H. Smith, W. Stewart, Inflammation and white matter degeneration persist for years after a single traumatic brain injury. *Brain* **136**, 28-42 (2013).
55. D. V. Agoston, How to Translate Time? The Temporal Aspect of Human and Rodent Biology. *Frontiers in Neurology* **8**, (2017).
56. E. Czeiter, K. Amrein, B. Y. Gravesteyn, F. Lecky, D. K. Menon, S. Mondello, V. F. J. Newcombe, S. Richter, E. W. Steyerberg, T. V. Vyvere, J. Verheyden, H. Xu, Z. Yang, A. I. R. Maas, K. K. W. Wang, A. Büki, Blood biomarkers on admission in acute traumatic brain injury: Relations to severity, CT findings and care path in the CENTER-TBI study. *EBioMedicine* **56**, (2020).
57. J. Kuhle, J. Gaiottino, D. Leppert, A. Petzold, J. P. Bestwick, A. Malaspina, C. H. Lu, R. Dobson, G. Disanto, N. Norgren, A. Nissim, L. Kappos, J. Hurlbert, V. W. Yong, G. Giovannoni, S. Casha, Serum neurofilament light chain is a biomarker of human spinal cord injury severity and outcome. *J Neurol Neurosurg Psychiatry* **86**, 273-279 (2015).
58. J. F. Malec, A. W. Brown, C. L. Leibson, J. T. Flaada, J. N. Mandrekar, N. N. Diehl, P. K. Perkins, The mayo classification system for traumatic brain injury severity. *J Neurotrauma* **24**, 1417-1424 (2007).
59. J. T. Wilson, L. E. Pettigrew, G. M. Teasdale, Structured interviews for the Glasgow Outcome Scale and the extended Glasgow Outcome Scale: guidelines for their use. *J Neurotrauma* **15**, 573-585 (1998).
60. A. M. Winkler, G. R. Ridgway, M. A. Webster, S. M. Smith, T. E. Nichols, Permutation inference for the general linear model. *Neuroimage* **92**, 381-397 (2014).
61. L. Siebold, A. Obenaus, R. Goyal, Criteria to define mild, moderate, and severe traumatic brain injury in the mouse controlled cortical impact model. *Exp Neurol* **310**, 48-57 (2018).

Figure legends

Fig. 1. Study design. (A) BIO-AX-TBI patient recruitment and retention. Univ: university; Hosp: hospital. (B) Assessments: typical patient pathway (top), centres performing additional advanced repeated neuroimaging and longitudinal blood biomarkers (middle), CT is performed clinically at all sites, and invasive cerebral microdialysis at two sites (lower). (C) Experimental injury model at London site.

Fig. 2. Demographics, Clinical Characteristics and Outcomes. (A) * Age and sex of patients after traumatic brain injury (TBI), healthy controls (HC) and with non-TBI-trauma (NTT). TBI patients were older in age than healthy controls ($W=10784$, $P=0.028$). *, **, *** indicates a significance of $P<0.05$, $P<0.01$ and $P<0.001$ compared to controls. (B) Relationship of age at injury to hospital admission duration and mortality (red:died, blue:survived). (C) Acute Glasgow Coma Scale (GCS) after TBI. (D) Pupillary assessment in emergency department (ED). N: normal; Uni dil.: unilaterally dilated; Bilat dil: bilaterally dilated. (E) Marshall classification of most severe CT in first day. (F) MRI pathologies subacutely (ie. at ten days to six weeks). (G) Glasgow Outcome Score Extended (GOSE) at six and twelve months post-TBI.

Fig. 3. Blood biomarker trends in TBI, non-TBI trauma and healthy controls. (A) Trends of blood NfL, TAU, UCH-L1, GFAP and S100B concentrations after TBI. (B) Local polynomial (LOESS) regression curves modelling concentrations in the first 40 days post-TBI. (C) Group comparisons of biomarkers in healthy controls (HC), TBI patients and patients with non-TBI traumatic injuries (NTT) at different timepoints. *, **, *** indicates a significance of $P<0.05$, $P<0.01$ and $P<0.001$ compared to controls. #, ##, ### indicates a significance of $P<0.05$, $P<0.01$ and $P<0.001$ compared to NTT at the same time point.

Fig. 4. Cerebral microdialysis evidence of axonal injury acutely following TBI. (A) CT images showing indicative placement of cerebral microdialysis catheter in deep white matter (B) Longitudinal trajectories NfL, TAU and UCH-L1 and GFAP in brain extracellular fluid

and plasma samples across all MD TBI patients over the first 10 days post-TBI. *** indicates a significance of $P < 0.001$ for concentrations in plasma compared to brain extracellular fluid.

Fig. 5. Axonal injury and neurodegeneration after moderate-severe TBI. (A) Fractional anisotropy (FA) in patients z-scored against local healthy controls in the whole white matter (WM, upper part) and the corpus callosum (CC, lower part), for subacute, 6 and 12 month visits in controls (blue) and patients (red). *** indicates a significance of $p < 0.001$ compared to controls. (B) Voxelwise comparison of zFA in patients vs controls with significant ($P < 0.05$) group differences in red, overlaid on the WM skeleton in green. Significant differences are seen at subacute (upper part), 6 month (middle) and 12 month (lower) visits. (C) Longitudinal change in FA measures between subacute and six month, and six to twelve month visits in WM and CC regions post-TBI. *** indicates a significance of $p < 0.001$ of zFA at timepoint 2 vs. timepoint 1. (D) Voxelwise map of regions with significant ($P < 0.05$) within-subject FA reductions between subacute and 6 months visits. (E) Jacobian determinant (JD) atrophy rates in grey matter (GM) and white matter (WM) in controls and patients. **, *** indicates a significance of $p < 0.01$, $p < 0.001$ compared to controls. (F) Voxelwise map of significant ($P < 0.05$) longitudinal reduction in WM volume (in blue) between 6-12 months post-injury.

Fig. 6. Relationship of TBI biomarkers brain atrophy rates and functional outcomes. Peak blood biomarker concentrations in the first six weeks post-TBI used for predictions, alongside the subacute z-scored FA in either corpus callosum (zFA CC) or white matter (zFA WM). (A) Peak biomarker concentrations and tissue-specific atrophy rates in grey matter from the subacute to six month visit (most strongly predicted by TAU), and in white matter from six to twelve months post injury (predicted by NfL). (B) Areas of significant correlation between subacute white matter FA and atrophy rates from 6-12 months post injury. White matter skeleton is shown in green. Significant (cluster corrected using TFCE with family-wise error rate of $P < 0.05$) areas of correlation voxelwise are shown in red colour. (C) Receiver operating characteristic curves predicting Glasgow Outcome Score Extended (GOSE) > 4 post-TBI at six (left) and twelve (right) months. For the six month

GOSE, significant ($P<0.05$) predictions are made using peak blood NfL, UCH-L1, GfAP, S100B and subacute FA in the corpus callosum (all $P<0.001$), subacute white matter FA ($P=0.001$) but not plasma TAU. For the twelve month GOSE, significant ($P<0.05$) predictors included peak blood NfL ($P<0.001$), UCH-L1 ($P=0.002$), GFAP ($P=0.001$), S100B, white matter and corpus callosum subacute FA (all $P<0.001$), but not plasma TAU.

Fig. 7. Correlations and factor structure of blood and neuroimaging biomarkers after TBI.

(A) Relationships of peak axonal injury plasma biomarker concentrations and subacute z-scored fractional anisotropy (zFA) in the corpus callosum (CC) showing significant correlations ($P<0.05$). (B) Factor analysis showing three factors explaining latent variable structure of biomarkers, with axonal markers NfL and corpus callosum zFA both loading strongly on Factor two.

Fig. 8. Axonal injury markers including NfL and DTI MRI in a controlled cortical impact model of TBI.

(A) NfL concentration in plasma at baseline, 3, 7, 10, 14 days post-injury in naïve, sham and after 1mm and 2mm CCI. *** indicates a significance of $P<0.001$ of 2mm CCI compared to controls. (B) CSF NfL 15 days post-injury (mean $\pm$ standard error of the mean). *** indicates a significance of $P<0.001$ compared to naïve/sham. (C) DTI-MRI images from naïve/sham animals, 1 and 2mm CCI (D) zFA after 1mm / 2mm CCI vs. controls. ** indicates a significance of $P<0.01$ of 2mm CCI compared to 1mm CCI. (E) Representative half brain photomicrographs of sections stained with Luxol Fast Blue from naïve/sham animals (left), 1mm (middle) and 2mm CCI (right). Magnifications (red rectangles) show ipsilateral corpus callosum (segments 2/3, dotted outline) of the respective groups. Black dotted scale bars correspond to 1 mm (half brain sections), black solid lines to 200 μ m (magnifications). (F) Quantification of Luxol Fast Blue staining showing thickness of segments of ipsilateral CC (% of the respective contralateral segment) measured every 500 μ m. *, *** indicates significance of $P<0.05$ and $P<0.001$ of 2mm CCI compared to naïve/sham. ## indicates significance of $P<0.01$ of 2mm CCI compared to 1mm CCI. (G) Neurofilament Alexa 568 immunofluorescence (grayscale) in naïve/sham (left), 1mm (middle) and 2mm CCI (right block). Magnifications (blue rectangles) show the ipsilateral corpus callosum (segments 2/3, dotted outline) with a HALO overlay

(threshold-based classification of intensity; red>orange>yellow). White dotted scale bars correspond to 1mm (half brain sections) and white solid lines to 100 μ m (inserts). **(H)** Quantification of neurofilament staining. Alexa 568 immunofluorescence intensity in the segments of the ipsilateral CC (% of the respective contralateral segment). *, *** indicates significance of $P<0.05$ and $P<0.001$ of 2mm CCI compared to naïve/sham. #, ### indicates significance of $P<0.05$, $P<0.001$ of 2mm CCI compared to 1mm CCI.

A TBI patient recruitment and study flow

B Patient assessment protocol

C Experimental injury protocol

A **B**

A Z-scored mean FA in patients vs controls **C** Z-scored mean FA change between scan visits **E** Comparison of JD in grey and white matter **B** Voxelwise comparison of z-scored mean FA in patients vs controls: Areas of reduced FA in patients **D** Areas of significant within subject reductions of z-scored FA between the acute and 6 month MRI scan **F** Areas of more negative JD within white matter of patients vs controls between 6 - 12 month scans

B Voxelwise correlation of subacute white matter FA and 6 to 12 month JD atrophy rate

A **B**

Supplementary Materials and Methods

Human blood biomarker assessment

Venous blood was sampled peripherally or by central access line if available, using ethylenediaminetetraacetic acid (EDTA) coated tubes and clot-activator for plasma and serum samples respectively. After 30 minutes at room temperature, samples were centrifuged at 2000-2500g at 4°C, transferred into 1.4ml aliquots and frozen at -80°C. NfL, total TAU, GFAP and UCH-L1 concentrations were assessed in plasma using a Neurology 4-plex B assay on the Simoa-HD1 platform, following the manufacturer's protocol (Quanterix Corp, Boston, MA). Serum S100B was measured using a Millipore ELISA kit (Merck Millipore, Darmstadt, Germany) with a lower limit of quantification of 2.7pg/ml. Samples were thawed on wet ice, centrifuged at $500 \times g$ for 5 minutes at 4°C. Calibrators (neat) and samples (1:4 dilution) were measured in duplicates. A four-parameter logistic curve fit data reduction method was used to generate a calibration curve. Two control samples of known concentrations of the protein of interest (high-control and low-control) were included as quality control. Samples were run in duplicate. The mean coefficient of variation across the study was 4.0% for NfL, 7.4% for total TAU, 16.5% for UCH-L1, 4.7% for GFAP, and 14% for S100B. The mean of both samples was used for further analyses. Any result falling below the lower limit of quantification (LLoQ) for a given biomarker was re-recorded as 50% of the LLoQ (GFAP: 9.38 pg/ml; NfL 0.5 pg/ml; S100B 2.7 pg/ml; TAU 0.125 pg/ml; UCH-L1 9.38 pg/ml). All results were included, with no removal of 'outliers' at lower or upper limits. Analyses were conducted at the biomarker lab at the UK Dementia Research Institute at UCL.

Experimental injury model blood biomarker assessment

Rats were inserted into a restrainer, the tail cleaned and approximately 200 μ L of whole blood withdrawn into EDTA-coated tubes. After centrifugation (10 min, 2000g), plasma was collected and immediately frozen. A single-plex Simoa NfL assay (Quanterix) was used to quantify rat plasma and CSF NfL and analysis conducted at the biomarker lab at the UK Dementia Research Institute at UCL using the same protocol as for human samples.

Human invasive cerebral microdialysis protocol and analysis

Microdialysis was initiated when clinically indicated and discontinued when no longer needed, typically five to seven days post-injury. 100kD cut-off MDialysis AB microdialysis catheters were surgically implanted through a triple lumen bolt cranial device into white matter regions that were not affected by gross focal pathologies such as contusions or haemorrhage on CT. This was typically in right, non-dominant frontal lobe. In patients

with frontal contusions, catheters were placed distant to intra-cerebral lesions. The placement was confirmed using a standard 2mm non-contrast CT head. Standardised commercially-available 3% Dextran 500 (M Dialysis, Stockholm, Sweden) solution was used to increase protein recovery, as indicated by the manufacturer (1). A standard flow rate of 0.3 mL/min was used in all patients, allowing hourly recovery of brain extracellular fluid (about 20 mL). Vials taken four hourly were immediately frozen and stored after collection and used exclusively for the measurement of biomarker concentrations. Samples were analysed using the Neurology 4-plex B assay on the the Quanterix Simoa-HD1 platform, as per blood biomarkers, but using a 20:1 dilution.

Human MRI acquisition

Participants in all sites had a high resolution structural T1 MPRAGE, T2 fluid-attenuated inversion recovery (FLAIR), susceptibility-weighted imaging (SWI) and diffusion tensor imaging (DTI). DTI sequences were aligned at each site and acquired in 64 directions with an isotropic voxel size of 2mm^3 , $b = 1000 \text{ s/mm}^2$, four images with $b = 0 \text{ s/mm}^2$ except in a single site (Santa Chiara Hospital – ‘Trento’) where only 60 directions were acquired and with a $1\text{mm} \times 2\text{mm} \times 2\text{mm}$ voxelwise. Two healthy volunteers were scanned at each site and oversaw harmonisation of scanning protocols. FA measures per white matter region of interest were all under 5% coefficient of variation (2). A physical N-tridecane phantom was scanned repeatedly at each site to ensure acquisition quality throughout the study. Further details for scanners and acquisition parameters can be found in Table S16 and the study protocol (3). Patients and controls were scanned using the same protocols within each centre and this protocol was kept consistent for the duration of the study. Structural MRI scans from all study sites were reviewed by a team of neuroradiologists in London and reported using electronic case report forms adapted from the National Institute of Neurological Disorders and Stroke (NINDS) TBI Imaging CDEs, providing a gold-standard readout of imaging abnormalities over time across the group.

Human MRI analysis - lesions

Focal lesions identified were manually segmented in native T1 space and scan/session-specific lesion masks were registered to MNI152 standard space and DTITK standard space using affine linear registration in FSL (fig. S5). An individual’s lesions were incorporated into the voxelwise analysis by including the lesioned voxels as voxelwise regressors in the general linear model.

Human MRI analysis – diffusion MRI including z-score normalisation across centres

Diffusion-weighted images were visually inspected before processing. Scans with uncorrectable artefacts or errors within acquisition were removed at this stage ($n = 6$). Scans were then processed following the standard tract based spatial statistics (TBSS) pipeline in the FMRIB Software Library (version 5.0.8) (4), including correcting susceptibility and eddy current induced distortions and diffusion tensor fitting. Brain extraction was

performed using HD-BET (Heidelberg University Hospital & Division of Medical Image Computing, German Cancer Research Center) (5). Tensor-based registration was performed using DTI-TK (6) involving the creation of a group template using affine and non-linear diffeomorphic registrations followed by registration of participant diffusion imaging to the template. Images were warped to 1mm isotropic space and the mean FA map produced was thresholded at 0.2 to produce a white matter skeleton. Subject FA data were projected onto the mean FA skeleton, with corpus callosum region of interest data generated using the Johns Hopkins University (JHU) white matter atlas. Images were repeatedly inspected at the brain extraction, eddy current correction and tensor registration stage. The mean Pearson correlation coefficient of the neighbouring diffusion-weighted images and slice-wise signal dropout for each slice was calculated for quality control checks. All data passed the criteria of baseline and follow-up having no difference in mean Pearson correlation coefficient greater than 0.1 while two scans from the same participant were removed due to signal dropout being higher than 0.1% of the total slice number (7). In total twelve diffusion scans were removed due to quality control issues.

Images were normalised across different centres to account for the inter-site scanner differences and enable groupwise comparisons in FA, with a z-scoring approach. In each site, raw FA maps in patients were normalised in a voxelwise manner to the local site-specific healthy control FA maps, who were age/sex matched to patients with TBI across the group. The following equation was used at each voxel:

$$zFA_{\text{patient}} = (\text{raw } FA_{\text{patient}} - \text{mean } FA_{\text{control group}}) / \text{standard deviation } FA_{\text{control group}}$$

This generates an individualised, normalised z-scored FA map for each patient which can then be compared between different sites. Mean z-scored FA was calculated for the whole white matter skeleton, and to account for any potential issues with scan quality diminishing at the boundaries, we also sampled zFA within the corpus callosum, a large, robust, central tract, while excluding sampling in areas with lesions. Voxelwise analysis of zFA was performed using threshold-free cluster enhancement (TFCE) multiple comparisons correction and a reported at a threshold significance level of $P < 0.05$. Longitudinal and cross sectional analysis of diffusion metrics was conducted using the general linear model with non-parametric permutation testing (10,000) in FSL Randomise (60), with age and gender included as a nuisance covariates in cross sectional analyses, and individualised lesion masking. In patients with microdialysis, the catheter tip was identified on CT imaging. CT scans were registered to DTITK space and a 10mm radius spherical region of interest was generated. The spherical region of interest was overlaid on the white matter skeleton and FA sampled within the overlapping voxels.

Human MRI analysis – volumetric MRI

Volumetric data were analysed using standard approaches (8). SPM 12 (UCL) was used to segment volumetric T1-weighted MRI images into grey matter, white matter and CSF to calculate volumes. A study-specific

template was generated using SPM DARTEL non-linear registration (9) to facilitate registration of segmented images to MNI152 space, with normalisation of volume and smoothing using an 8mm kernel. SPM12 longitudinal registration was used to assess within subject volume change over time. Briefly, the tool produces a temporal-average image by iteratively registering baseline and follow-up scans. A Jacobian determinant (JD) rate map is produced, encoding the change in volume from baseline to follow-up at each voxel, divided by the time separating the two scans (10). The JD rate maps are then registered to normal space, including smoothing using an 8mm kernel, where statistical analyses were performed. Region of interest atrophy rates were calculated by sampling mean grey matter JD values within left and right thalamus, and left and right hippocampus regions, defined in the Harvard-Oxford subcortical atlas (FSL). Focally lesioned voxels were not included in this estimation.

Experimental injury model overview

Experiments were conducted under a Home Office license and were in keeping with the Animal [Scientific Procedures] Act 1986 and EU legislation. Thirty-four Male Sprague Dawley rats (~8-9 weeks old) were housed in standard conditions (15). MRI was performed in 18 animals prior to group assignment. Animals were then randomised to sham-operation (n=4), mild/1 mm CCI (n=15) and moderate/2 mm CCI (n=15). Standard guidelines were followed to calibrate the CCI model, with 1mm and 2mm injuries previously established as corresponding with mild and moderate injuries in humans (61). Surgery and CCI were performed as previously (24). Anaesthesia was induced at least 30 min before the first incision and body temperature was adjusted and maintained at 37°C. Surgery and Controlled Cortical Impact were performed as previously published (11). A ~6 mm unilateral rectangular craniotomy, -0.5mm to -6.5 mm posterior and + 3.5 mm lateral to Bregma, was performed with a high-speed microdrill. The bone flap was stored in sterile saline. Immediately thereafter, CCI was induced using the 5 mm flat electromagnetically driven steel impactor (23° angle) performed at ~4 m/s for 100 milliseconds with an impact depth of either 1 or 2 mm. The dura was then inspected for signs of rupture, which occurred in one animal, in which the craniotomy was covered with absorbable gelatine sponges. The bone flap was re-implanted and the craniotomy was sealed with a nontoxic light-curing resin. After thorough irrigation and cleaning, the subcutaneous tissue and scalp were sutured and anaesthesia stopped. After surgery, all animals were single-housed and checked for signs of pain and discomfort every 30 minutes for the following 4 hours. For the next three days, all animals were weighed and inspected twice daily and afterward every morning. All animals were then housed with the same cage-mate as before. Further details can be found in a previous report (12). Sham-operated animals were subjected to all drugs and surgical procedures except craniotomy and impact. Blood was taken at baseline (two to four days prior to surgery), 3, 7, 10 and 14 days post-injury (fig. S6). Fourteen days post-impact, animals underwent a second MRI. The next day, the rats underwent terminal anaesthesia followed by transcardial perfusion and tissue

harvesting. Ten additional naïve animals served as histology controls and were only subjected to anaesthesia prior to transcardial perfusion.

Experimental injury model MRI acquisition

MRI scanning was performed in a 9.4T Bruker BioSpec scanner, equipped with a 4-channel phase array receiver coil and Paravision 6.1 software (Supplementary Figure 5A). Briefly, after localiser scans, high-resolution structural imaging was acquired with the following parameters: 3D T1 (TE=5 ms, TR=60 ms, $0.2 \times 0.2 \times 0.2 \text{ mm}^3$ resolution, 12 min acquisition) and 2D T2 (TE=33 ms, TR=5.5 s, RARE factor=8, $0.2 \times 0.2 \times 0.5 \text{ mm}^3$ resolution, 20 slices, acquisition time approximately 3 min). For diffusion, the multi-shell protocol included one shell with 40 gradient directions and $b = 1000 \text{ s/mm}^2$ and another with 40 directions and $b = 2000 \text{ s/mm}^2$. The protocol also contained four images without diffusion weighting ($b = 0 \text{ s/mm}^2$) and a single reversed-phase encoding image without diffusion weighting. The EPI readout (TE = 21 ms, TR = 4 s) had a resolution of $0.25 \times 0.25 \times 0.40 \text{ mm}^3$. A total of 34 contiguous slices were acquired for whole-brain coverage.

Experimental injury model MRI analysis – diffusion MRI

Standard post-processing steps were undertaken using FSL and FA extracted from the corrected multi-shell diffusion data. The corpus callosum was selected as a region of interest and segmented into 5 equal sections using MATLAB. Segments closest to the midline were labelled as segment 1. CC region of interest z-FA scores were generated at 14 days post injury using the mean and standard deviation of baseline scans of all animals as a ‘healthy control’ reference, to facilitate comparisons with the human neuroimaging data.

Experimental injury model histology quantification

Histology quantification was performed using methods previously described (12). Sections were stained with Luxol Fast Blue (LFB) and neurofilaments were stained using immunofluorescence. LFB-stained slides were imaged using on a slide scanner while immunofluorescence was imaged on a motorized fluorescent microscope. Corpus callosum (CC) thickness analysis was performed in FIJI and HALO®. Thickness of the CC was measured every 500 μm , beginning from the midline. Measurements per segment were averaged and values were normalized to the contralateral segment of the same section. Data is % change of contralateral. Neurofilament fluorescence intensity was quantified with the HALO® area quantification for fluorescence. Segments were calculated by taking the distance from the midline to the extent of the CC at the auditory cortex and dividing by five, and the average fluorescence intensity measured in each segment. Intensity was determined for each pixel and then normalized for the number of pixels. All measures were normalised to the contralateral segment of the same section and the final value expressed as % change of contralateral.

Sensitivity Analysis

A sensitivity analysis was performed to exclude significant influences of age on biomarker differences between patients with TBI and healthy controls. A group of 96 healthy individuals was matched to the TBI group with a median age of 47 ($P=0.346$). This had no effect on the significance of the results for all biomarkers (Supplementary Table 5).

Supplementary Tables

Supplementary Table 1. Participant demographics. *TBI: traumatic brain injury; NTT: non-TBI trauma patients; HC: healthy controls; ICP: intracranial pressure. Of the eighteen participants in the microdialysis arm, five are also in the main group ('all'). There was no significant difference ($P>0.05$) in demographics between the TBI ($n=197$), imaging ($n=146$) and microdialysis groups ($n=18$), or in clinical characteristics between TBI and imaging groups. Neurosurgical intervention and refractory raised intracranial pressures were significantly more common the microdialysis group (all P values <0.001 ; for neurosurgery and raised ICP respectively: TBI vs microdialysis $\chi^2=24.7$, 21.1 ; imaging vs microdialysis $\chi^2=25.1$, 20.2)*

	TBI			NTT	Healthy controls	
	All	Imaging	Microdialysis	All	Cross sectional	Longitudinal
Total, N	197	146	18	25	128	30
Age, median (IQR)	47 (30)	44 (29.5)	34 (27.5)	41 (29)	39 (27)	39.9 (26.9)
Male, N (%)	152 (77.2)	116 (79.5)	14 (77.8)	23 (92.0)	78 (61)	21 (70)
Centre, N						
Florence	13	11	0	0	13	0
Lausanne	22	19	14	0	15	0
Ljubljana	32	19	4	0	15	0
London	54	39	0	25	39	30
Niguarda	40	25	0	0	15	0
Policlinico	14	13	0	0	16	0
Trento	22	20	0	0	15	0
GCS (admission), median (IQR)	9 (8)	9 (7)	8 (8)	-	-	-
CT evidence of DAI, N (%)	35 (17.8)	29 (19.7)	7 (38.9)	-	-	-
CT midline shift, N (%)	40 (20.3)	28 (19.2)	7 (38.9)	-	-	-
Neurosurgery, N (%)	70 (35.5)	58 (39.5)	18 (100)	-	-	-
Refractory raised ICP, N (%)	27 (13.7)	26 (17.7)	11 (61.1)	-	-	-
Died in first month post-TBI, N (%)	10 (5.1)	0 (0.0)	5 (27.8)	-	-	-

Supplementary Table 2. Clinical characteristics of patients with TBI. TBI: traumatic brain injury; GOSE: Glasgow Outcome Scale Extended; SDH: subdural haematoma; SAH: subarachnoid haemorrhage; ICH: intracerebral haemorrhage; DAI: diffuse axonal injury; Evac: evacuation; All values are N (%) unless otherwise stated.

Demographics		Injury details	
Total (N)	197	Biophysics	110
Age, median (IQR)	47 (30)	High energy TBI	(55.8)
Male	152 (77.2)	Low energy TBI	68 (34.5)
Imaging subgroup (N)	146	Unknown	19 (9.6)
Age, median (IQR)	44 (29.5)	Injury cause	180
Male	116 (79.5)	Accidental	(91.4)
Microdialysis group (N)	18	Violence/assault	13 (6.6)
Age, median (IQR)	34 (27.5)	Self-harm	1 (0.5)
Male	14 (77.8)	Other/unknown, N (%)	3 (1.5)
Premorbid employment		Occupational injury	22 (11%)
Unfit for work	16 (8.1)	Road traffic accident	93 (47)
Home-maker	2 (1.0)	Bicycle	22
Retired	2 (1.0)	Car/Van	23
Student	11 (5.6)	Motorbike	23
Jobseeker	31 (15.7)	Other	2
Unknown	36 (18.3)	Pedestrian	23
Worker	99 (50.3)	Blunt weapon	4 (2.0)
Ethnicity		Gunshot	1 (0.5)
Asian	12 (6.0)	Crush	2 (1.0)
Black/African/Caribbean	5 (2.5)		
Other	1 (0.5)		
Unknown	15 (7.6)		
White	164 (83.2)		
Education			
University	36 (18.3)		
Secondary education	30 (15.2)		
Primary education	65 (33.0)		
Unknown	65 (33.0)		
No formal education	1 (0.5)		
Clinical Characteristics		Retrograde amnesia	
Initial GCS		definite	57 (28.9)
3 to 8	89 (45.2)	suspected	63 (32.0)
9 to 13	55 (27.9)	absent	19 (9.6)
14 to 15	51 (25.9)	unknown	58 (29.4)
unknown	2 (1.0)	Post traumatic amnesia	136
Lowset GCS first 24h, median (IQR)	7.5 (8)	definite/suspected	(69.0)
Pupils in ED	138	duration 1-30 mins	3 (2.2)
bilaterally reactive	(70.1)	duration 30-60 mins	4 (2.9)
unilaterally dilated/non-reactive	25 (12.7)	duration 1-24 hours	10 (7.4)
bilaterally dilated/nonreactive	1 (0.5)	duration 1-7 days	20 (14.7)
unknown	34 (17.3)	duration > 7 days	23 (16.9)
Loss of consciousness	121	duration unknown	76 (55.9)
definite	(61.4)	absent	14 (7.1)
suspected	26 (13.2)	unknown	61 (31.0)
absent	27 (13.7)		
unknown	23 (11.7)		
Neuroimaging reports		MRI performed (10 days - 6 weeks), N	
CT (acute), N	197	Subdural - acute	88 (62.9)

DAI	35 (17.8)	Subdural - subacute	7 (5.0)
Extradural	44 (22.3)	Contusion	119 (85.0)
SDH	113 (57.4)	Intracerebral haemorrhage	8 (5.7)
SAH	118 (59.9)	Intraventricular haemorrhage	39 (27.9)
Contusion/ICH	130 (66.0)	SAH	30 (21.4)
Midline shift	41 (20.8)	Extradural	6 (4.3)
CT Marshall classification (acute)		Microhaemorrhages	97 (69.3)
I (DI, no vis. Pathology)	23 (12.0)	Oedema	38 (27.1)
II (DI)	85 (44.5)	Midline shift	30 (21.4)
III (DI, oedema)	12 (6.3)	Cisternal compression	2 (1.4)
IV (DI, shift > 5mm)	6 (3.1)	Venous sinus injury	2 (1.4)
V (EML)	51 (26.7)	Brain atrophy	3 (2.1)
VI (NEML)	14 (7.3)	Ischaemia	1 (0.7)
Management and clinical course		Neurosurgery	
	135		
Intubated & ventilated	(68.5)	evac of EDH	18 (33.5)
Intracranial hypertension	58 (29.4)	evac of SDH	31 (15.7)
persistent	31 (15.7)	evac of parenchymal haematoma	8 (4.1)
		secondary decompression	8 (4.1)
Outcomes		GOSE, 12 months	
GOSE, 6 months			
dead (1)	13 (6.6)	dead (1)	13 (6.6)
vegetative state (2)	6 (3.0)	vegetative state (2)	6 (3.0)
severe disability (3,4)	48 (24.4)	severe disability (3,4)	28 (14.2)
moderate disability (5,6)	60 (30.5)	moderate disability (5,6)	52 (26.4)
good recovery (7,8)	58 (29.4)	good recovery (7,8)	62 (31.5)
unknown	12 (6.1)	unknown	36 (18.3)

Supplementary Table. 3. Characteristics of non-TBI trauma patients. Injury description for extracranial injury trauma patients. RTA: road traffic accident; low energy defined as <30kmph or fall from less than 3m.

Age	Sex	Mechanism	Energy	Description	Admission duration (days)	Management
30	Male	RTA	Low	Tibia & fibula fracture	14	Operative
52	Male	RTA	Low	Multiple rib fractures and clavicular fracture	2	Conservative
37	Female	Fall	High	Lumbar vertebral and calcaneal fractures	11	Operative
25	Male	Fall	Low	Tibia & fibula fracture	1	Operative
41	Male	RTA	High	Multiple rib fractures with haemopneumothorax and splenic laceration	4	Operative
53	Male	RTA	Low	Multiple rib fractures with haemopneumothorax; pulmonary contusion; scapular fracture;	7	Conservative
20	Male	Penetrating	-	Haemopneumothorax; pulmonary laceration; splenic laceration; renal injury; multiple rib fractures	7	Operative
69	Male	Fall	Low	Multiple rib fractures with haemopneumothorax; scapular fracture, pelvic fractures; thoracic spinal process fracture; sacral canal fracture	12	Operative
69	Male	RTA	Low	Hip and metatarsal fractures	9	Operative
46	Male	Fall	Low	Hepatic haematoma, rib fracture	2	Conservative
26	Male	Penetrating	-	Penetrating knee wound with vascular injury	5	Operative
23	Male	Fall	Low	Liver laceration; multiple rib fractures; splenic laceration; lumbar spinal transverse process fracture; pulmonary contusions	2	Conservative
53	Male	Fall	Low	Multiple rib fractures with haemopneumothorax; scapular fracture, pelvic fractures; thoracic spinal process fracture; sacral canal fracture	7	Conservative
55	Male	RTA	Low	Tibia & fibula fracture	8	Operative
36	Male	RTA	Low	Tibia & fibula fracture; soft tissue injury	16	Operative
33	Male	Fall	High	Multiple rib fractures with pneumothorax	3	Conservative
54	Male	Penetrating	-	Bilateral deep ankle wounds	6	Operative
68	Female	Fall	Low	Tibia & fibula fracture	5	Operative
69	Male	Fall	Low	Hip fracture	26	Operative
23	Male	Crush	-	Femoral fracture; soft tissue injury	2	Operative
68	Male	Fall	Low	Multiple rib fractures, humeral fracture	16	Conservative
37	Male	RTA	Low	Rib fractures with pneumothorax, humeral fracture, spinal endplate fractures	10	Operative
22	Male	RTA	Low	Tibia & fibula fracture	11	Operative
23	Male	Penetrating	-	Neck wound	5	Operative
60	Male	Fall	Low	Tibia & fibula fracture	11	Operative

Supplementary Table. 4. Blood biomarkers after TBI, extracranial injuries and in healthy controls.
Acute 1: first testing timepoint in days 0-10 post-trauma; Acute 2: second testing timepoint in days 0-10 post-trauma; subacute: ten days to six weeks post-injury. NTT: non-TBI trauma. HC: healthy control.

Biomarker		TBI					NTT		HC
		Acute 1	Acute 2	Subacute	Six month	One year	Acute 1	Acute 2	Visit 1
NfL plasma	Days since injury, median (IQR)	1 (1)	5 (2)	20 (14)	188 (24)	376.5 (30.25)	2 (1.25)	4 (2)	-
	n	167	149	141	53	25	24	19	96
	conc, median (IQR)	60.3 (103.6)	152.8 (230.7)	507.3 (728.2)	29.7 (30.5)	11.6 (6.5)	12.2 (11.3)	15.9 (10.7)	5.9 (4.6)
	P vs HC	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	-
	W	352.00	42.00	6.00	387.00	431.00	459.00	209.00	
	P vs NTT	<0.001	<0.001	-	-	-	-	-	-
TAU plasma	W	539	224	-	-	-	-	-	-
	n	168	150	141	53	25	24	19	96
	conc, median (IQR)	7.0 (18.8)	3.0 (4.1)	3.0 (3.5)	1.4 (1.1)	1.2 (1.1)	1.3 (1.0)	1.3 (0.7)	1.2 (1.0)
	P vs HC	<0.001	<0.001	<0.001	1.00	1.00	1.00	1.00	-
	W	1630.00	2708.00	2451.00	1777.00	1210.00	1039.00	944.00	
	P vs NTT	<0.001	<0.001	-	-	-	-	-	-
UCH-L1 plasma	W	466.00	469.00	-	-	-	-	-	-
	n	168	150	141	47	22	23	17	91
	conc, median (IQR)	97.6 (155.9)	30.6 (35.4)	27.6 (31.4)	4.7 (9.0)	7.3 (9.7)	13.1 (17.0)	10.8 (8.8)	4.7 (6.2)
	P vs HC	<0.001	<0.001	<0.001	1.00	1.00	0.002	1.00	-
	W	1098.00	1341	1668	1986	834	534	577	
	P vs NTT	<0.001	<0.001	-	-	-	-	-	-
GFAP plasma	W	595.00	367.00	-	-	-	-	-	-
	n	167	148	141	53	25	24	19	96
	conc, median (IQR)	12590.5 (31756.0)	2052.9 (5768.9)	330.2 (434.5)	76.7 (74.5)	77.8 (65.6)	72.7 (47.0)	77.0 (77.9)	57.0 (37.4)
	P vs HC	<0.001	<0.001	<0.001	0.081	0.041	0.66	1.00	-
	W	162.00	163.00	469.00	1773.00	691.00	792.00	741.00	
	P vs NTT	<0.001	<0.001	-	-	-	-	-	-
S100B serum	W	69.00	67.00						
	n	181	150	138	51	25	24	19	96
	conc, median (IQR)	215.1 (338.3)	90.2 (92.9)	50.2 (48.9)	45.8 (18.9)	46.8 (24.3)	69.7 (42.5)	53.9 (41.9)	44.5 (23.5)
	P vs HC	<0.001	<0.001	0.67	1.00	1.00	0.35	0.29	-
	W	1783.00	3166.00	5951.00	2456.00	1298.00	757.00	560.00	
	P vs NTT	<0.001	0.56	-	-	-	-	-	-
W		862.00	938.00						

Supplementary Table 5. Blood biomarker concentrations: lesion related analysis. Analysis of TBI patients either in Marshall Grade I compared to Marshall Grade II-VI on CT, or presence of lesions on MRI to no presence of lesion on MRI. Conc. = concentration.

	Blood biomarker (peak within 6 weeks of injury)				
	NfL	TAU	UCH-L1	GFAP	S100B
Marshall Grade I					
N	19	19	19	19	19
conc, median (IQR)	53.2 (134)	3.57 (7.99)	111 (145)	1980 (10600)	79.6 (98.3)
Marshall Grade II-VI					
N	154	155	155	153	159
conc, median (IQR)	147 (243)	8.42 (22.0)	35.8 (73.4)	15200 (31800)	257 (357)
W	1321	976	910	779	776
P	0.084	0.083	0.033	0.005	0.0027
Lesions present on MRI					
N	112	113	113	112	115
conc, median (IQR)	153 (257)	7.52 (19.9)	108 (150)	15800 (29300)	252 (343)
No lesions present					
N	16	16	16	16	16
conc, median (IQR)	46.2 (147)	1.19 (2.89)	22.9 (37.1)	1710 (2200)	65.3 (112)
W	1547	1518	1427.5	1484	1469
P	0.052	<0.001	<0.001	<0.001	<0.001
Correlation with lesion volume on MRI					
ρ	0.5	0.52	0.52	0.53	0.45
P	<0.001	<0.001	<0.001	<0.001	<0.001

Supplementary Table 6. Blood biomarker concentrations: neurosurgery analysis. Analysis of TBI patients with and without neurosurgery, comparing peak biomarker concentrations in the first six weeks post-injury.

	Intervention		Statistical comparison	
	No surgery	Surgery	W	P
NfL				
<i>N</i>	122	73		
median	290.2	574.9	2920	<0.001
TAU				
<i>N</i>	122	73		
median	5	13.5	2960	<0.001
UCH-L1				
<i>N</i>	122	73		
median	58.4	123.2	3140.5	0.0023
GFAP				
<i>N</i>	122	72		
median	6411.8	18402.5	3125	0.0032
S100B				
<i>N</i>	121	75		
median	144.2	281.4	3110.5	<0.001

Supplementary Table 7. Blood biomarker concentrations: neurosurgery analysis accounting for white matter FA. Analysis of TBI patients with and without neurosurgery, comparing peak biomarker concentrations in the first six weeks post-injury while accounting for injury severity as measured by white matter FA.

	Biomarker concentration		Statistical comparison	
	No surgery (n=82)	Surgery (n=49)	W	P
NfL				
peak in 6 weeks (pg/ml)	372.6	759.8	1168	<0.001
residuals	-196.6	22.9	1168	<0.001
TAU				
peak in 6 weeks (pg/ml)	3.8	9.8	1192	<0.001
residuals	-14.3	-12.7	1738	0.792
UCH-L1				
peak in 6 weeks (pg/ml)	45.9	114	1137	0.011
residuals	-58.2	-10.9	1627	0.28
GFAP				
peak in 6 weeks (pg/ml)	6016.4	17435.1	1361	0.0083
residuals	-10068.4	-1994.4	1705	0.6
S100B				
peak in 6 weeks (pg/ml)	131.9	235.4	1416	0.019
residuals	-106.7	-43.1	1812	1

Supplementary Table 8. Blood biomarker concentrations: age-related sensitivity analysis. Analysis with sub-group of healthy controls. Acute 1: first testing timepoint in days 0-10 post-trauma; Acute 2: second testing timepoint in days 0-10 post-trauma; subacute: ten days to six weeks post-injury. NTT: non-TBI trauma. HC: healthy control.

	NfL (plasma)		TAU (plasma)		UCH-L1 (plasma)		GFAP (plasma)		S100B (serum)	
	P vs HC	W	P vs HC	W	P vs HC	W	P vs HC	W	P vs HC	W
TBI										
Acute 1	<0.001	359	<0.001	197	<0.001	724	<0.001	124	<0.001	1453
Acute 2	<0.001	38	<0.001	1898	<0.001	830	<0.001	128	<0.001	2525
Subacute	<0.001	5	<0.001	1705	<0.001	1098	<0.001	370	1.00	4573
Six month	<0.001	348	1.00	1625	1.00	1474	1.00	1460	1.00	1890
One year	<0.001	395	1.00	887	1.00	618	0.60	591	1.00	985
NTT										
Acute 1	0.003	424	1.00	24	0.001	377	1.00	663	1.00	599
Acute 2	<0.001	193	1.00	678	1.00	442	1.00	589	1.00	448

Supplementary Table 9. Cerebral microdialysis: patient characteristics and biomarker trends. GCS: Glasgow Coma Scale; MD: microdialysate; F.R.: front right; F.L.: front left. Hosp.: hospital.

Age	Sex	GCS (pre- hosp)	ICU duration (days)	CT Marshall	Outcome (1 month)	Catheter location	GFAP (MD conc., pg/ml)		NfL (MD conc., pg/ml)		TAU (MD conc., pg/ml)		UCH-L1 (MD conc., pg/ml)		MRI days since injury	zFA (mean)		
							Mean	SD	Mean	SD	Mean	SD	Mean	SD		WM	CC	Catheter
54	Male	3 – 8	5	III	Survived	F. R.	3972	6021	2981	2321	119	106	2783	1037	43	-0.3	0.5	-0.4
31	Male	14 – 15	17	III	Survived	F. L.	126746	242115	18990	27717	2878	6656	9371	18705	33	-0.5	0.7	-0.7
25	Male	3 – 8	6	VI	Survived	F. L.	1351607	880197	75023	22366	14869	7212	15555	8323	-	-	-	-
20	Male	3 – 8	23	III	Survived	F. R.	175535	661049	15277	25097	3212	11485	5277	13099	16	-0.4	0.5	-0.8
31	Male	3 – 8	16	VI	Survived	F. R.	85331	97671	15187	6078	1272	1041	6938	7934	-	-	-	-
32	Male	3 – 8	21	II	Survived	F. R.	52624	57901	27320	14726	5577	7359	11627	8237	21	-0.5	1.4	-1.0
72	Male	3 – 8	13	VI	Died	F. R.	134079	167941	13384	10137	2136	3646	40917	35088	-	-	-	-
65	Male	3 – 8	16	VI	Died	F. R.	31839	26877	15097	5630	428	311	8643	10898	12	-1.3	1.8	-3.6
36	Female	3 – 8	9	VI	Survived	F. R.	64823	84861	17003	11861	4968	5691	31900	72024	21	-0.6	0.7	-0.4
19	Female	3 – 8	16	VI	Died	F. L.	5483	8526	2811	1823	49	24	1857	433	8	-1.0	2.1	-0.2
54	Male	3 – 8	23	II	Survived	F. R.	63711	125744	7773	10809	112	137	3437	2325	13	-1.6	3.0	-1.5
47	Male	3 – 8	24	V	Died	F. R.	1045822	668936	43747	40076	2541	1563	53287	33443	-	-	-	-
34	Male	3 – 8	18	VI	Survived	F. R.	13215	11387	15749	5461	707	266	3995	3248	-	-	-	-
22	Female	3 – 8	5	II	Survived	F. R.	7672	6281	4677	1513	484	73	4025	1480	20	-0.8	1.1	-0.6
24	Male	3 – 8	7	III	Survived	F. R.	24775	59407	1577	467	459	446	16654	23360	30	-1.8	3.4	-1.9
61	Male	3 – 8	28	III	Survived	F. R.	78632	154003	4964	5099	1469	6152	34475	100679	35	-2.6	5.2	-2.3
34	Male	9 – 13	42	IV	Survived	F. R.	9092	32427	2619	4301	588	1714	10559	27910	43	-2.6	6.2	-4.1
56	Female	14 – 15	23	VI	Died	F. R.	120219	99717	11839	4984	1291	2331	17027	27276	-	-	-	-

Supplementary Table 10. Biomarker concentrations in microdialysate and plasma. Relationships over ten days post injury in 18 patients undergoing invasive cerebral microdialysis. Ratio of microdialysate: plasma reflects median values. MD: microdialysate.

Biomarker	Location	Biomarker concentration (pg/ml)				Ratio MD : plasma (of median)	Statistics (MD vs plasma)	
		minimum	maximum	median	IQR		W	P
NfL	Microdialysate	86.7	163224.0	10059.0	16040.0	106.6	108017	< 0.001
	Plasma	17.8	910.0	94.4	167.0			
TAU	Microdialysate	4.5	61597.0	488.0	1730.0	105.6	107992	< 0.001
	Plasma	0.6	149.0	4.6	6.5			
UCH-L1	Microdialysate	1106.0	503373.0	3849.0	10412.0	105.5	108035	< 0.001
	Plasma	1.8	299.0	36.5	50.3			
GFAP	Microdialysate	178.0	3484315.0	19678.0	77501.0	4.3	77669	< 0.001
	Plasma	89.9	138100.0	4601.0	13416.0			

Supplementary Table. 11. MRI measures of white matter integrity and progressive atrophy. Cross-sectional assessments of fractional anisotropy (FA) and progressive brain atrophy (JD, Jacobian determinant atrophy rate) after TBI and in healthy controls.

	Traumatic Brain Injury			Healthy Control
<i>Cross sectional assessments</i>	<i>Subacute visit</i>	<i>6 month visit</i>	<i>12 month visit</i>	<i>Single visit</i>
Days since injury, median (IQR)	20 (14.5)	189 (23)	372 (20)	-
Diffusion tensor imaging, N	132	53	39	121
mean WM zFA, median (IQR)	-0.338 (0.798)	-0.329 (0.585)	-0.242 (0.649)	0.004 (0.360)
test statistic vs HC (W)	11720	4556	3112	-
significance (P)	<0.001	<0.001	<0.001	-
mean callosum zFA, median (IQR)	-0.999 (1.278)	-0.662 (0.810)	-0.599 (0.957)	0.002 (0.402)
test statistic vs HC (W)	12485	4720	3138	-
significance (P)	<0.001	<0.001	<0.001	-
<i>Longitudinal assessments</i>	<i>N/A</i>	<i>Subacute to 6 month</i>	<i>6 to 12 months</i>	
Serial volumetric imaging, N	-	58	38	30
GM JD, median (IQR)	-	-0.007 (0.015)	-0.001 (0.005)	0.000 (0.002)
test statistic vs HC (W)	-	469	504	-
significance (P)	-	<0.001	NS (0.421)	-
WM JD, median (IQR)	-	0.000 (0.021)	-0.007 (0.016)	-0.001 (0.005)
test statistic vs HC (W)	-	803	309	-
significance (P)	-	NS (0.558)	0.001	-
CSF JD, median (IQR)	-	0.016 (0.08)	0.012 (0.016)	0.004 (0.006)

test statistic vs HC (W)	-	1214	769	-	
significance (P)	-	0.002	0.005	-	
Hippocampus JD, median (IQR)	-			0.000 (0.004)	
test statistic vs HC (W)	-	-0.018 (0.038)	-0.002 (0.015)	-	
significance (P)	-	0	0.212	-	
Thalamus JD, median (IQR)	-	-0.006 (0.017)	-0.001 (0.007)	0.000 (0.002)	
test statistic vs HC (W)	-	1151	663	-	
significance (P)	-	0.014	0.253	-	
	Traumatic Brain Injury				Healthy Control
Cross sectional assessments	Subacute visit	6 month visit	12 month visit	Single visit	
Days since injury, median (IQR)	20 (14.5)	189 (23)	372 (20)	-	
Diffusion tensor imaging, N	132	53	39	121	
mean WM zFA, median (IQR)	-0.338 (0.798)	-0.329 (0.585)	-0.242 (0.649)	0.004 (0.360)	
test statistic vs HC (W)	11720	4556	3112	-	
significance (P)	<0.001	<0.001	<0.001	-	
mean callosum zFA, median (IQR)	-0.999 (1.278)	-0.662 (0.810)	-0.599 (0.957)	0.002 (0.402)	
test statistic vs HC (W)	12485	4720	3138	-	
significance (P)	<0.001	<0.001	<0.001	-	
Longitudinal assessments	N/A	Subacute to 6 month	6 to 12 months		
Serial volumetric imaging, N	-	58	38	30	
GM JD, median (IQR)	-	-0.007 (0.015)	-0.001 (0.005)	0.000 (0.002)	
test statistic vs HC (W)	-	469	504	-	
significance (P)	-	<0.001	NS (0.421)	-	
WM JD, median (IQR)	-	0.000 (0.021)	-0.007 (0.016)	-0.001 (0.005)	
test statistic vs HC (W)	-	803	309	-	
significance (P)	-	0.558	0.001	-	
CSF JD, median (IQR)	-	0.016 (0.08)	0.012 (0.016)	0.004 (0.006)	
test statistic vs HC (W)	-	1214	769	-	
significance (P)	-	0.002	0.005	-	

Hippocampus JD, median (IQR)	-	-0.018 (0.038)	-0.002 (0.015)	0.000 (0.004)
test statistic vs HC (W)	-	1287	672	-
significance (P)	-	<0.001	0.212	-
Thalamus JD, median (IQR)	-	-0.006 (0.017)	-0.001 (0.007)	0.000 (0.002)
test statistic vs HC (W)	-	1151	663	-
significance (P)	-	0.014	0.253	-

Supplementary Table 12. Relationship of blood biomarkers and long-term outcomes after TBI. Relationships of peak blood biomarkers sampled in first six weeks after injury and outcome measures. zFA: z-scored fractional anisotropy measure of white matter integrity. Correlations for atrophy rates are using the tissue-specific Jacobian determinant atrophy rates. AUC: area under receiver operating characteristic curve.

Outcome and Timepoint		Biomarker Predictors						
		Fluid markers					DTI MRI markers	
		NfL (peak)	TAU (peak)	UCH-L1 (peak)	GFAP (peak)	S100B (peak)	zFA WM all (subacute)	zFA CC (subacute)
Function: GOSE								
6m (n=126)	R ² *	0.24	0.04	0.12	0.12	0.16	0.11	0.18
	AUC	0.83	0.77	0.77	0.73	0.81	0.69	0.76
	95% CI	0.75-0.91	0.68-0.87	0.68-0.86	0.63-0.82	0.72-0.89	0.59-0.80	0.66-0.85
	P	<0.001	0.13	<0.001	<0.001	<0.001	0.001	<0.001
12m (n=108)	R ² *	0.28	0.05	0.12	0.13	0.16	0.22	0.25
	AUC	0.85	0.85	0.81	0.78	0.83	0.79	0.81
	95% CI	0.77-0.94	0.78-0.92	0.72-0.89	0.69-0.86	0.74-0.91	0.68-0.91	0.70-0.92
	P	<0.001	0.17	0.002	0.001	<0.001	<0.001	<0.001
Neurodegeneration: Atrophy Rate								
subacute to 6 months (n=53)								
Grey matter	Adj. R ²	0.20	0.34	0.25	0.27	0.21	0.00	0.00
	P	0.001	<0.001	0.001	<0.001	0.004	1.000	1.000
6 to 12 months (n=33)								
WM JD	Adj. R ²	0.14	-0.03	-0.02	0.02	0.06	0.08	0.06
	P	0.0374	1.0000	1.0000	1.0000	1.0000	0.667	1.0000

Supplementary Table 13. Cerebral Microdialysis: DTI imaging measures. Z-scored FA in the whole white matter skeleton, corpus callosum and within a region of interest placed around the catheter in microdialysis patients compared to healthy controls. Due to variable placement of the catheter per individual patient, a mean z-score of FA within the ROI's for patients is shown only.

	Microdialysis TBI patients	Healthy Control
Days since injury, median (IQR)	25 (12)	-
Diffusion tensor imaging, N	12	116
mean WM zFA, median (IQR)	-0.9(1.15)	0.004 (0.365)
test statistic vs HC (W)	1336	-
significance (P)	<0.001	-
mean callosum zFA, median (IQR)	-1.6 (2.4)	0.02 (0.5)
test statistic vs HC (W)	1378	-
significance (P)	<0.001	-
mean catheter ROI zFA, median (IQR)	-0.9 (1.45)	-
test statistic vs 0 (V)	0	-
significance (P)	0.003	-

Supplementary Table 14. Blood and imaging markers in an experimental injury model. CCI: controlled cortical injury.

Biomarker	Timepoint	Naïve		Sham		P vs naïve	1mm CCI			P vs naïve	P vs sham	2mm CCI			P vs naïve	P vs sham
		n	conc, median (IQR)	n	median (IQR)		n	median (IQR)	P vs naïve			n	median (IQR)	P vs naïve		
NfL (plasma, pg/ml)	Baseline	10	9.4 (5.5)	1	17.1 (0)	-	12	6.52 (4.8)	0.4	-		12	7.12 (4.1)	1.00	-	
	Day 3	-	-	4	23.0 (38.4)	-	14	140.2 (142.4)	-	0.24		13	469.8 (226.4)	-	<0.001	
	Day 7	-	-	4	14.7 (11.2)	-	14	44.5 (66.1)	-	0.024		13	137.0 (119.8)	-	<0.001	
	Day 10	-	-	4	16.1 (3.9)	-	10	50.15 (46.1)	-	0.15		10	84.2 (48.1)	-	<0.001	
	Day 14/15	-	-	4	12.9 (1.8)	-	9	22.08 (40.2)	-	0.033		13	54.9 (40.2)	-	<0.001	
NfL (CSF, pg/ml)	Day 15	9	791.4 (713.9)	3	1283.1 (1154.1)	1.00	13	2530.6 (3174.8)	0.52			14	10003.0 (6010.6)	<0.001	<0.001	
Corpus Callosum zFA	Baseline	3	0.44 (0.02)	-	-	-	8	0.45 (0.04)	0.26	-		7	0.43 (0.06)	0.74	-	
	Day 14	-	-	-	-	-	13	0.44 (0.1)	-	-		7	0.35 (0.06)	-	-	

Supplementary Table 15. Plasma biomarkers after experimental injury: mixed effects models results. LFB: Luxol fast blue white matter tissue quantification. CCI: controlled cortical impact. ns: not significant

Biomarker		1mm CCI		2mm CCI	
		q(N)	P	q(N)	P
Plasma NfL (pg/ml)	Timepoint				
	Day 3	q(10.00)=4.9	0.014	q(9.00)=6.769	0.002
	Day 7	q(10.00)=4.8	0.018	q(10.00)=8.095	<0.001
	Day 10	q(5.00)=3.7	ns		
	Day 14/15	q(11.00)=5.0	0.009	q(9.00)=5.970	0.005
LFB-positive tissue (% change vs contralateral)	Segment				
	S1	-	ns	-	ns
	S2	-	ns	q(80.00)=5.878	0.002
	S3	-	ns	q(80.00)=8.906	<0.001
	S4	-	ns	q(80.00)=3.926	ns
	S5	-	ns	-	ns
NF staining (% change vs contralateral)	Segment				
	S1	-	ns	-	ns
	S2	-	ns	q(80.00)=3,533	ns
	S3	-	ns	q(80.00)=5.760	0.003
	S4	-	ns	q(80.00)=7.223	<0.001
	S5	-	ns	q(80.00)=7.280	<0.001

Supplementary Table 16. Imaging acquisition parameters and scanner details. MPRAGE: Magnetization Prepared RApid Gradient Echo; DTI: diffusion tensor imaging; FLAIR: FLuid Attenuated Inversion Recovery; SWI: Susceptibility Weighted Imaging.

	Research Site						
	Florence	Lausanne	Ljubljana	London	Niguarda	Policlinico	Trento
Scanner							
manufacturer	Siemens	Siemens	Siemens	Siemens	Philips	Philips	GE
model	Aera	Skyra Fit	Trio Tim	Verio	Achieva	Achieva	Optima
field strength	1.5 Tesla	3 Tesla	3 Tesla	3 Tesla	1.5 Tesla	3 Tesla	1.5 Tesla
software	VE11a	MR B19	MR B19	MR B17	Rel. 5 Neuroradio	Rel. 5 Neuroradio	DV25.1_R03_1802.a
Voxelsize (mm)							
MPRAGE	1x1x1	1x1x1	1x1x1	1x1x1	1x1x1	1x1x1	0.5x0.5x1
DTI	2x2x2	2x2x2	2x2x2	2x2x2	2x2x2	2x2x2	1x2x2
FLAIR	0.5x0.5x1	0.5x0.5x1	1x1x1	1x1x1	1.2x1.2x0.7	1x1x1	1.2x0.5x0.5
SWI	0.8x0.8x0.8	0.3x0.3x1.6	0.6x0.5x1.2	0.6x0.5x1.2	1x1x1	0.5x0.5x1.2	0.5x0.5x1
DTI parameters							
receiver coil channels	20	64	32	32	8	32	8
directions	64	64	64	64	64	64	60
b-value	1000	1000	1000	1000	1000	1000	1000

Sup Fig. 1. Spatial pattern of MRI-defined focal lesions in patients TBI. Colourbar denotes number of patients. **(A)** Subacute patterns (ten days to six weeks post-injury) of focal lesions on MRI. **(B)** Longitudinal evolution of lesions (composite of all types) in 58 patients with serial structural MRI.

Sup. Fig 2. No significant relationship between age at injury and plasma NfL concentration at six and twelve months post-TBI. Plasma NfL at six months (left panel) and twelve months after TBI (right panel) related to age at injury. Linear regression shows non-significant relationship ($P>0.05$) in both cases. Adjusted R^2 's shown.

Sup. Fig 3. Individual fluid biomarker trends in cerebral microdialysate and blood. Biomarker concentration in cerebral microdialysate and plasma. Each row represents an individual patient.

Sup Fig 4. Experimental injury model fluid biomarker, neuroimaging and histology relationships.

Correlations between plasma NFL, CSF NFL, zFA from DTI and histopathological results. CC: corpus callosum. Asterisk denotes significant result at $P < 0.05$.

Sup. Fig. 5. Neuroimaging acquisition and analysis pipelines. MRI data including diffusion weighted imaging (DWI) and T1-volumetric structural images were processed using the following approaches. **(A)** DWI images are eddy-corrected, undergo brain extraction, FSL DTIFIT extraction of tensors and DTITK registration. **(B)** Volumetric images are segmented in SPM producing grey matter, white matter and CSF maps. Custom lesion masks are created per subject in native space. Scans are registered to a custom study template (non linear, via DARTEL in SPM). **(C)** Longitudinal structural images are assessed using the SPM longitudinal tool, whereby pairs of images are co-registered iteratively to produce an average space image and change rate map (the Jacobian determinant map). These are segmented and registered to MNI space. Statistical comparisons are undertaken using summary measures, region of interest approaches or voxelwise analyses.

Sup Fig 6. Controlled cortical injury methods. (A) Timeline of experimental model. (B) Timeline of histology examination in relation to injury and imaging. Skull and brain show the location of the craniotomy and injury. From T2 MRI, a 3D printed with brain matrix was created. Blocks of 2 mm thickness were cut, paraffinized and sectioned (7 μ m thickness), with block 4 (contusion core) being the main interest. In vivo DTI-MRI data was acquired from this block and sections were stained for white matter integrity (Luxol Fast Blue, LFB; Neurofilaments) and glia cells (microglia, astrocytes). Additional blocks 2, 3 and 5 were only investigated for white matter integrity. (C) Histology analysis: Stereotaxic coordinate for the 4 different blocks with regions of interest outlined in red. For assessment of white matter integrity, sections were stained with Luxol Fast Blue and the thickness measured in the individual segments using FIJI/HALO. For the quantification of Neurofilament fluorescence intensity and numbers of microglia and astrocytes, the CC segment was outlined and analysed in HALO.

Appendix:

Contributor list: The BIO-AX-TBI Group

In addition to those members of the consortium on the authorship list, we acknowledge the contribution of the following individuals.

ASST Grande Ospedale Metropolitano Niguarda

Francesco Curto
Giorgio Chevallard
Marco Sacchi
Anna Politi
Cristiana Cipolla
Beatrice Alberti
Francesca Turchetto
Luciana Gennari
Maurizio Sberna
Fortunata Imbesi
Alberto Torresin
Francesca Fossi

Azienda Ospedaliero-Universitaria Careggi

Luca Bucciardini
Vanni Orzalesi
Alessandra De Luca
Caterina Frosini
Silvia Marchiani
Fabio Picciafuochi
Stefano Chiti

Centre Hospitalier Universitaire Vaudois

Dr. Nawfel Ben-Hamouda
Dr. Vincent Dunet
Eva Favre

Fondazione IRCCS, Ca' Granda Ospedale Maggiore Policlinico

Prof. Nino Stocchetti
Julian Dallmeier
Marco Carbonara

Imperial College London & Imperial College Healthcare NHS Trust

Zayneb Al-Saadi
Michela Cicchetti
Andrea D'Mello
Irene Faiman

Sonia Fernandez-Lopez
Hoi Lun Helen Lai
Heather McLachlan
Sebastian Popescu
Alberto Rigoni
Margherita Rosnati
Rhys Thomas
Rachel Williams
Louise Young
Karolina Zimmerman

Istituto di Ricerche Farmacologiche Mario Negri IRCCS

Joanne Fleming

Ospedale S. Chiara di Trento, Azienda Provinciale per i Servizi Sanitari - APSS

Umberto Rozzanigo
Daniele Penzo
Elena Baruzzo
Costanza Papagno

University College London

Dr Martha Foiani
Dr Henrietta Wellington

University Medical Centre Ljubljana / Univerzitetni Klinični Center Ljubljana

Tomislav Mirkovic, MD, PhD
Vesna Novak Jankovic, MD, PhD
Suada Filekovic, MD, PhD
Katarina Surlan Popovic, MD, PhD
Matej Vrabec, MD, PhD
Andrej Sirnik
Sabina Salihagic
Sabina Kovacic
Pajo Bucic
Natalija Bruncic
Roman Bosnjak, MD, PhD
Tilen Žele, MD, PhD
Boštjan Matos, MD, Msc
Barbara Smrke, MD

Supplementary References

1. S. Giorgi-Coll, E. P. Thelin, C. Lindblad, T. Tajsic, K. L. H. Carpenter, P. J. A. Hutchinson, A. Helmy, Dextran 500 Improves Recovery of Inflammatory Markers: An In Vitro Microdialysis Study. *J Neurotrauma* **37**, 106-114 (2020).
2. E. M. Palacios, A. J. Martin, M. A. Boss, F. Ezekiel, Y. S. Chang, E. L. Yuh, M. J. Vassar, D. M. Schnyer, C. L. MacDonald, K. L. Crawford, A. Irimia, A. W. Toga, P. Mukherjee, Toward Precision and Reproducibility of Diffusion Tensor Imaging: A Multicenter Diffusion Phantom and Traveling Volunteer Study. *AJNR Am J Neuroradiol* **38**, 537-545 (2017).
3. N. S. Graham, K. A. Zimmerman, G. Bertolini, S. Magnoni, M. Oddo, H. Zetterberg, F. Moro, D. Novelli, A. Heslegrave, A. Chierigato, E. Fainardi, J. M. Fleming, E. Garbero, S. Abed-Maillard, P. Gradisek, A. Bernini, D. J. Sharp, Multicentre longitudinal study of fluid and neuroimaging BIOMarkers of AXonal injury after traumatic brain injury: the BIO-AX-TBI study protocol. *BMJ Open* **10**, e042093 (2020).
4. S. M. Smith, M. Jenkinson, H. Johansen-Berg, D. Rueckert, T. E. Nichols, C. E. Mackay, K. E. Watkins, O. Ciccarelli, M. Z. Cader, P. M. Matthews, T. E. Behrens, Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *Neuroimage* **31**, 1487-1505 (2006).
5. F. Isensee, M. Schell, I. Pflueger, G. Brugnara, D. Bonekamp, U. Neuberger, A. Wick, H.-P. Schlemmer, S. Heiland, W. Wick, M. Bendszus, K. H. Maier-Hein, P. Kickingeder, Automated brain extraction of multisequence MRI using artificial neural networks. *Human Brain Mapping* **40**, 4952-4964 (2019).
6. H. Zhang, B. B. Avants, P. A. Yushkevich, J. H. Woo, S. Wang, L. F. McCluskey, L. B. Elman, E. R. Melhem, J. C. Gee, High-dimensional spatial normalization of diffusion tensor images improves the detection of white matter differences: an example study using amyotrophic lateral sclerosis. *IEEE Trans Med Imaging* **26**, 1585-1597 (2007).
7. F.-C. Yeh, I. M. Zaydan, V. R. Suski, D. Lacomis, R. M. Richardson, J. C. Maroon, J. Barrios-Martinez, Differential tractography as a track-based biomarker for neuronal injury. *NeuroImage* **202**, 116131 (2019).
8. J. H. Cole, A. Jolly, S. de Simoni, N. Bourke, M. C. Patel, G. Scott, D. J. Sharp, Spatial patterns of progressive brain volume loss after moderate-severe traumatic brain injury. *Brain* **141**, 822-836 (2018).
9. J. Ashburner, A fast diffeomorphic image registration algorithm. *Neuroimage* **38**, 95-113 (2007).
10. J. Ashburner, G. R. Ridgway, Symmetric diffeomorphic modeling of longitudinal structural MRI. *Front Neurosci* **6**, 197 (2012).
11. C. K. Donat, G. Scott, S. M. Gentleman, M. Sastre, Microglial Activation in Traumatic Brain Injury. *Front Aging Neurosci* **9**, 208 (2017).
12. C. K. Donat, M. Yanez-Lopez, M. Sastre, N. Baxan, M. Goldfinger, R. Seeamber, F. Müller, P. Davies, P. Hellyer, P. Siegkas, S. Gentleman, D. J. Sharp, M. Ghajari, From biomechanics to pathology: predicting axonal injury from patterns of strain after traumatic brain injury. *Brain* **144**, 70-91 (2021).