

The In Situ Signature of Cyclotron Resonant Heating

Trevor A. Bowen,^{1,*} Benjamin D. G. Chandran,² Jonathan Squire,³ Stuart D. Bale,^{1,4} Die Duan,⁵ Kristopher G. Klein,⁶ Davin Larson,¹ Alfred Mallet,¹ Michael D. McManus,^{1,4} Romain Meyrand,³ Jaye L. Verniero,⁷ and Lloyd D. Woodham⁸

¹*Space Sciences Laboratory, University of California, Berkeley, CA 94720-7450, USA*

²*Department of Physics & Astronomy, University of New Hampshire, Durham, NH 03824, USA*

³*Department of Physics, University of Otago, 730 Cumberland St., Dunedin 9016, New Zealand*

⁴*Physics Department, University of California, Berkeley, CA 94720-7300, USA*

⁵*School of Earth and Space Sciences, Peking University, Beijing, 100871, China*

⁶*Department of Planetary Sciences & Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721*

⁷*NASA Goddard Space Flight Center*

⁸*Department of Physics, The Blackett Laboratory,*

Imperial College London, London, SW7 2AZ, UK

The dissipation of magnetized turbulence is an important paradigm for describing heating and energy transfer in astrophysical environments such as the solar corona and wind; however, the specific collisionless processes behind dissipation and heating remain relatively unconstrained by measurements. Remote sensing observations have suggested the presence of strong temperature anisotropy in the solar corona consistent with cyclotron resonant heating. In the solar wind, in situ magnetic field measurements reveal the presence of cyclotron waves, while measured ion velocity distribution functions have hinted at the active presence of cyclotron resonance. Here, we present Parker Solar Probe observations that connect the presence of ion-cyclotron waves directly to signatures of resonant damping in observed proton-velocity distributions using the framework of quasilinear theory. We show that the quasilinear evolution of the observed distribution functions should absorb the observed cyclotron wave population with a heating rate of 10^{-14} W/m³, indicating significant heating of the solar wind.

PACS numbers:

Introduction Observations of the solar corona reveal plasma that is millions of degrees hotter than the black-body temperature of the solar surface. While the energy required to heat the corona, and accelerate the solar wind originates from solar convection and the magnetic fields produced by the solar dynamo, the specific pathways to heating and particle acceleration remain elusive [1]. The dissipation of Alfvénic turbulence at kinetic scales has become a common paradigm in explaining the dynamics of coronal heating and solar wind acceleration [2–5]; possible dissipative mechanisms include Landau or cyclotron resonant damping [6–10], stochastic heating [11], or magnetic reconnection [12, 13]. Additionally, the portion of energy deposited by these processes at ion scales, versus that which is subject to a kinetic cascade and dissipated by electrons, remains an open question [10, 14–17].

It is well known that the observed ion temperature profiles in the solar wind require significant perpendicular heating [18, 19], which is likely initiated at ion kinetic scales where particles interact efficiently with electromagnetic waves [9, 16, 20–24]. Cyclotron resonant coupling of electromagnetic fluctuations with ion gyromotion [25], has received particular attention as a potential perpendicular heating mechanism [26–30]. Ultraviolet spectroscopic measurements of coronal ion temperature anisotropy suggest large T_{\perp}/T_{\parallel} , consistent with cyclotron resonant heating [28, 31–33]. The presence of ion-cyclotron waves has been well documented in in-situ

observations throughout the heliosphere both as solitary waves and as part of the background spectrum of fluctuations [34–40]. Observations of magnetic-helicity at ion scales have been interpreted as evidence for active cyclotron damping of quasi-parallel Alfvénic fluctuations, which contribute to turbulent heating [8, 9, 41, 42].

Theoretical signatures of resonant interactions in particle distribution functions are often studied in the framework of quasilinear (QL) diffusion [15, 43–45]; observations of the solar wind have suggested evidence for QL cyclotron resonant diffusion in signatures of the proton velocity distribution function $f_p(\mathbf{v})$ [46–48]. While the generation of cyclotron waves through instabilities has been widely discussed [37, 38, 40, 49, 50] and signatures of cyclotron resonant dissipation have been suggested [8, 9, 46, 48, 51–53], definitive cyclotron resonant heating sufficient to power the solar wind has not been observed.

In this Letter, we apply the QL theory of resonant cyclotron interactions [43, 44] to empirically measured cyclotron wave spectra and ion-distribution functions. Our results provide evidence of substantial heating at levels comparable with bulk solar wind heating rates, providing a compelling picture of ion-heating in the solar wind.

Data Parker Solar Probe (PSP, [54]) observations from the electromagnetic FIELDS [55] and Solar Wind Electron Alpha and Proton (SWEAP, [56]) instruments aim to constrain fundamental processes of around coronal heating and solar wind acceleration. PSP has re-




FIG. 1: (a) Magnetic field measurements from PSP/FIELDS. (b) Velocity measurements from PSP/SPANi. (c) Spectra of magnetic field data. Spectral indices at $-3/2$, -4 and -2.7 are shown. Wavelet coefficients for total power are shown as black \diamond . Wavelet coefficients filtered by left/right handed σ_b is shown in red/blue \diamond .

vealed prevalent ion-scale electromagnetic waves [39, 40], ion-distributions out of thermal equilibrium [57, 58], and evidence for resonant wave-particle interactions predicted by QL theory [59]. To constrain cyclotron resonant heating, we study a stream from PSP perihelion 4 from 2020-01-30/00:00-08:00 with resolved proton distributions. During the interval PSP was $\sim 30R_\odot$ from the solar surface. We use merged search coil and fluxgate magnetometer data from PSP/FIELDS [55, 60] enabling measurement of the inertial, transition, and kinetic scales of turbulence; the merged data set only has two axes available [61], thus we use vector-fluxgate magnetometer data to study wave-polarization. Fig 1(a) shows \mathbf{B} in Radial-Tangential-Normal (RTN) coordinates. Proton velocity distribution functions $f_p(\mathbf{v})$ are obtained from the PSP/SWEAP Solar Probe ANalyzer (SPANi). The proton population is often parameterized with a pair of drifting bi-Maxwellian fits to model $f_p(\mathbf{v})$ using separate thermal (core) and non-thermal (beam) populations [1]. Fits to a proton core and field-aligned beam provide estimates of bulk velocity \mathbf{u} , anisotropic temperatures perpendicular and parallel the background magnetic field $T_{\perp,\parallel}$, and the beam-to-core proton density ratio n_b/n_c [56, 58]. Fig 1(b) shows measurements of \mathbf{u} in RTN coordinates. The stream is relatively slow with an average speed of ~ 220 km/s, and moderately Alfvénic with a

cross helicity of ~ 0.85 .

The phase-space density of $f_p(\mathbf{v})$ is calibrated to quasi-thermal noise from FIELDS to recover the absolute density [55, 62]. The mean proton density is $n_p = 1100/\text{cm}^3$, SPANi gives an average beam to core density ratio of 0.48. The core/beam have T_\perp of 15 eV/22 eV and T_\parallel of 12 eV/30 eV. The average drift of the beam relative to the core is 83 km/s. The individual core and beam have $\beta_c = 0.65$ and $\beta_b = 1.1$. The mean magnetic field was directed sunward, with an Alfvén speed of 60 km/s. Fig 1(c) shows the magnetic field spectra of the interval with a steep transition range at ion-kinetic scales [63–65].

We apply a Morlet wavelet transform to the vector magnetic field data rotated into field aligned coordinates [66]. Signatures of circular polarization are found using

$$\sigma_B(f, t) = -2\text{Im}(B_{\perp 1}B_{\perp 2}^*)/(B_{\perp 1}^2 + B_{\perp 2}^2), \quad (1)$$

with left/right handed waves corresponding to positive/negative σ_B [35, 36, 67, 68]. Circular polarization is measured in the spacecraft frame, such that the measured sign may not correspond to the innate plasma frame polarization [67]. A sign change in σ_B occurs if the wave is Doppler shifted to negative frequencies in the spacecraft frame. However, it has been demonstrated that the majority of waves propagate outward, and thus, that Doppler shift does not change their handedness when observed in the spacecraft frame [69].

Previous work has shown that circularly polarized ion-scale waves are parallel propagating and evident when $\theta_{vB} \sim 0$ [40]. However, observations of parallel propagating, circular polarized, waves are strongly inhibited when the angle between the solar wind and the mean magnetic field is oblique. This effect occurs because a) the wave polarization plane is not well resolved by the spacecraft and b) the turbulence is anisotropic with increasing power with larger θ_{vB} [67, 70–72]. The lack of circular polarization signatures when θ_{vB} is moderately oblique is consistent with sampling effects of quasi-parallel waves at oblique angles in anisotropic turbulence [40], suggesting that ion-scale waves can persist at oblique θ_{vB} . In order to estimate the parallel propagating left-hand polarized spectrum, wavelet power with $\sigma_B > 0.9$ is identified when $\theta_{vB} < 15^\circ$. We assume homogeneity, and stationarity, such that the circularly polarized spectrum measured at $\theta_{vB} < 15^\circ$ represents the wave spectrum at all times (i.e. when $\theta_{vB} > 15^\circ$). Fig 1(c) shows the power spectrum of circularly polarized fluctuations with $\sigma_B > 0.9$ and $\sigma_B < -0.9$, corresponding to strong left and right-handed power. The right handed modes have been shown to be statistically consistent with a fast magnetosonic mode [69].

Distribution Functions Fig 2 shows the SPANi proton distribution function $f_p(v_\perp, v_\parallel)$ from the stream at 2020-01-30/04:10:21. Figure 2(a) shows an interpolation of the 3D measurements in the $v_\perp - v_\parallel$ plane constructed by identifying values of v_\perp and v_\parallel for each 3D energy




FIG. 2: (a) Interpolation of $f_p(\mathbf{v})$ from SPANi in the $v_{\perp} - v_{\parallel}$ plane, with the mean magnetic field pointing vertically. Points show SPANi measurement locations. Solid lines show $v_{\parallel th}$, and set of cyclotron-resonant diffusion contours are plotted. (b) Drifting bi-Maxwellian fit to $f_p(\mathbf{v})$. (c) Hermite representation of $f_p(\mathbf{v})$. Integration of $f_p(\mathbf{v})$ in a-c over $v_{\perp} - v_{\parallel}$ is normalized to the QTN density. (d) Contours of $f_p(\mathbf{v})$ determined by interpolating the gyrotropic distribution along QL diffusion contours for parallel cyclotron resonance. (e-f) Drifting bi-Maxwellian and the Hermite representations of $f_p(\mathbf{v})$ evaluated along cyclotron resonance contours.

bin, assuming gyrotropy. A drifting two-component bi-Maxwellian fit, assuming gyrotropy, is shown in Fig 2(b). The drifting bi-Maxwellian fit provides an approximation to $f_p(\mathbf{v})$ using two individual proton populations, though this parametrization may not resolve all non-thermal features that affect resonant interaction with cyclotron waves. Indeed, the presence of strong and persistent cyclotron-resonant interactions should affect the shape of $f_p(\mathbf{v})$, leading to an equilibrium distribution that deviates from a bi-Maxwellian [44]. To explore a nonpara-

metric $f_p(v_{\perp}, v_{\parallel})$, which may better represent the data [73–75], we fit a set of orthonormal Hermite functions using linear least square methods:

$$f_p(v_{\perp}, v_{\parallel}) = \sum_{m,n} g_{mn} \phi_m(v_{\perp}/v_{\perp th}) \phi_n(v_{\parallel}/v_{\parallel th}) \quad (2)$$

$$H_n(v) = (-1)^n e^{v^2} \frac{d^n}{dx^n} e^{-v^2} \quad (3)$$

$$\phi_m = \frac{H_m(v)}{\sqrt{2^m \pi^{1/2} m!}} e^{-v^2} \quad (4)$$

[76, 77]. Fig 2(c) shows the best-fit estimate to $f_p(\mathbf{v})$ for Hermite functions of order $m_{max} = 6, n_{max} = 6$. The distributions are shown in field aligned coordinates with $\hat{\mathbf{B}}_0$ along the vertical axis. In carrying out this fit, we effectively extend $f(v_{\perp}, v_{\parallel})$ to negative values of v_{\perp} by treating $f(v_{\perp}, v_{\parallel})$ as an even function of v_{\perp} , thereby omitting the terms in the sum corresponding to odd values of m . Our use of Hermite functions is meant solely as a nonparametric interpolative scheme, and is not intended to represent a natural-basis for $f_p(\mathbf{v})$ [76]. Over the studied interval, the average χ^2 residual of the Hermite representation is 90% of the bi-Maxwellian fit. No intervals were significantly better fit by the drifting bi-Maxwellian, though some distributions are equally well represented by either approximation.

The ion cyclotron resonance condition is $\omega(k_{\parallel}) = \Omega + k_{\parallel} v_{\parallel}$ such that outward-propagating cyclotron waves resonate with the inward propagating portion of the distribution function. The evolution of $f_p(\mathbf{v})$ in the presence of resonant interactions can be described by QL diffusion theory [43, 44]. In a reference frame moving with a wave, the particles conserve kinetic energy as they scatter off that wave, tracing contours in v_{\perp} and v_{\parallel} that can be computed using the wave dispersion relation and resonance condition [29, 43, 44, 46]. The QL diffusion contours [44, 78] are overlaid on $f_p(\mathbf{v})$ in Fig 2(a-c). If $f_p(\mathbf{v})$ decreases as v_{\perp} increases along the contours, cyclotron resonance diffuses energy across in the region where resonant waves are present, heating the plasma. Conversely, if $f_p(\mathbf{v})$ increases as v_{\perp} increases, then $f_p(\mathbf{v})$ is unstable, generating waves and cooling the plasma. Cyclotron resonant equilibrium corresponds to a flattening of $f_p(\mathbf{v})$ along the contours. Fig 2(d-f) shows $f_p(\mathbf{v})$ evaluated along QL diffusion contours, parameterized on v_{\perp} ; for each representation, $f_p(\mathbf{v})$ is characteristically flat along contours, suggesting that $f_p(\mathbf{v})$ has been processed by cyclotron-resonance [46, 48].

The QL proton heating rate due to resonance with parallel-propagating cyclotron waves is given by

$$\mathcal{H} = \int \frac{m_p v^2}{2} \frac{\partial f_p(\mathbf{v})}{\partial t} d^3 \mathbf{v} = \frac{\pi e^2}{2m_p^2} \int_0^{\infty} dk_{\parallel} \frac{1}{v_{\perp}} \hat{G}_k v_{\perp} \delta(\omega_k - k_{\parallel} v_{\parallel} - \Omega_p) \frac{\omega_k^2}{k_{\parallel}^2 c^2} I(k_{\parallel}) \hat{G}_k f_p(\mathbf{v}) d^3 \mathbf{v} \quad (5)$$

with

$$\hat{G}_k = \left(1 - \frac{k_{\parallel} v_{\parallel}}{\omega}\right) \frac{\partial}{\partial v_{\perp}} + \frac{k_{\parallel} v_{\perp}}{\omega} \frac{\partial}{\partial v_{\parallel}} \quad (6)$$

[43]. Using the observed left-handed spectrum in Fig 1, the cold-plasma dispersion relation, along with the bulk velocity to correct for Doppler shift [69], an average parallel left-handed cyclotron spectrum $I(k_{\parallel})$ is established.

For each observed $f_p(\mathbf{v})$, a value of \mathcal{H} is obtained numerically through Eq 5 using bi-Maxwellian and Hermite representations of $f(v_{\perp}, v_{\parallel})$. For the distribution shown in Fig 2, a heating rate of 10^{-14} W/m³ is found using the Hermite representation and 4×10^{-15} W/m³ using the drifting bi-Maxwellian spectrum. The measured \mathcal{H} is similar to estimates of bulk ion heating due to turbulent dissipation at the spacecraft’s location ($30R_{\odot}$) [79–81].

Fig 3(a-d) shows the differential volumetric heating rate \mathcal{H} as a function of resonant parallel velocity measured in each distribution function. The top panels, Fig 3(a-b), show positive \mathcal{H} , the “heating” rate, as a function of v_{\parallel} and time for the bi-Maxwellian (a) and Hermite (b) representations. The bottom panels Fig 3(c-d) shows negative \mathcal{H} , the “cooling” rate over the interval as a function of resonant v_{\parallel} due to emission of waves through instability. Fig 3(e) shows the net integrated \mathcal{H} for each measured distribution function. The integrated \mathcal{H} is uniformly positive, indicating that cyclotron waves present in the plasma are likely absorbed. There is very little cyclotron resonant emission from this plasma. However, the Hermite representation shows that cyclotron-instability, when present, is focused at the parallel thermal speed, $v_{\parallel th}$. The median heating rate is 3×10^{-15} W/m³ for the bi-Maxwellian fits, and 1×10^{-14} W/m³ for the Hermite representation. Using third order moments of the inertial range turbulence, [82], a cascade rate of 4.7×10^{-14} W/m³ is measured, which is consistent with previous measurements of the energy cascade rate at a similar radius [83, 84]. The measured cyclotron heating \mathcal{H} ranges from approximately 10-20% percent of the measured cascade rate. While uncertainties on the cascade rate exist due to the assumption of stationarity, isotropy, and homogeneity [82], previous work has suggested that the cascade rate estimates through third-order moments are accurate to a factor of approximately two or three [83, 85, 86].

Observations from SPANi are partial measurements of $f_p(\mathbf{v})$ and are subject to both uncertainties and ongoing calibration work. However, during this interval, $f_p(\mathbf{v})$ is well resolved, and while uncertainties in $f_p(\mathbf{v})$ will dominate uncertainties in our measured heating rates, we argue that the measurements reliably suggest net energy transfer from waves to the particles. Specifically, the biMaxwellian fit removes fine-structure in $f_p(\mathbf{v})$ that is present in the measurements and replicated by the Hermite function; importantly, we find that removing fine structure (i.e. the bi-Maxwellian fit) produces heating




FIG. 3: (a) Differential volumetric heating rates (positive \mathcal{H}) as a function of resonant parallel velocity computed from observed cyclotron spectra and drifting biMaxwellian fit to ion-distribution functions. The total heating rate is the sum over v_{\parallel} . Solid lines show the average $v_{\parallel th}$ for the interval (49 km/s). (b) Measured heating rates for a Hermite function approximation. (c-d) Corresponding cooling rates (negative \mathcal{H}). (e) Distribution of total \mathcal{H} for biMaxwellian and Hermite representations; respective lines show median value of \mathcal{H} . The turbulent cascade rate estimated from third-order moments is shown in green.

rates of the same sign and order of magnitude as when fine structure in $f_p(\mathbf{v})$ is included. In essence, while fine structure in $f_p(\mathbf{v})$ is observed by SPAN, it neither drastically affects the order of magnitude, nor, importantly, the sign of the cyclotron heating rate, \mathcal{H} . However, these results highlight the importance of modeling fine structure and local gradients in $f_p(\mathbf{v})$, e.g. with Hermite functions, which may yield heating rates more than double those found by smoother approximations to $f_p(\mathbf{v})$, e.g. a bi-Maxwellian fit.

Discussion Cyclotron resonance may play a significant role in shaping observed magnetic field spectra and distribution functions [8, 9, 23, 42, 46, 48, 52] observed in the solar wind. While flattening along QL diffusion contours has been previously reported [46, 48, 59] our observations directly couple an observed spectrum of cyclotron waves to heating rates in measured distribution functions. Our measured heating rates are on the order of the measured energy cascade rates ($\sim 10^{-14}$ W/m³), obtained through third order moments of the observed turbulence [82]; while this estimate may have significant uncertainty, we find good agreement with previous esti-

mates of the cascade rate [83, 84]. We furthermore show that incorporating fine, nonthermal, structures in the distribution function using a Hermite functional decomposition introduces relatively little effect on the extent or sign of measured cyclotron heating. We thus argue that the measured levels of cyclotron heating provide significant evidence for the mediation of turbulent dissipation through cyclotron resonance that is potentially sufficient to power the solar wind [79–81]. Studying the radial scaling of the turbulent energy cascade alongside quasilinear cyclotron heating rates promises to further constrain cyclotron resonance as a dissipation process.

There are several sources of uncertainty in this analysis, rising predominantly in the estimate of the cascade rate and in the gradients of $f_p(\mathbf{v})$ due to limited resolution. Furthermore, if the occurrence rates of cyclotron waves decreases with θ_{vB} , then the heating rate may be limited at oblique θ_{vB} . Additionally, there is the potential that heating by oblique kinetic Alfvén waves or oblique cyclotron waves may generate parallel cyclotron waves as a secondary process [87, 88]. The spectrum of oblique cyclotron waves is difficult to distinguish due to the strong anisotropy of the background turbulence [40], though future work will explore signatures of oblique cyclotron resonance. In any case, observed ion distributions are often flat along the quasi-parallel cyclotron diffusion contours and are rarely unstable to the growth of the waves. This flattening suggests that even if other physical processes contribute to bulk heating, the parallel-cyclotron resonance [43, 44] plays a significant role in shaping the distribution functions.

The measured heating rate indicates a near total lack of cyclotron emission through instabilities; thus, the origin of cyclotron waves remains an important unresolved point. Our observations show that 95% of the time left handed signatures are present, the net heating rate is positive, suggesting absorption of waves [53]. We note the studied interval does not have strong cyclotron wave storms [40, 57], though application of our method to a similar interval with more significant cyclotron wave events similarly suggests net heating. There are two main possible physical origins for these Alfvén/ion cyclotron waves. First, it is possible they are excited by beam instabilities [89], though recent work has suggested that dominant instability associated with the strong beam is associated with right-handed modes [57, 58]. Second, they may be generated by turbulence, though canonical theories of Alfvénic nonlinearity preferentially transport energy to large k_\perp but not large k_\parallel [90, 91], which is a hurdle to the turbulent generation of cyclotron resonant waves. However, recent work suggests that imbalance, i.e. the dominance of the outward Alfvén mode, may prevent energy from cascading to kinetic scales [92]. Fully kinetic simulations in the presence of such a barrier [88] show the generation of quasi-parallel cyclotron modes, similar to those observed in the solar wind, providing a

novel method for generating cyclotron waves that is consistent with a variety of observations [39, 40, 65, 93–98]. Further work is required to specifically investigate the origin of the observed cyclotron waves and their connection to turbulence, though our observation of localized instability at the proton-core thermal speed (Fig 3) may be a clue regarding the origin of the waves and their connection to the net heating measured in this study.

This work explicitly shows that the measured distribution functions in the solar wind contain evidence of cyclotron resonant heating at a level that may power the expanding solar wind. These results are significant step towards understanding the underlying physics of collisionless heating and a kinetic description of astrophysical plasmas.

TAB was supported by NASA PSP-GI Grant 80NSSC21K1771. KGK was supported by NASA ECIP Grant 80NSSC19K0912. BDGC was supported in part by NASA grant 80NSSC19K0829. JLV was supported by NASA grant NNH20ZDA001N-PSPGI. LDW was supported by STFC consolidated grant ST/S000364/1. DD acknowledges NSFC grant 42204166. The authors additionally acknowledge PSP/SWEAP & FIELDS contract NNN06AA01C.

* Electronic address: tbowen@berkeley.edu

- [1] Eckart Marsch. Kinetic Physics of the Solar Corona and Solar Wind. *Living Reviews in Solar Physics*, 3(1):1, July 2006. doi: 10.12942/lrsp-2006-1.
- [2] Jr. Coleman, Paul J. Turbulence, Viscosity, and Dissipation in the Solar-Wind Plasma. *The Astrophysical Journal*, 153:371, Aug 1968. doi: 10.1086/149674.
- [3] W. H. Matthaeus and M. L. Goldstein. Measurement of the rugged invariants of magnetohydrodynamic turbulence in the solar wind. *Journal of Geophysical Research*, 87:6011–6028, Aug 1982. doi: 10.1029/JA087iA08p06011.
- [4] S. R. Cranmer and A. A. van Ballegoijen. Alfvénic Turbulence in the Extended Solar Corona: Kinetic Effects and Proton Heating. *The Astrophysical Journal*, 594(1): 573–591, Sep 2003. doi: 10.1086/376777.
- [5] Benjamin D. G. Chandran, Bo Li, Barrett N. Rogers, Eliot Quataert, and Kai Germaschewski. Perpendicular Ion Heating by Low-frequency Alfvén-wave Turbulence in the Solar Wind. *The Astrophysical Journal*, 720(1): 503–515, Sep 2010. doi: 10.1088/0004-637X/720/1/503.
- [6] A. Barnes. Collisionless Damping of Hydromagnetic Waves. *Physics of Fluids*, 9:1483–1495, August 1966. doi: 10.1063/1.1761882.
- [7] A. Barnes. *Hydromagnetic waves and turbulence in the solar wind*, volume 1, pages 249–319. 1979.
- [8] M. L. Goldstein, D. A. Roberts, and C. A. Fitch. Properties of the fluctuating magnetic helicity in the inertial and dissipation ranges of solar wind turbulence. *Journal of Geophysical Research*, 99(A6):11519–11538, Jun 1994. doi: 10.1029/94JA00789.
- [9] Robert J. Leamon, William H. Matthaeus, Charles W.

- Smith, and Hung K. Wong. Contribution of Cyclotron-resonant Damping to Kinetic Dissipation of Interplanetary Turbulence. *The Astrophysical Journal Letters*, 507(2):L181–L184, Nov 1998. doi: 10.1086/311698.
- [10] G. G. Howes, J. M. Tenborge, W. Dorland, E. Quataert, A. A. Schekochihin, R. Numata, and T. Tatsuno. Gyrokinetic Simulations of Solar Wind Turbulence from Ion to Electron Scales. *Physical Review Letters*, 107(3):035004, Jul 2011. doi: 10.1103/PhysRevLett.107.035004.
- [11] B. D. G. Chandran, D. Verscharen, E. Quataert, J. C. Kasper, P. A. Isenberg, and S. Bourouaine. Stochastic Heating, Differential Flow, and the Alpha-to-proton Temperature Ratio in the Solar Wind. *ApJ*, 776(1):45, October 2013. doi: 10.1088/0004-637X/776/1/45.
- [12] Alfred Mallet, Alexander A. Schekochihin, and Benjamin D. G. Chandran. Disruption of Alfvénic turbulence by magnetic reconnection in a collisionless plasma. *Journal of Plasma Physics*, 83(6):905830609, Dec 2017. doi: 10.1017/S0022377817000812.
- [13] Nuno F. Loureiro and Stanislav Boldyrev. Role of Magnetic Reconnection in Magnetohydrodynamic Turbulence. *Physical Review Letters*, 118(24):245101, Jun 2017. doi: 10.1103/PhysRevLett.118.245101.
- [14] Eliot Quataert and Andrei Gruzinov. Turbulence and Particle Heating in Advection-dominated Accretion Flows. *The Astrophysical Journal*, 520(1):248–255, Jul 1999. doi: 10.1086/307423.
- [15] Benjamin D. G. Chandran, Peera Pongkitiwanchakul, Philip A. Isenberg, Martin A. Lee, Sergei A. Markovskii, Joseph V. Hollweg, and Bernard J. Vasquez. Resonant Interactions Between Protons and Oblique Alfvén/Ion-cyclotron Waves in the Solar Corona and Solar Flares. *ApJ*, 722(1):710–720, October 2010. doi: 10.1088/0004-637X/722/1/710.
- [16] Benjamin D. G. Chandran, Timothy J. Dennis, Eliot Quataert, and Stuart D. Bale. Incorporating Kinetic Physics into a Two-fluid Solar-wind Model with Temperature Anisotropy and Low-frequency Alfvén-wave Turbulence. *The Astrophysical Journal*, 743(2):197, Dec 2011. doi: 10.1088/0004-637X/743/2/197.
- [17] O. Alexandrova, C. Lacombe, A. Mangeney, R. Grappin, and M. Maksimovic. Solar Wind Turbulent Spectrum at Plasma Kinetic Scales. *ApJ*, 760(2):121, December 2012. doi: 10.1088/0004-637X/760/2/121.
- [18] John D. Richardson, Karolen I. Paularena, Alan J. Lazarus, and John W. Belcher. Radial evolution of the solar wind from IMP 8 to Voyager 2. *Geophys. Research Letters*, 22(4):325–328, Feb 1995. doi: 10.1029/94GL03273.
- [19] Lorenzo Matteini, Simone Landi, Petr Hellinger, Filippo Pantellini, Milan Maksimovic, Marco Velli, Bruce E. Goldstein, and Eckart Marsch. Evolution of the solar wind proton temperature anisotropy from 0.3 to 2.5 AU. *GRL*, 34(20):L20105, October 2007. doi: 10.1029/2007GL030920.
- [20] Eliot Quataert. Particle Heating by Alfvénic Turbulence in Hot Accretion Flows. *The Astrophysical Journal*, 500(2):978–991, Jun 1998. doi: 10.1086/305770.
- [21] Robert J. Leamon, Charles W. Smith, Norman F. Ness, William H. Matthaeus, and Hung K. Wong. Observational constraints on the dynamics of the interplanetary magnetic field dissipation range. *Journal of Geophysical Research*, 103(A3):4775–4788, Mar 1998. doi: 10.1029/97JA03394.
- [22] S. Peter Gary. Collisionless dissipation wavenumber: Linear theory. *JGR*, 104(A4):6759–6762, April 1999. doi: 10.1029/1998JA900161.
- [23] R. J. Leamon, W. H. Matthaeus, C. W. Smith, G. P. Zank, D. J. Mullan, and S. Oughton. MHD-driven Kinetic Dissipation in the Solar Wind and Corona. *The Astrophysical Journal*, 537(2):1054–1062, Jul 2000. doi: 10.1086/309059.
- [24] G. G. Howes, S. C. Cowley, W. Dorland, G. W. Hammett, E. Quataert, and A. A. Schekochihin. A model of turbulence in magnetized plasmas: Implications for the dissipation range in the solar wind. *Journal of Geophysical Research (Space Physics)*, 113(A5):A05103, May 2008. doi: 10.1029/2007JA012665.
- [25] Thomas Howard Stix. *Waves in plasmas*. 1992.
- [26] Joseph V. Hollweg and Walter Johnson. Transition region, corona, and solar wind in coronal holes: Some two-fluid models. *JGR*, 93(A9):9547–9554, September 1988. doi: 10.1029/JA093iA09p09547.
- [27] C. Y. Tu and E. Marsch. Two-Fluid Model for Heating of the Solar Corona and Acceleration of the Solar Wind by High-Frequency Alfvén Waves. *Sol Phys*, 171(2):363–391, April 1997. doi: 10.1023/A:1004968327196.
- [28] Steven R. Cranmer. Ion Cyclotron Wave Dissipation in the Solar Corona: The Summed Effect of More than 2000 Ion Species. *The Astrophysical Journal*, 532(2):1197–1208, Apr 2000. doi: 10.1086/308620.
- [29] Joseph V. Hollweg and Philip A. Isenberg. Generation of the fast solar wind: A review with emphasis on the resonant cyclotron interaction. *Journal of Geophysical Research (Space Physics)*, 107(A7):1147, July 2002. doi: 10.1029/2001JA000270.
- [30] Steven R. Cranmer. Ensemble Simulations of Proton Heating in the Solar Wind via Turbulence and Ion Cyclotron Resonance. *ApJS*, 213(1):16, July 2014. doi: 10.1088/0067-0049/213/1/16.
- [31] J. L. Kohl, G. Noci, E. Antonucci, G. Tondello, M. C. E. Huber, L. D. Gardner, P. Nicolosi, L. Strachan, S. Fineschi, J. C. Raymond, M. Romoli, D. Spadaro, A. Panasyuk, O. H. W. Siegmund, C. Benna, A. Ciaravella, S. R. Cranmer, S. Giordano, M. Karovska, R. Martin, J. Michels, A. Modigliani, G. Naletto, C. Pernechele, G. Poletto, and P. L. Smith. First Results from the SOHO Ultraviolet Coronagraph Spectrometer. *Sol Phys*, 175(2):613–644, October 1997. doi: 10.1023/A:1004903206467.
- [32] J. L. Kohl, G. Noci, E. Antonucci, G. Tondello, M. C. E. Huber, S. R. Cranmer, L. Strachan, A. V. Panasyuk, L. D. Gardner, M. Romoli, S. Fineschi, D. Dobrzycka, J. C. Raymond, P. Nicolosi, O. H. W. Siegmund, D. Spadaro, C. Benna, A. Ciaravella, S. Giordano, S. R. Habbal, M. Karovska, X. Li, R. Martin, J. G. Michels, A. Modigliani, G. Naletto, R. H. O’Neal, C. Pernechele, G. Poletto, P. L. Smith, and R. M. Suleiman. UVCS/SOHO Empirical Determinations of Anisotropic Velocity Distributions in the Solar Corona. *ApJL*, 501(1):L127–L131, July 1998. doi: 10.1086/311434.
- [33] Steven R. Cranmer, George B. Field, and John L. Kohl. Spectroscopic Constraints on Models of Ion Cyclotron Resonance Heating in the Polar Solar Corona and High-Speed Solar Wind. *ApJ*, 518(2):937–947, June 1999. doi: 10.1086/307330.
- [34] Lan K. Jian, Christopher T. Russell, Janet G. Luh-

- mann, Robert J. Strangeway, Jared S. Leisner, and Antoinette B. Galvin. Ion Cyclotron Waves in the Solar Wind Observed by STEREO Near 1 AU. *ApJL*, 701(2):L105–L109, August 2009. doi: 10.1088/0004-637X/701/2/L105.
- [35] J. J. Podesta and S. P. Gary. Magnetic Helicity Spectrum of Solar Wind Fluctuations as a Function of the Angle with Respect to the Local Mean Magnetic Field. *The Astrophysical Journal*, 734(1):15, Jun 2011. doi: 10.1088/0004-637X/734/1/15.
- [36] Jiansen He, Eckart Marsch, Chuanyi Tu, Shuo Yao, and Hui Tian. Possible Evidence of Alfvén-cyclotron Waves in the Angle Distribution of Magnetic Helicity of Solar Wind Turbulence. *The Astrophysical Journal*, 731(2):85, Apr 2011. doi: 10.1088/0004-637X/731/2/85.
- [37] R. T. Wicks, R. L. Alexander, M. Stevens, III Wilson, L. B., P. S. Moya, A. Viñas, L. K. Jian, D. A. Roberts, S. O’Modhrain, J. A. Gilbert, and T. H. Zurbuchen. A Proton-cyclotron Wave Storm Generated by Unstable Proton Distribution Functions in the Solar Wind. *ApJ*, 819(1):6, March 2016. doi: 10.3847/0004-637X/819/1/6.
- [38] Lloyd D. Woodham, Robert T. Wicks, Daniel Verscharen, Christopher J. Owen, Bennett A. Maruca, and Benjamin L. Alterman. Parallel-propagating Fluctuations at Proton-kinetic Scales in the Solar Wind Are Dominated By Kinetic Instabilities. *ApJL*, 884(2):L53, October 2019. doi: 10.3847/2041-8213/ab4adc.
- [39] SD Bale, ST Badman, JW Bonnell, TA Bowen, D Burgess, AW Case, CA Cattell, BDG Chandran, CC Chaston, CHK Chen, et al. Highly structured slow solar wind emerging from an equatorial coronal hole. *Nature*, pages 1–6, 2019.
- [40] Trevor A. Bowen, Alfred Mallet, Jia Huang, Kristopher G. Klein, David M. Malaspina, Michael Stevens, Stuart D. Bale, J. W. Bonnell, Anthony W. Case, Benjamin D. G. Chandran, C. C. Chaston, Christopher H. K. Chen, Thierry Dudok de Wit, Keith Goetz, Peter R. Harvey, Gregory G. Howes, J. C. Kasper, Kelly E. Korreck, Davin Larson, Roberto Livi, Robert J. MacDowall, Michael D. McManus, Marc Pulupa, J. L. Verniero, and Phyllis Whittlesey. Ion-scale Electromagnetic Waves in the Inner Heliosphere. *ApJS*, 246(2):66, February 2020. doi: 10.3847/1538-4365/ab6c65.
- [41] Philip A. Isenberg. Investigations of a turbulence-driven solar wind model. *JGR*, 95(A5):6437–6442, May 1990. doi: 10.1029/JA095iA05p06437.
- [42] Lloyd D. Woodham, Robert T. Wicks, Daniel Verscharen, and Christopher J. Owen. The Role of Proton Cyclotron Resonance as a Dissipation Mechanism in Solar Wind Turbulence: A Statistical Study at Ion-kinetic Scales. *The Astrophysical Journal*, 856(1):49, Mar 2018. doi: 10.3847/1538-4357/aab03d.
- [43] C. F. Kennel and F. Engelmann. Velocity Space Diffusion from Weak Plasma Turbulence in a Magnetic Field. *Physics of Fluids*, 9(12):2377–2388, December 1966. doi: 10.1063/1.1761629.
- [44] Philip A. Isenberg and Martin A. Lee. A dispersive analysis of bispherical pickup ion distributions. *JGR*, 101(A5):11055–11066, May 1996. doi: 10.1029/96JA00293.
- [45] Philip A. Isenberg. The kinetic shell model of coronal heating and acceleration by ion cyclotron waves: 2. Inward and outward propagating waves. *JGR*, 106(A12):29249–29260, December 2001. doi: 10.1029/2001JA000176.
- [46] E. Marsch and C. Y. Tu. Evidence for pitch angle diffusion of solar wind protons in resonance with cyclotron waves. *JGR*, 106(A5):8357–8362, May 2001. doi: 10.1029/2000JA000414.
- [47] E. Marsch and S. Bourouaine. Velocity-space diffusion of solar wind protons in oblique waves and weak turbulence. *Annales Geophysicae*, 29(11):2089–2099, November 2011. doi: 10.5194/angeo-29-2089-2011.
- [48] Jiansen He, Linghua Wang, Chuanyi Tu, Eckart Marsch, and Qiugang Zong. Evidence of Landau and Cyclotron Resonance between Protons and Kinetic Waves in Solar Wind Turbulence. *ApJ*, 800(2):L31, February 2015. doi: 10.1088/2041-8205/800/2/L31.
- [49] S. Peter Gary, Ruth M. Skoug, John T. Steinberg, and Charles W. Smith. Proton temperature anisotropy constraint in the solar wind: ACE observations. *GRL*, 28(14):2759–2762, January 2001. doi: 10.1029/2001GL013165.
- [50] S. Peter Gary, Lan K. Jian, Thomas W. Broiles, Michael L. Stevens, John J. Podesta, and Justin C. Kasper. Ion-driven instabilities in the solar wind: Wind observations of 19 March 2005. *Journal of Geophysical Research (Space Physics)*, 121(1):30–41, January 2016. doi: 10.1002/2015JA021935.
- [51] J. C. Kasper, A. J. Lazarus, and S. P. Gary. Hot Solar-Wind Helium: Direct Evidence for Local Heating by Alfvén-Cyclotron Dissipation. *PRL*, 101(26):261103, December 2008. doi: 10.1103/PhysRevLett.101.261103.
- [52] Daniele Telloni, Francesco Carbone, Roberto Bruno, Gary P. Zank, Luca Sorriso-Valvo, and Salvatore Mancuso. Ion Cyclotron Waves in Field-aligned Solar Wind Turbulence. *ApJL*, 885(1):L5, November 2019. doi: 10.3847/2041-8213/ab4c44.
- [53] D. Vech, M. M. Martinović, K. G. Klein, D. M. Malaspina, T. A. Bowen, J. L. Verniero, K. Paulson, T. Dudok de Wit, J. C. Kasper, J. Huang, M. L. Stevens, A. W. Case, K. Korreck, F. S. Mozer, K. A. Goodrich, S. D. Bale, P. L. Whittlesey, R. Livi, D. E. Larson, M. Pulupa, J. Bonnell, P. Harvey, K. Goetz, and R. MacDowall. Wave-particle energy transfer directly observed in an ion cyclotron wave. *A&AP*, 650:A10, June 2021. doi: 10.1051/0004-6361/202039296.
- [54] N. J. Fox, M. C. Velli, S. D. Bale, R. Decker, A. Driesman, R. A. Howard, J. C. Kasper, J. Kinnison, M. Kusterer, D. Lario, M. K. Lockwood, D. J. McComas, N. E. Raouafi, and A. Szabo. The solar probe plus mission: Humanity’s first visit to our star. *Space Science Reviews*, 204(1):7–48, Dec 2016. ISSN 1572-9672. doi: 10.1007/s11214-015-0211-6. URL <https://doi.org/10.1007/s11214-015-0211-6>.
- [55] S. D. Bale, K. Goetz, P. R. Harvey, P. Turin, J. W. Bonnell, T. Dudok de Wit, R. E. Ergun, R. J. MacDowall, M. Pulupa, M. Andre, M. Bolton, J.-L. Bougeret, T. A. Bowen, D. Burgess, C. A. Cattell, B. D. G. Chandran, C. C. Chaston, C. H. K. Chen, M. K. Choi, J. E. Connerney, S. Cranmer, M. Diaz-Aguado, W. Donakowski, J. F. Drake, W. M. Farrell, P. Ferreau, J. Fermin, J. Fischer, N. Fox, D. Glaser, M. Goldstein, D. Gordon, E. Hanson, S. E. Harris, L. M. Hayes, J. J. Hinze, J. V. Hollweg, T. S. Horbury, R. A. Howard, V. Hoxie, G. Jannet, M. Karlsson, J. C. Kasper, P. J. Kellogg, M. Kien, J. A. Klimchuk, V. V. Krasnoselskikh, S. Krucker, J. J. Lynch, M. Maksimovic, D. M. Malaspina, S. Marker, P. Martin, J. Martinez-Oliveros, J. McCauley, D. J. McComas,

- T. McDonald, N. Meyer-Vernet, M. Moncuquet, S. J. Monson, F. S. Mozer, S. D. Murphy, J. Odom, R. Oliverson, J. Olson, E. N. Parker, D. Pankow, T. Phan, E. Quataert, T. Quinn, S. W. Ruplin, C. Salem, D. Seitz, D. A. Sheppard, A. Siy, K. Stevens, D. Summers, A. Szabo, M. Timofeeva, A. Vaivads, M. Velli, A. Yehle, D. Werthimer, and J. R. Wygant. The FIELDS Instrument Suite for Solar Probe Plus. Measuring the Coronal Plasma and Magnetic Field, Plasma Waves and Turbulence, and Radio Signatures of Solar Transients. *Space Science Rev.*, 204:49–82, December 2016. doi: 10.1007/s11214-016-0244-5.
- [56] Justin C. Kasper, Robert Abiad, Gerry Austin, Marianne Balat-Pichelin, Stuart D. Bale, John W. Belcher, Peter Berg, Henry Bergner, Matthieu Berthomier, Jay Bookbinder, Etienne Brodu, David Caldwell, Anthony W. Case, Benjamin D. G. Chandran, Peter Cheimets, Jonathan W. Cirtain, Steven R. Cranmer, David W. Curtis, Peter Daigneau, Greg Dalton, Brahmananda Dasgupta, David DeTomaso, Millan Diaz-Aguado, Blagoje Djordjevic, Bill Donaskowski, Michael Effinger, Vladimir Florinski, Nichola Fox, Mark Freeman, Dennis Gallagher, S. Peter Gary, Tom Gauron, Richard Gates, Melvin Goldstein, Leon Golub, Dorothy A. Gordon, Reid Gurnee, Giora Guth, Jasper Halekas, Ken Hatch, Jacob Heerikuisen, George Ho, Qiang Hu, Greg Johnson, Steven P. Jordan, Kelly E. Korreck, Davin Larson, Alan J. Lazarus, Gang Li, Roberto Livi, Michael Ludlam, Milan Maksimovic, James P. McFadden, William Marchant, Bennet A. Maruca, David J. McComas, Luciana Messina, Tony Mercer, Sang Park, Andrew M. Peddie, Nikolai Pogorelov, Matthew J. Reinhart, John D. Richardson, Miles Robinson, Irene Rosen, Ruth M. Skoug, Amanda Slagle, John T. Steinberg, Michael L. Stevens, Adam Szabo, Ellen R. Taylor, Chris Tiu, Paul Turin, Marco Velli, Gary Webb, Phyllis Whittlesey, Ken Wright, S. T. Wu, and Gary Zank. Solar wind electrons alphas and protons (sweap) investigation: Design of the solar wind and coronal plasma instrument suite for solar probe plus. *Space Science Reviews*, 204(1):131–186, Dec 2016. ISSN 1572-9672. doi: 10.1007/s11214-015-0206-3. URL <https://doi.org/10.1007/s11214-015-0206-3>.
- [57] J. L. Verniero, D. E. Larson, R. Livi, A. Rahmati, M. D. McManus, P. Sharma Pyakurel, K. G. Klein, T. A. Bowen, J. W. Bonnell, B. L. Alterman, P. L. Whittlesey, David M. Malaspina, S. D. Bale, J. C. Kasper, A. W. Case, K. Goetz, P. R. Harvey, K. E. Korreck, R. J. MacDowall, M. Pulupa, M. L. Stevens, and T. Dudok de Wit. Parker Solar Probe Observations of Proton Beams Simultaneous with Ion-scale Waves. *ApJS*, 248(1):5, May 2020. doi: 10.3847/1538-4365/ab86af.
- [58] K. G. Klein, J. L. Verniero, B. Alterman, S. Bale, A. Case, J. C. Kasper, K. Korreck, D. Larson, E. Lichko, R. Livi, M. McManus, M. Martinović, A. Rahmati, M. Stevens, and P. Whittlesey. Inferred Linear Stability of Parker Solar Probe Observations Using One- and Two-component Proton Distributions. *ApJ*, 909(1):7, March 2021. doi: 10.3847/1538-4357/abd7a0.
- [59] J. L. Verniero, B. D. G. Chandran, D. E. Larson, K. Paulson, B. L. Alterman, S. Badman, S. D. Bale, J. W. Bonnell, T. A. Bowen, T. Dudok de Wit, J. C. Kasper, K. G. Klein, E. Lichko, R. Livi, M. D. McManus, A. Rahmati, D. Verscharen, J. Walters, and P. L. Whittlesey. Strong Perpendicular Velocity-space Diffusion in Proton Beams Observed by Parker Solar Probe. *ApJ*, 924(2):112, January 2022. doi: 10.3847/1538-4357/ac36d5.
- [60] T. A. Bowen et al. A merged search-coil and fluxgate magnetometer data product for parker solar probe fields. *JGR*, 125(5):e2020JA027813, 2020.
- [61] T. Dudok de Wit, V. V. Krasnoselskikh, O. Agapitov, C. Froment, A. Larosa, S. D. Bale, T. Bowen, K. Goetz, P. Harvey, G. Jannet, M. Kretzschmar, R. J. MacDowall, D. Malaspina, P. Martin, B. Page, M. Pulupa, and C. Revillet. First Results From the SCM Search-Coil Magnetometer on Parker Solar Probe. *Journal of Geophysical Research (Space Physics)*, 127(4):e30018, April 2022. doi: 10.1029/2021JA030018.
- [62] M. Pulupa, S. D. Bale, J. W. Bonnell, T. A. Bowen, N. Carruth, K. Goetz, D. Gordon, P. R. Harvey, M. Maksimovic, J. C. Martínez-Oliveros, M. Moncuquet, P. Saint-Hilaire, D. Seitz, and D. Sundkvist. The Solar Probe Plus Radio Frequency Spectrometer: Measurement requirements, analog design, and digital signal processing. *Journal of Geophysical Research (Space Physics)*, 122(3):2836–2854, March 2017. doi: 10.1002/2016JA023345.
- [63] F. Sahaoui, M. L. Goldstein, P. Robert, and Yu. V. Khotyaintsev. Evidence of a Cascade and Dissipation of Solar-Wind Turbulence at the Electron Gyroscale. *Physical Review Letters*, 102(23):231102, Jun 2009. doi: 10.1103/PhysRevLett.102.231102.
- [64] K. H. Kiyani, S. C. Chapman, Yu. V. Khotyaintsev, M. W. Dunlop, and F. Sahaoui. Global Scale-Invariant Dissipation in Collisionless Plasma Turbulence. *Physical Review Letters*, 103(7):075006, Aug 2009. doi: 10.1103/PhysRevLett.103.075006.
- [65] Trevor A. Bowen, Alfred Mallet, Stuart D. Bale, J. W. Bonnell, Anthony W. Case, Benjamin D. G. Chandran, Alexandros Chasapis, Christopher H. K. Chen, Die Duan, Thierry Dudok de Wit, Keith Goetz, Jasper S. Halekas, Peter R. Harvey, J. C. Kasper, Kelly E. Korreck, Davin Larson, Roberto Livi, Robert J. MacDowall, David M. Malaspina, Michael D. McManus, Marc Pulupa, Michael Stevens, and Phyllis Whittlesey. Constraining Ion-Scale Heating and Spectral Energy Transfer in Observations of Plasma Turbulence. *PRL*, 125(2):025102, July 2020. doi: 10.1103/PhysRevLett.125.025102.
- [66] Christopher Torrence and Gilbert P. Compo. A Practical Guide to Wavelet Analysis. *Bulletin of the American Meteorological Society*, 79(1):61–78, January 1998. doi: 10.1175/1520-0477(1998)079<textless{}0061:APGTWA<textgreater{}>2.0.CO;2.
- [67] G. G. Howes and E. Quataert. On the Interpretation of Magnetic Helicity Signatures in the Dissipation Range Of Solar Wind Turbulence. *The Astrophysical Journal Letters*, 709(1):L49–L52, Jan 2010. doi: 10.1088/2041-8205/709/1/L49.
- [68] Kristopher G. Klein, Gregory G. Howes, Jason M. TenBarge, and John J. Podesta. Physical Interpretation of the Angle-dependent Magnetic Helicity Spectrum in the Solar Wind: The Nature of Turbulent Fluctuations near the Proton Gyroradius Scale. *ApJ*, 785(2):138, April 2014. doi: 10.1088/0004-637X/785/2/138.
- [69] Trevor A. Bowen, Stuart D. Bale, J. W. Bonnell, Davin Larson, Alfred Mallet, Michael D. McManus, Forrest S. Mozer, Marc Pulupa, Ivan Y. Vasko, J. L. Verniero, Psp/Fields Team, and Psp/Sweap Teams. The Elec-

- tromagnetic Signature of Outward Propagating Ion-scale Waves. *ApJ*, 899(1):74, August 2020. doi: 10.3847/1538-4357/ab9f37.
- [70] R. W. Fredricks and F. V. Coroniti. Ambiguities in the deduction of rest frame fluctuation spectrums from spectrums computed in moving frames. *JGR*, 81(A31):5591–5595, November 1976. doi: 10.1029/JA081i031p05591.
- [71] Timothy S. Horbury, Miriam Forman, and Sean Oughton. Anisotropic Scaling of Magnetohydrodynamic Turbulence. *Phys. Rev. Lett.*, 101(17):175005, October 2008. doi: 10.1103/PhysRevLett.101.175005.
- [72] T. S. Horbury, R. T. Wicks, and C. H. K. Chen. Anisotropy in Space Plasma Turbulence: Solar Wind Observations. *Space Sci. Rev.*, 172(1-4):325–342, November 2012. doi: 10.1007/s11214-011-9821-9.
- [73] C. T. Dum, E. Marsch, and W. Pilipp. Determination of wave growth from measured distribution functions and transport theory. *Journal of Plasma Physics*, 23(1):91–113, February 1980. doi: 10.1017/S0022377800022170.
- [74] Adolfo F. Viñas and Chris Gurgiolo. Spherical harmonic analysis of particle velocity distribution function: Comparison of moments and anisotropies using Cluster data. *Journal of Geophysical Research (Space Physics)*, 114(A1):A01105, January 2009. doi: 10.1029/2008JA013633.
- [75] S. Servidio, A. Chasapis, W. H. Matthaeus, D. Perrone, F. Valentini, T. N. Parashar, P. Veltri, D. Gershman, C. T. Russell, B. Giles, S. A. Fuselier, T. D. Phan, and J. Burch. Magnetospheric Multiscale Observation of Plasma Velocity-Space Cascade: Hermite Representation and Theory. *Phys. Rev. Lett.*, 119(20):205101, November 2017. doi: 10.1103/PhysRevLett.119.205101.
- [76] Joseph T. Parker and Paul J. Dellar. Fourier-Hermite spectral representation for the Vlasov-Poisson system in the weakly collisional limit. *Journal of Plasma Physics*, 81(2):305810203, April 2015. doi: 10.1017/S0022377814001287.
- [77] A. A. Schekochihin, J. T. Parker, E. G. Highcock, P. J. Dellar, W. Dorland, and G. W. Hammett. Phase mixing versus nonlinear advection in drift-kinetic plasma turbulence. *Journal of Plasma Physics*, 82(2):905820212, April 2016. doi: 10.1017/S0022377816000374.
- [78] J. Squire, B. D. G. Chandran, and R. Meyrand. In-situ Switchback Formation in the Expanding Solar Wind. *ApJL*, 891(1):L2, March 2020. doi: 10.3847/2041-8213/ab74e1.
- [79] Benjamin D. G. Chandran and Jean C. Perez. Reflection-driven magnetohydrodynamic turbulence in the solar atmosphere and solar wind. *Journal of Plasma Physics*, 85(4):905850409, Aug 2019. doi: 10.1017/S0022377819000540.
- [80] Petr Hellinger, Pavel M. Trávníček, Štěpán Štverák, Lorenzo Matteini, and Marco Velli. Proton thermal energetics in the solar wind: Helios reloaded. *Journal of Geophysical Research (Space Physics)*, 118(4):1351–1365, April 2013. doi: 10.1002/jgra.50107.
- [81] K. Sasikumar Raja, Prasad Subramanian, Madhusudan Ingale, R. Ramesh, and Milan Maksimovic. Turbulent Proton Heating Rate in the Solar Wind from 5–45 R_{\odot} . *Astrophys. J.*, 914(2):137, June 2021. doi: 10.3847/1538-4357/abfcd1.
- [82] H. Politano and A. Pouquet. von Kármán-Howarth equation for magnetohydrodynamics and its consequences on third-order longitudinal structure and correlation functions. *Phys Rev E.*, 57(1):R21–R24, January 1998. doi: 10.1103/PhysRevE.57.R21.
- [83] Riddhi Bandyopadhyay, M. L. Goldstein, B. A. Maruca, W. H. Matthaeus, T. N. Parashar, D. Ruffolo, R. Chhiber, A. Usmanov, A. Chasapis, R. Qudsi, Stuart D. Bale, J. W. Bonnell, Thierry Dudok de Wit, Keith Goetz, Peter R. Harvey, Robert J. MacDowall, David M. Malaspina, Marc Pulupa, J. C. Kasper, K. E. Korreck, A. W. Case, M. Stevens, P. Whittlesey, D. Larson, R. Livi, K. G. Klein, M. Velli, and N. Raouafi. Enhanced Energy Transfer Rate in Solar Wind Turbulence Observed near the Sun from Parker Solar Probe. *ApJs*, 246(2):48, February 2020. doi: 10.3847/1538-4365/ab5dae.
- [84] N. Andrés, F. Sahraoui, S. Huang, L. Z. Hadid, and S. Galtier. The incompressible energy cascade rate in anisotropic solar wind turbulence. *A&A*, 661:A116, May 2022. doi: 10.1051/0004-6361/202142994.
- [85] Joshua E. Stawarz, Charles W. Smith, Bernard J. Vasquez, Miriam A. Forman, and Benjamin T. MacBride. The Turbulent Cascade and Proton Heating in the Solar Wind at 1 AU. *ApJ*, 697(2):1119–1127, June 2009. doi: 10.1088/0004-637X/697/2/1119.
- [86] K. T. Osman, M. Wan, W. H. Matthaeus, J. M. Weygand, and S. Dasso. Anisotropic Third-Moment Estimates of the Energy Cascade in Solar Wind Turbulence Using Multispacecraft Data. *Phys. Rev. Lett.*, 107(16):165001, October 2011. doi: 10.1103/PhysRevLett.107.165001.
- [87] Benjamin D. G. Chandran, Bo Li, Barrett N. Rogers, Eliot Quataert, and Kai Germaschewski. Perpendicular Ion Heating by Low-frequency Alfvén-wave Turbulence in the Solar Wind. *The Astrophysical Journal*, 720(1):503–515, Sep 2010. doi: 10.1088/0004-637X/720/1/503.
- [88] Jonathan Squire, Romain Meyrand, Matthew W. Kunz, Lev Arzamasskiy, Alexander A. Schekochihin, and Eliot Quataert. High-frequency heating of the solar wind triggered by low-frequency turbulence. *Nature Astronomy*, 6:715–723, March 2022. doi: 10.1038/s41550-022-01624-z.
- [89] Daniel Verscharen, Sofiane Bourouaine, and Benjamin D. G. Chandran. Instabilities Driven by the Drift and Temperature Anisotropy of Alpha Particles in the Solar Wind. *ApJ*, 773(2):163, August 2013. doi: 10.1088/0004-637X/773/2/163.
- [90] J. V. Shebalin, W. H. Matthaeus, and D. Montgomery. Anisotropy in MHD turbulence due to a mean magnetic field. *Journal of Plasma Physics*, 29(3):525–547, June 1983. doi: 10.1017/S0022377800000933.
- [91] P. Goldreich and S. Sridhar. Toward a Theory of Interstellar Turbulence. II. Strong Alfvénic Turbulence. *The Astrophysical Journal*, 438:763, Jan 1995. doi: 10.1086/175121.
- [92] R. Meyrand, J. Squire, A. A. Schekochihin, and W. Dorland. On the violation of the zeroth law of turbulence in space plasmas. *Journal of Plasma Physics*, 87(3):535870301, May 2021. doi: 10.1017/S0022377821000489.
- [93] Joshua E. Stawarz, Charles W. Smith, Bernard J. Vasquez, Miriam A. Forman, and Benjamin T. MacBride. The Turbulent Cascade for High Cross-Helicity States at 1 AU. *ApJ*, 713(2):920–934, April 2010. doi: 10.1088/0004-637X/713/2/920.
- [94] C. H. K. Chen, S. D. Bale, J. W. Bonnell, D. Borovikov, T. A. Bowen, D. Burgess, A. W. Case, B. D. G. Chandran, T. Dudok de Wit, K. Goetz, P. R. Harvey, J. C. Kasper, K. G. Klein, K. E. Korreck, D. Larson, R. Livi,

- R. J. MacDowall, D. M. Malaspina, A. Mallet, M. D. McManus, M. Moncuquet, M. Pulupa, M. L. Stevens, and P. Whittlesey. The Evolution and Role of Solar Wind Turbulence in the Inner Heliosphere. *ApJS*, 246(2):53, February 2020. doi: 10.3847/1538-4365/ab60a3.
- [95] Die Duan, Trevor A. Bowen, Christopher H. K. Chen, Alfred Mallet, Jiansen He, Stuart D. Bale, Daniel Vech, J. C. Kasper, Marc Pulupa, John W. Bonnell, Anthony W. Case, Thierry Dudok de Wit, Keith Goetz, Peter R. Harvey, Kelly E. Korreck, Davin Larson, Roberto Livi, Robert J. MacDowall, David M. Malaspina, Michael Stevens, and Phyllis Whittlesey. The Radial Dependence of Proton-scale Magnetic Spectral Break in Slow Solar Wind during PSP Encounter 2. *ApJS*, 246(2):55, February 2020. doi: 10.3847/1538-4365/ab672d.
- [96] Die D, Jiansen He, Trevor A. Bowen, Lloyd D. Woodham, Tieyan Wang, Christopher H. K. Chen, Alfred Mallet, and Stuart D. Bale. Anisotropy of Solar Wind Turbulence in the Inner Heliosphere at Kinetic Scales: PSP Observations. *ApJL*, 915(1):L8, July 2021. doi: 10.3847/2041-8213/ac07ac.
- [97] F. S. Mozer, O. V. Agapitov, S. D. Bale, J. W. Bonnell, T. A. Bowen, and I. Vasko. DC and Low-Frequency Electric Field Measurements on the Parker Solar Probe. *Journal of Geophysical Research (Space Physics)*, 125(9):e27980, September 2020. doi: 10.1029/2020JA027980.
- [98] Michael D. McManus, Trevor A. Bowen, Alfred Mallet, Christopher H. K. Chen, Benjamin D. G. Chandran, Stuart D. Bale, Davin E. Larson, Thierry Dudok de Wit, J. C. Kasper, Michael Stevens, Phyllis Whittlesey, Roberto Livi, Kelly E. Korreck, Keith Goetz, Peter R. Harvey, Marc Pulupa, Robert J. MacDowall, David M. Malaspina, Anthony W. Case, and J. W. Bonnell. Cross Helicity Reversals in Magnetic Switchbacks. *ApJS*, 246(2):67, February 2020. doi: 10.3847/1538-4365/ab6dce.