

A TWO-STEP MECHANISM FOR THE FOLDING OF ACTIN BY THE YEAST CYTOSOLIC CHAPERONIN

Sarah F Stuart^{1,2}, Robin J Leatherbarrow¹, Keith R Willison²

¹Institute of Chemical Biology, Imperial College London, London SW7 2AZ; ²Section of Cell and Molecular Biology, Institute of Cancer Research, London SW3 6JB.

Running head: Two-step mechanism for CCT-actin folding


Address correspondence to: K R Willison, 237 Fulham Road, London, SW3 6JB, UK.

Tel. 44-(0)207-878-3855; Email: keith.willison@icr.ac.uk

Actin requires the chaperonin containing TCP1 (CCT), a hexadecameric ATPase essential for cell viability in eukaryotes, to fold to its native state. Following binding of unfolded actin to CCT, the cavity of the chaperone closes and actin is folded and released in an ATP-dependent folding cycle. In yeast, CCT forms a ternary complex with the phosphatidylethanolamine transferase-like protein PLP2p to fold actin, and together they can return nascent or chemically denatured actin to its native state in a pure *in vitro* folding assay. The complexity of the CCT-actin system makes the study of the actin folding mechanism technically challenging. We have established a novel spectroscopic assay through selectively labeling the C-terminus of yeast actin with acrylodan and observe significant changes in the acrylodan fluorescence emission spectrum as actin is chemically unfolded and then refolded by the chaperonin. The variation in the polarity of the environment surrounding the fluorescent probe during the unfolding/folding processes has allowed us to monitor actin as it folds on CCT. The rate of actin folding at a range of temperatures and ATP concentrations has been determined for both wild type CCT and a mutant CCT, CCT4anc2 defective in folding actin *in vivo*. Binding of the non-hydrolysable ATP analogue AMP-PNP to the ternary complex leads to threefold faster release of actin from CCT following addition of ATP, suggesting a two-step folding process with a conformational change occurring upon closure of the cavity and a subsequent final folding step involving packing of the C-terminus to the native-like state.

The cytoskeletal protein actin is one of the most highly conserved in all eukaryotes and is involved in many essential cellular processes such as cell motility and cytokinesis. It exists in two forms: at low ionic concentrations, the G-actin monomer is stable, whereas in the presence of KCl, MgCl₂ or CaCl₂ and ATP, the F-actin

polymer predominates. The actin monomer consists of two domains, the so-called large domain and the small domain, which surround a nucleotide binding cleft and a high affinity divalent cation binding site (1). Actin can be further divided into subdomains, with the N- and C-termini co-located at the base of subdomain 1 (Fig. 1). G-actin can be unfolded thermally or chemically in the presence of denaturant or EDTA. Studies of the unfolding kinetics of rabbit skeletal muscle α -actin (ActA) with EDTA have shown that following loss of the cation and nucleotide from native actin, the actin then unfolds to an intermediate I₃, which cannot refold spontaneously (2) (Eq. 1). Unfolding actin by EDTA treatment allows folding studies to be performed under physiological conditions.



In eukaryotic cells, nascent actin is folded by the cytosolic chaperonin containing TCP1 (CCT or TRiC for TCP1-ring complex). CCT is a 1 MDa protein complex made up of two rings, each of which consists of 8 different subunits, known as CCT α – CCT θ in mammals and CCT1 – CCT8 in yeast. Recent studies of the CCT interactome have revealed a wide number of interacting proteins within the cell (3, 4), although only a limited number of obligate substrates have been characterized, including actin and tubulin as well as a number of WD40 repeat containing proteins (5). Actin folding by CCT *in vivo* in yeast requires the phosphatidylethanolamine transferase-like protein PLP2p (6), which contains a thioredoxin fold and forms a stable ternary structure with actin and CCT *in vitro* (7). The precise manner by which PLP2p assists actin folding is as yet unknown; PLP2p may be involved in the initial loading of actin onto CCT and ensuring that the correct engagement is reached, and/or it could have a role during folding or release. Although both EDTA-unfolded ActA and *S. cerevisiae* actin (Act1) can be bound by yeast CCT only Act1 can be productively refolded, indicating species-specific differences in actin folding behaviour and providing compelling evidence for the

absolute dependence of actin on CCT for its folding to the native state (8).

Here, we discuss the development of a spectroscopic actin folding assay which allows actin folding by CCT to be monitored in real time through labeling with the environmentally sensitive dye acrylodan. This approach allows greater sensitivity in the measurement of actin folding kinetics than gel electrophoresis-based assays, and has been used to observe the effects of ATP concentration and temperature on the rate of actin folding. Actin folding by CCT4anc2, a mutated form of CCT found in the temperature sensitive yeast strain *anc2-1* (9), has also been investigated. We present evidence for an actin folding intermediate within a functional folding cycle.

EXPERIMENTAL PROCEDURES

Purification and Labeling of Act1- Act1 was purified from *S. cerevisiae* lysate by affinity chromatography using a DNase I-Affigel column followed by ion exchange chromatography (10). The purified protein was eluted with approximately 220 mM KCl, and was polymerized by addition of 1 mM ATP and 2 mM MgCl₂ and incubation on ice for 30 min. A five-fold molar excess of Alexa Fluor 488-C₅-maleimide (Invitrogen) or a twenty-fold molar excess of acrylodan (Anaspec) was added to the polymerized actin and incubated on ice overnight. The labeled actin was centrifuged (2 h, 4°C, 100 000 x g) then the pellet was resuspended in 10 mM Tris pH 7.5, 0.5 mM β-mercaptoethanol, 0.2 mM CaCl₂, 50 μM ATP and dialyzed against the same buffer for 70 h at 4°C. After centrifugation, the supernatant was collected and the actin concentration and degree of labeling were calculated using the extinction coefficient 26 600 M⁻¹cm⁻¹ for actin at 290 nm (2), 18 500 M⁻¹cm⁻¹ for acrylodan at 385 nm (11), and 71 000 M⁻¹cm⁻¹ for Alexa Fluor 488 at 494 nm (2). Labeling efficiencies of 60 - 80 % and 92 % were obtained for ^{Acryl}Act1 and ⁴⁸⁸Act1 respectively.

Purification of CCT- CCT and CCT4anc2 were purified through a calmodulin binding peptide (CBP) tag inserted into subunit CCT3 (12). *S. cerevisiae* lysate expressing the tagged CCT was bound to calmodulin resin (Stratagene) and washed firstly with 20 mM HEPES pH 8, 1 M NaCl, 2 mM CaCl₂, 1 mM TCEP, 0.01 %v/v lauryldimethylamine oxide (LDAO), 20 %v/v glycerol, 5 mM ATP, 0.5 mM ADP, 15 mM MgCl₂, followed by two washes with 20 mM

HEPES pH 8, 150 mM KCl, 0.1 mM CaCl₂, 1 mM TCEP, 0.01 %v/v LDAO, 20 %v/v glycerol, 5 mM ATP, 0.5 mM ADP, 15 mM MgCl₂ and a final wash with 20 mM HEPES pH 8, 150 mM KCl, 0.1 mM CaCl₂, 1 mM TCEP, 0.01 %v/v LDAO, 20 %v/v glycerol. CCT was eluted with 20 mM HEPES pH 8, 150 mM KCl, 2 mM EGTA, 1 mM TCEP, 0.01 %v/v LDAO, 20 %v/v glycerol. To ensure that only the intact complex was isolated, CCT was further purified by sucrose gradient (20 mM HEPES pH 8, 150 mM KCl, 1 mM TCEP, 0.01 %v/v LDAO, 15 %v/v glycerol, 10 - 37.5 %w/v sucrose). The CCT containing fractions were concentrated, and the concentration was determined using the extinction coefficient 320 240 M⁻¹ cm⁻¹ at 280 nm (7).

Purification of PLP2p- Polyhistidine (6His)-tagged PLP2p was expressed in BL21(DE3) *E. coli* cells and purified by affinity chromatography using His-Spin Protein Miniprep columns (Zymo Research). After washing with His-Binding Buffer (50mM sodium phosphate pH 7.8, 300 mM NaCl, 10 mM imidazole, 0.03 % Triton X-100), PLP2p was eluted from the column with His-Elution Buffer (50mM sodium phosphate pH 7.8, 300 mM NaCl, 250 mM imidazole) and dialysed into 20mM HEPES pH 8, 75 mM KCl.

Polymerization of ^{Acryl}Act1- ^{Acryl}Act1 (8.5 μM, 0.35 mg/ml) was incubated at room temperature in the presence of 100 mM KCl, 2 mM MgCl₂, 1 mM ATP.

Unfolding of ^{Acryl}Act1- ^{Acryl}Act1 was diluted in 20mM HEPES pH 8, 75 mM KCl, 1 mM TCEP, 2 μM ATP to concentrations of no more than 18 μg/ml (440 nM). Unfolding was carried out at 24°C by addition of 1.5 mM EDTA.

⁴⁸⁸Act1 Folding Assay- ⁴⁸⁸Act1 (6.5 μg/ml, 160 nM) was unfolded in the presence of CCT (460 μg/ml, 460 nM) and PLP2p (15 μg/ml, 450 nM) in 20 mM HEPES pH 8, 75 mM KCl, 1 mM TCEP, 1.5 mM EDTA, 10 %v/v glycerol. Samples were protected from light and incubated at room temperature for 3 h. For release from CCT, 2 mM ATP and 10 mM MgCl₂ were added and the mixture was incubated at 30°C. Samples for gels were added to 20 mM HEPES pH 8, 10 %v/v glycerol, 1 mM CaCl₂ and kept on ice until loading onto a 6% native polyacrylamide gel containing 1 mM ATP. (7)

^{Acryl}Act Folding Assay- ^{Acryl}Act1 or ^{Acryl}ActA (14.4 μg/ml, 350 nM) was unfolded for 3 h in the presence of CCT or CCT4anc2 (400 μg/ml, 400 nM) and PLP2p (13.5 μg/ml, 400 nM) as for

⁴⁸⁸Act1. ATP and MgCl₂ were added so that the final concentrations were 1, 2 or 4 mM ATP, and 5, 10 or 20 mM MgCl₂ respectively, with the final glycerol concentration at 8 %v/v. Folding was monitored spectroscopically, either by repeatedly scanning the fluorescence emission of the sample or by monitoring emission at 470 nm for the duration of the reaction.

Co-polymerisation of ⁴⁸⁸Act1 and ^{Acryl}Act1 with unlabeled Act1- Refolded ^{Acryl}Act1 or ⁴⁸⁸Act1 was incubated overnight on ice with Act1 to a final Act1 concentration of 250 µg/ml (6 µM) in the presence of 115 mM KCl, 1.7 mM ATP and 6 mM MgCl₂. After centrifugation (2 h, 4°C, 100 000 x g), the pellet was analysed by SDS PAGE. *AMP-PNP Actin Folding Assay-* ^{Acryl}Act1 or ⁴⁸⁸Act1 were unfolded as described previously. Adenylyl imidodiphosphate (AMP-PNP) and MgCl₂ were added to final concentrations of 10 mM and 50 mM respectively, and samples were incubated at 30°C for 10 min. After this time, ATP or ADP and MgCl₂ were added so that the final concentrations were 8 mM AMP-PNP, 2 mM ATP or ADP, 50 mM MgCl₂, 70 mM KCl, 8 %v/v glycerol, 0.64 mM TCEP and folding was monitored either spectroscopically for ^{Acryl}Act1 or by native PAGE for ⁴⁸⁸Act1.

Fluorescence measurements- Fluorescence emission was measured with the PTI QuantaMaster 40 spectrofluorometer with a Haake DC10-K15 refrigerated circulator attached. The excitation wavelength used was 380 nm, and the excitation slits were set to 2 nm. Emission spectra were collected from 425 - 525 nm, and time-based emission was monitored at 470 nm with the emission slits set to 4 nm.

Data Analysis- The change in the intensity of acrylodan fluorescence emission at 470 nm over time was fitted to a single exponential using GraphPad Prism 5 (GraphPad Software), with the first 50 seconds of time-based scans being excluded since a mixing phase could be seen. Rate constants were calculated as the mean of at least three independent experiments with error bars representing the standard deviation.

RESULTS

Acrylodan Labeling and Chemical Unfolding of Act1. Dye labeling of Act1 at Cys374 has a large impact on the kinetics of chemical unfolding with EDTA, resulting in much more rapid unfolding of the labeled protein to the I₃ state in comparison to unlabeled Act1. This is a useful experimental property since yeast actin unfolds on a much slower time-scale than ActA, taking

several hours to denature completely (2, 7). Actin was selectively labeled at Cys374 by polymerization followed by incubation with Alexa Fluor 488 or acrylodan, a thiol-reactive derivative of Prodan. Cys374 is located in a flexible stretch of residues at the C-terminus of actin which packs against subdomain 1 in the crystal structure of the actin-rhodamine dye complex (1) (Fig. 1).

Prodan, which has an emission maximum of 531 nm in water (13), is highly sensitive to changes in its environment. In less polar solvents than water a blue shift in emission, accompanied by a large increase in the fluorescence intensity, occurs and this allows changes in the environment of the probe to be monitored spectroscopically. The emission of native acrylodan-labeled Act1 (^{Acryl}Act1_{NAT}) is dramatically different to that of EDTA-unfolded ^{Acryl}Act1 (^{Acryl}Act1_{I3}), which has a very similar emission spectrum to CCT-bound ^{Acryl}Act1 (Fig. 2A). The maximum emission wavelength for ^{Acryl}Act1_{NAT} is at 489 nm, and the Stokes shift of this form is 98 nm, in contrast to the EDTA-unfolded protein which exhibits maximum fluorescence emission at 463 nm and has a Stokes shift of 81 nm. Interestingly, the maximum emission wavelength of the unfolded ^{Acryl}Act1 is very similar to the value of 465 nm reported for acrylodan-labeled rabbit skeletal muscle F-actin (14), indicating that the environment in which the dye is found in F-actin is similar to that of the CCT-bound ^{Acryl}Act1. Within acrylodan-labeled yeast F-actin, fluorescence emission is also observed to be blue-shifted (Fig. 2B), although not to the same extent as that of the acrylodan-labeled rabbit F-actin (14). This could reflect only partial incorporation of ^{Acryl}Act1 into the filaments, however it is known that actin labeled in this manner is capable of polymerisation since that is how it is purified. This blue shift in fluorescence emission in the unfolded, CCT-bound and polymerized ^{Acryl}Act1 indicates that the probe is in a less polar environment in all these three states compared to the native G-actin monomer. The rate of EDTA unfolding of ^{Acryl}Act1_{NAT} was determined by fitting a single exponential to the fluorescence emission at 470 nm (Fig. 2C). This gives the observed rate constant for the unfolding reaction to be $1.48 \times 10^{-3} \pm 0.09 \times 10^{-3} \text{ s}^{-1}$ in the presence of the 10 % v/v glycerol required to stabilize CCT. Stopped flow analysis of the unfolding kinetics of rabbit skeletal muscle actin revealed a transition through

intermediates (2), such as the calcium free, ATP bound intermediate I_1 , which is not visible in this analysis of yeast actin unfolding. This could be due to the short time scale over which this transition takes place relative to the mixing time. Alternatively, it is also possible that the calcium-unbinding transition does not affect the local environment surrounding the fluorescent probe at the C-terminus to the same degree as the four tryptophan residues in subdomain 1 that constitute the entire intrinsic fluorescence signal of actin (2).

Release of Fluorescently Labeled Act1 from CCT. It has been shown that CCT can form a stable ternary complex with $^{488}\text{Act1}$ and PLP2p following chemical unfolding of $^{488}\text{Act1}_{\text{NAT}}$ with EDTA, and that upon addition of ATP and MgCl_2 native $^{488}\text{Act1}$ can be released in the pure *in vitro* folding assay system (7) (Fig. 3A). We then experimented with various environment-sensitive probes but found that some labeled actins behaved differently with respect to release from CCT; for example Act1, fluorescently labeled with 5-((2-((iodoacetyl)amino)ethyl)amino)naphthalene-1-sulfonic acid (IAEDANS), $^{\text{AEDANS}}\text{Act1}$, could be loaded onto CCT, but native $^{\text{AEDANS}}\text{Act1}$ could not be released from the complex (data not shown). In contrast, $^{\text{Acryl}}\text{Act1}$ could be loaded onto CCT and released in the same manner as $^{488}\text{Act1}$ (Fig. 3B). It can be seen by native polyacrylamide gel electrophoresis (native PAGE) that the folding reaction is complete within approximately 10 minutes, and around 90 % of actin is released. The refolded $^{\text{Acryl}}\text{Act1}$ can be bound by DNase I and vitamin D binding protein (VDBP) (Fig. 3B), and copolymerized with unlabeled actin (Fig. 3D), demonstrating that native G-actin is formed (7). Further support for the natural behaviour of the folding system used here is that it is unable to fold $^{\text{Acryl}}\text{ActA}$ to the native state (Fig. 3C). This had only previously been shown in the coupled IVT transcription/translation system (8) but is a property of this pure component system as well. Upon addition of ATP and MgCl_2 to the $^{\text{Acryl}}\text{Act1}$ -CCT-PLP2p complex, a decrease in acrylodan emission accompanied by a shift in the maximum emission wavelength is observed (Fig. 2A and 4A), thereby allowing the folding reaction to be monitored spectroscopically. This shift occurs over a similar time period to the release of native actin from CCT as observed by native PAGE, and as such can be used to make more accurate measurements of the rate of

release from the chaperonin. A single exponential can be fitted to the decrease in emission as $^{\text{Acryl}}\text{Act1}$ is released from CCT (Fig. 4A). The observed rate constant of release (k) of $^{\text{Acryl}}\text{Act1}$ from CCT over a range of temperatures was determined based on fluorescence emission at 470 nm. By fitting of these rate constants to the Arrhenius equation (Fig. 4B), the activation energy of actin release by CCT was found to be $34 \pm 8 \text{ kJ mol}^{-1}$, $32 \pm 8 \text{ kJ mol}^{-1}$ and $29 \pm 19 \text{ kJ mol}^{-1}$ for 4 mM, 2 mM and 1 mM ATP respectively. It can be seen that although both temperature and the concentration of ATP have an effect on the rate of release of $^{\text{Acryl}}\text{Act1}$, the temperature dependence appears similar over the different ATP concentrations indicating that the reaction is proceeding by the same mechanism. The spectroscopic folding assay was not performed using $^{488}\text{Act1}$ as only a very small change in the fluorescence emission is observed upon unfolding (2) and since a mixture of states exist during the folding process, the changes in intensity would be too small to accurately measure.

Release of Act1 from CCT4anc2. CCT4anc2 contains the mutation G345D in subunit CCT4, located in the external region of the apical domain, and causes actin defects in the temperature sensitive yeast strain *anc2-1* (9, 15). Using the spectroscopic folding assay, the rate of release of $^{\text{Acryl}}\text{Act1}$ by CCT4anc2 was found to be the same as for CCT at 2 mM ATP; however different behaviour was observed at the higher ATP concentration. At 4 mM ATP the rate of release of $^{\text{Acryl}}\text{Act1}$, while faster than from the mutant at 2 mM ATP, was slower than that from wild-type CCT. Furthermore, the rate of release increased with temperature up to 32.5°C , but at temperatures above this there was a subsequent decrease in the rate of release (Fig. 5), consistent with the heat-sensitive behaviour *in vivo* of CCT4anc2 mutant cells (9, 15).

Release of Act1 from CCT following addition of AMP-PNP. Addition of the non-hydrolysable ATP analogue AMP-PNP to CCT-bound $^{\text{Acryl}}\text{Act1}$ had no significant effect on the acrylodan emission spectrum. However, upon addition of ATP to a final concentration of 2 mM, actin release could be seen to occur at a faster rate than observed when ATP was added directly to the $^{\text{Acryl}}\text{Act1}$ -CCT complex in the absence of AMP-PNP (Fig. 6A). Fitting of a single exponential to the decrease in fluorescence gave an observed rate constant k of $11.8 \times 10^{-3} \pm 0.7 \times 10^{-3} \text{ s}^{-1}$, which is an

approximately three-fold increase relative to the release of ^{Acryl}Act1 from CCT without the pre-incubation step ($3.8 \times 10^{-3} \pm 0.3 \times 10^{-3} \text{ s}^{-1}$). Following addition of ADP in place of ATP, no release was observed (Fig. 6A). The same effect of pre-incubation of the loaded CCT complex with AMP-PNP was exhibited for release of ⁴⁸⁸Act1 as demonstrated by native PAGE analysis (Fig. 6, B and C), with virtually all ⁴⁸⁸Act1 released from CCT and returned to its native state within 5 minutes of addition of ATP compared with 10 to 15 minutes with ATP alone.

DISCUSSION

Conformational changes at the C-terminus of actin during folding. The selective labeling of actin at its C-terminal cysteine is useful for observing changes in conformation of this flexible region of the protein. Labeling of Cys374 with bulky fluorophores has an effect on the local structure of the amino acid within the actin molecule (16), with the label acting as a mutation which destabilizes the actin molecule sufficiently to allow chemical unfolding over a shorter time scale. It should be noted that chemically unfolded actin, not nascent actin, has been used in these assays, and the solution structures of these two denatured actin states, which are unknown, may well differ (12). Nonetheless, binding of the fluorescently labeled actin to DNase I and VDBP and its ability to polymerize demonstrates that neither the introduction of the fluorophore, nor the manner of unfolding of actin, prevents folding of actin to a functional monomer.

The proximity of the probe to the hinge region in the actin monomer makes it ideal for monitoring folding since EM structures demonstrate that actin bound to CCT takes on a more open structure, with the large and small domains binding in a 1,4 conformation across the cavity connected only by this hinge (17). It has also been hypothesised that reconfiguration of the C-terminus is an important step in the chaperone-mediated folding of actin (18). The contribution of the C-terminus to the stability of Act1 is clearly reflected in the variability of folding and unfolding behaviours depending on the nature of modification at Cys374. The inability of CCT to refold ^{AEDANS}Act1, while ⁴⁸⁸Act1 and ^{Acryl}Act1 can be refolded, demonstrates that the nature of the probe attached to the surface is very important and provides further evidence towards

the role of the C-terminus both in the folding and the stability of native actin.

Interestingly, the acrylodan moiety is located in a less polar environment in both the unfolded I₃ intermediate and in F-actin relative to G-actin. This demonstrates the differences in conformation between the monomeric and polymeric actin states, but also similarities between unfolded actin and F-actin. Perhaps F-actin re-explores the unfolding landscape of actin during its functional cycles as proposed by Altschuler and Willison (19). CCT subunits have been shown to associate with the F-actin filaments, which may indicate exposure of similar aggregation-prone, CCT-binding regions in both the unfolded and polymerized actin (20). *Actin Folding by CCT4anc2.* The G345D mutation in CCT4anc2 affects the inter-ring allostery of the chaperonin, and results in a slower rate of ATP hydrolysis (15). Actin folding by CCT4anc2 is defective *in vivo* above the permissive temperature of 30°C, with actin aggregates being observed in cells (9). Folding assays with radiolabeled human β-actin demonstrated that while the rate of actin folding by CCT4anc2 is similar to that of wild-type CCT, there is an approximately two-fold reduction in the yield of native actin (15). The data obtained from spectroscopic folding assay in general agrees with this, with the rate of actin release by the mutant affected by both changes in ATP concentration and temperature. A drop in the rate of actin release by CCT4anc2 is observed above the permissive temperature (highlighted in Fig. 5) and is more pronounced in the presence of 4 mM ATP compared to 2 mM ATP. We note that the intracellular concentrations of ATP in *S. cerevisiae* ranges from 1.1 to 4.5 mM (21) and therefore the maximum attainable level of activity of the enzyme may be being reached at 37°C in 4 mM ATP and hence the maximum effect of the anc2 mutation should be observed at these extremes (Fig. 5). Nevertheless it is striking that the range of activity varies only two-fold over this physiological relevant range of both nucleotide concentration and temperature. A lower yield of native actin is observed in the presence of CCT4anc2 compared to with CCT (data not shown), but it is unclear whether this is due to a decrease in loading of the unfolded actin, less effective release from the chaperonin or a combination of the two.

Mechanism of Actin Folding by CCT. The C-terminus of Act1 appears to be in a very similar

environment in both the unfolded ^{Acryl}Act1₁₃ and ^{Acryl}Act1-CCT based on the acrylodan emission spectra. This is the case for both the nucleotide-free and AMP-PNP bound ^{Acryl}Act1-CCT complexes, so it seems probable that no significant conformational rearrangement at the C-terminus of actin occurs upon binding of AMP-PNP and closure of the lid of the chaperonin (22). The differing kinetics of ATP-dependent release of native ^{Acryl}Act1 from CCT when pre-incubated with AMP-PNP relative to direct release from the complex indicates that a conformational change in actin does occur upon lid closure which corresponds to an initial folding step that is independent of the C-terminus of actin.

Neiryneck et al (18) developed a multi-step model for the CCT-mediated folding of human β -actin based on their analysis of mutants from an alanine scan of the entire actin polypeptide chain and previous structural electron microscopy models of CCT-actin complexes (17, 22). Their model proposes two major transitions. During transition 1, the CCT-captured, extended form of actin rotates about the interdomain hinge region and rebinds as a more compact actin mass, as observed after closure of the CCT cavity by AMP-PNP (22). During transition 2, the actin C-terminus which is still in direct contact with a CCT-subunit binding site, is released from CCT to establish its correct contacts with subdomain 1 of actin and this step completes folding. The kinetic behaviour of ^{Acryl}Act1 folding on CCT in the presence of AMP-PNP followed by ATP supports a sequential model of actin folding (Fig. 7). Firstly, incubation of the pre-assembled CCT-actin-PLP2 ternary complex in AMP-PNP allows actin to progress along its folding trajectory to a more closed state, consistent with both the structural (22) and Förster resonance energy transfer studies (23) and secondly the addition of ATP permits the correct packing of the actin C-terminus and thus release. We suggest that an ATP hydrolysis step by a particular CCT subunit(s) is required for C-terminal release which is why AMP-PNP does not facilitate this process.

The reduced polarity of the environment surrounding the probe in the ^{Acryl}Act1-CCT complex and the similarity of the emission spectra to that of acrylodan labeled rabbit F-actin (14) suggests that the C-terminus is unlikely to be 'free' in solution. Instead, it is likely to be fixed somewhere, probably on a particular CCT subunit binding site, and as such the acrylodan

fluorescence remains unaffected by the preliminary folding step induced by binding of AMP-PNP. The acrylodan moiety could be tethered to CCT in this preliminary folding step, or bound to the actin molecule itself. This CCT-actin folding intermediate complex can convert actin into its native form through rearrangement at the C-terminus of actin, perhaps locking the molecule into the correct conformation. This native or native-like conformation would be released from the post-ATP hydrolysis, ADP state of CCT which could no longer bind it as it was no longer in a non-native conformation.

Based on the relative rates with or without pre-incubation with AMP-PNP, it appears that a slow rearrangement takes place upon binding of the nucleotide to CCT. This rearrangement does not include the labeled C-terminus, and most likely would be observed as a lag phase in the initial stages of the folding reaction which could well be masked in this analysis by the mixing period observed initially. Development of the spectroscopic folding assay for analysis by stopped flow might allow further insight into this stage of the reaction. This is followed by a more rapid rearrangement of the C-terminus as ATP is hydrolysed.

The mechanism by which ATP displaces AMP-PNP in the chaperone complex is as yet unknown. Due to the variation in ATP binding sites across the subunits of CCT, it is possible that AMP-PNP is not bound at all the potential ATP binding sites within the chaperone. Binding and hydrolysis of ATP at an unoccupied site could then bring about the allosteric rearrangement required to instigate the final folding step and release of actin. Alternatively, the release process could be dependent on the dissociation rate of AMP-PNP and subsequent binding of ATP.

The folding of actin by CCT is undoubtedly an intricate process and is likely to be composed of many further rearrangements before the native actin monomer can be released, and more work is required to characterize the elaborate folding pathway of actin and the contribution of cofactors such as PLP2. Here we have shown evidence that an actin folding intermediate can be trapped within a functional folding cycle, elucidating the importance of the packing of the C-terminal of actin as a final step in the folding process in order to ensure release of native actin from CCT.

REFERENCES

1. Otterbein LR, Graceffa P, Dominguez R (2001) *Science* **293**(5530), 708-711
2. Altschuler G, Klug D, Willison K (2005) *J Mol Biol* **353**(2), 385-396
3. Dekker C, Stirling PC, McCormack EA, Filmore H, Paul A, Brost RL, Costanzo M, Boone C, Leroux MR, Willison KR (2008) *EMBO J* **27**(13), 1827-1839
4. Yam AY, Xia Y, Lin HTJ, Burlingame A, Gerstein M, Frydman J (2008) *Nat Struct Mol Biol* **15**(12), 1255-1262
5. Valpuesta JM, Martin-Benito J, Gomez-Puertas P, Carrascosa JL, Willison KR (2002) *FEBS Lett* **529**(1), 11-16
6. Stirling PC, Srayko M, Takhar KS, Pozniakovskiy A, Hyman AA, Leroux MR (2007) *Mol Biol Cell* **18**(6), 2336-2345
7. McCormack EA, Altschuler GM, Dekker C, Filmore H, Willison KR (2009) *J Mol Biol* **391**(1), 192-206
8. Altschuler GM, Dekker C, McCormack EA, Morris EP, Klug DR, Willison KR (2009) *FEBS Lett* **583**(4), 782-786
9. Vinh DB, Drubin DG (1994) *Proc Natl Acad Sci U S A* **91**(19), 9116-9120
10. Goode BL (2002) *Guide to Yeast Genetics and Molecular and Cell Biology, Pt C* **351**, 433-441
11. Roy P, Rajfur Z, Jones D, Marriott G, Loew L, Jacobson K (2001) *J Cell Biol* **153**(5), 1035-1048
12. Pappenberger G, McCormack EA, Willison KR (2006) *J Mol Biol* **360**(2), 484-496
13. Weber G, Farris FJ (1979) *Biochemistry* **18**(14), 3075-3078
14. Marriott G, Zechel K, Jovin TM (1988) *Biochemistry* **27**(17), 6214-6220
15. Shimon L, Hynes GM, McCormack EA, Willison KR, Horovitz A (2008) *J Mol Biol* **377**(2), 469-477
16. Yasunaga T, Wakabayashi T (2001) *J Biochem (Tokyo)* **129**(2), 201-204
17. Llorca O, McCormack EA, Hynes G, Grantham J, Cordell J, Carrascosa JL, Willison KR, Fernandez JJ, Valpuesta JM (1999) *Nature* **402**(6762), 693-696
18. Neirynek K, Waterschoot D, Vandekerckhove J, Ampe C, Rommelaere H (2006) *J Mol Biol* **355**(1), 124-138
19. Altschuler GM, Willison KR (2008) *Journal of the Royal Society Interface* **5**(29), 1391-1408
20. Grantham J, Ruddock LW, Roobol A, Carden MJ (2002) *Cell Stress Chaperon* **7**(3), 235-242
21. Larsson C, Nilsson A, Blomberg A, Gustafsson L (1997) *J Bacteriol* **179**(23), 7243-7250
22. Llorca O, Martin-Benito J, Grantham J, Ritco-Vonsovici M, Willison KR, Carrascosa JL, Valpuesta JM (2001) *EMBO J* **20**(15), 4065-4075
23. Villebeck L, Persson M, Luan S-L, Hammarstrom P, Lindgren M, Jonsson B-H (2007) *Biochemistry* **46**(17), 5083-5093

ABBREVIATIONS

488, Alexa Fluor 488; Acryl, Acrylodan; Act1, yeast actin; ActA, rabbit skeletal muscle α -actin; AMP-PNP, adenylyl-imidodiphosphate; CBP, calmodulin binding peptide; CCT, chaperonin containing TCP1; IAEDANS, 5-((2-((iodoacetyl)amino)ethyl)amino)naphthalene-1-sulfonic acid, PLP2, phosphocin-like protein 2; VDBP, Vitamin D binding protein.

ACKNOWLEDGEMENTS

Sarah Stuart is funded by the EPSRC through the Institute of Chemical Biology.

FIGURE LEGENDS

Fig. 1. The crystal structure of G-actin (1), with ADP (light blue) and Ca^{2+} (orange) bound. Actin is labelled at C374 with tetramethylrhodamine (purple).

Fig. 2. (A) Fluorescence emission spectra of $^{\text{Acryl}}\text{Act1}_{\text{NAT}}$ (---), $^{\text{Acryl}}\text{Act1}_{\text{I}_3}$ (—) and $^{\text{Acryl}}\text{Act1-CCT}$ (···) exciting at 380 nm; (B) Polymerisation of $^{\text{Acryl}}\text{Act1}$: fluorescence emission spectra of $^{\text{Acryl}}\text{Act1}$ immediately after addition of MgCl_2 and ATP (---) and after 10 minutes polymerisation (—); (C) Representative scan showing the change in acrylodan emission at 470 nm as $^{\text{Acryl}}\text{Act1}$ is unfolded in the presence of 10 %v/v glycerol.

Fig. 3. (A) and (B) Time course of ATP-dependent release of (A) $^{488}\text{Act1}$ and (B) $^{\text{Acryl}}\text{Act1}$ from CCT. Act1 is unfolded onto CCT in the presence of EDTA and released by addition of 2 mM ATP and 10 mM MgCl_2 . The time is shown in minutes following addition of ATP. In the final lane the formation of native actin is confirmed by binding to DNase I (D) and VDBP (V) as shown by a shift on the native gel. Coomassie stained native gels are shown on the left hand side as well as in-gel fluorescence of (A) $^{488}\text{Act1}$ (imaged on Typhoon 9410) and (B) $^{\text{Acryl}}\text{Act1}$ (imaged by UV lamp) of the same gel on the right hand side. Unlabelled PLP2 is visible as marked by *; (C) In-gel fluorescence by UV lamp of the release of $^{\text{Acryl}}\text{Act1}$ and $^{\text{Acryl}}\text{ActA}$ from CCT, showing native actin (N), EDTA unfolded actin (U), and the time in minutes after addition of 2 mM ATP and 10 mM MgCl_2 to CCT-bound actin. $^{\text{Acryl}}\text{ActA}_{\text{NAT}}$ is not released from CCT; (D) Refolded $^{488}\text{Act1}$ and $^{\text{Acryl}}\text{Act1}$ can be copolymerized with unlabeled Act1, as can be shown by centrifugation at 100 000 x g of these filaments. The Coomassie stained SDS gel is shown on the left hand side, followed by in-gel fluorescence of $^{488}\text{Act1}$ and $^{\text{Acryl}}\text{Act1}$. A small amount of aggregated CCT, formed overnight during the actin polymerisation step, pellets.

Fig. 4. (A) Representative scan of fluorescence emission of $^{\text{Acryl}}\text{Act1}$ at 470 nm following release from CCT with 2 mM ATP, 10 mM MgCl_2 , fitted to a single exponential to allow the rate constant k to be determined; (B) Arrhenius plot of the rate constants for release of $^{\text{Acryl}}\text{Act1}$ from CCT with 1 mM ATP, 5 mM MgCl_2 (circles), 2 mM ATP, 10 mM MgCl_2 (squares) or 4 mM ATP, 20 mM MgCl_2 (diamonds). Error bars show the standard deviation.

Fig. 5. Arrhenius plot of the rate constants for release of $^{\text{Acryl}}\text{Act1}$ from CCT4anc2 in the presence of 4 mM ATP (open circles) or 2 mM ATP (solid circles). Temperatures of 35°C and higher are non-permissive *in vivo* in CCT4anc2 (9). Error bars show the standard deviation.

Fig. 6. (A) Representative scans of the fluorescence emission at 470 nm for incubation of $^{\text{Acryl}}\text{Act1-CCT}$ with AMP-PNP followed by addition of ADP (triangles), and release of $^{\text{Acryl}}\text{Act1}$ from CCT by addition of ATP following pre-incubation in the presence (solid circles) or absence (open circles) of AMP-PNP; (B) In-gel fluorescence of a native gel showing release of $^{488}\text{Act1}$ from CCT, showing $^{488}\text{Act1}$ unfolded onto CCT (U), then incubation of this complex for 10 minutes with AMP-PNP and time points following addition of ATP to allow release of $^{488}\text{Act1}_{\text{NAT}}$; (C) quantification of the $^{488}\text{Act1}_{\text{NAT}}$ band from the gel (B).

Fig. 7. Schematic representation of the proposed model of $^{\text{Acryl}}\text{Act1}$ unfolding and binding to CCT. The fluorescent probe is in a more polar environment in native actin than in the EDTA-unfolded intermediate I_3 and in the complex with CCT. (A) Actin binds in a 1,4-conformation across the cavity of the open CCT complex (black) as shown by EM (17). (B) Binding of ATP or AMP-PNP closes the lid of CCT (gray). Step 1: Nucleotide binding induces a conformational change within the actin molecule (23), but does not facilitate release and the environment of the C-terminus remains unaffected. Step 2: Hydrolysis of ATP is required for the final packing of the actin C-terminus and release of native actin from CCT.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5






Figure 6

(A)



(B)



(C)




Figure 7

