

Research Article

# Effect of different $\gamma$ -subunit isoforms on the regulation of AMPK

Robin Willows<sup>1</sup>, Naveenan Navaratnam<sup>1</sup>, Ana Lima<sup>1</sup>, Jon Read<sup>2</sup> and David Carling<sup>1,3</sup>

<sup>1</sup>Cellular Stress Group, Medical Research Council London Institute of Medical Sciences, Hammersmith Hospital, Imperial College, London W12 0NN, U.K.; <sup>2</sup>AstraZeneca R&D, Discovery Sciences, Darwin Building, 310 Cambridge Science Park, Milton Road, Cambridge CB4 0WG, U.K.; and <sup>3</sup>Institute of Clinical Sciences, Faculty of Medicine, Imperial College London, Du Cane Road, London W12 0NN, U.K.

Correspondence: David Carling (david.carling@lms.mrc.ac.uk)



AMP-activated protein kinase (AMPK) plays a key role in integrating metabolic pathways in response to energy demand. AMPK activation results in a wide range of downstream responses, many of which are associated with improved metabolic outcome, making AMPK an attractive target for the treatment of metabolic diseases. AMPK is a heterotrimeric complex consisting of a catalytic subunit ( $\alpha$ ) and two regulatory subunits ( $\beta$  and  $\gamma$ ). The  $\gamma$ -subunit harbours the nucleotide-binding sites and plays an important role in AMPK regulation in response to cellular energy levels. In mammals, there are three isoforms of the  $\gamma$ -subunit and these respond differently to regulation by nucleotides, but there is limited information regarding their role in activation by small molecules. Here, we determined the effect of different  $\gamma$ -isoforms on AMPK by a direct activator, 991. In cells, 991 led to a greater activation of  $\gamma$ 2-containing AMPK complexes compared with either  $\gamma$ 1 or  $\gamma$ 3. This effect was dependent on the long N-terminal region of the  $\gamma$ 2-isoform. We were able to rule out an effect of Ser<sup>108</sup> phosphorylation, since mutation of Ser<sup>108</sup> to alanine in the  $\beta$ 2-isoform had no effect on activation of AMPK by 991 in either  $\gamma$ 1- or  $\gamma$ 2-complexes. The rate of dephosphorylation of Thr<sup>172</sup> was slower for  $\gamma$ 2- compared with  $\gamma$ 1-complexes, both in the absence and presence of 991. Our studies show that activation of AMPK by 991 depends on the nature of the  $\gamma$ -isoform. This finding may have implications for the design of isoform-selective AMPK activators.

## Introduction

AMP-activated protein kinase (AMPK) is a highly conserved heterotrimeric serine/threonine kinase widely characterised as a sensor of cellular energetic stress [1,2]. In response to activation by a decrease in ATP levels, AMPK phosphorylates many downstream targets involved in a diverse range of metabolic pathways [1,3,4]. The overall effect of AMPK activation is to turn off anabolic, ATP-utilising pathways and turn on catabolic, ATP-generating pathways. For instance, AMPK phosphorylates and inactivates acetyl-CoA carboxylase (ACC), which leads to inhibition of fatty acid synthesis and/or activation of fatty acid oxidation [1]. AMPK is a heterotrimeric complex consisting of a catalytic  $\alpha$ -subunit and two regulatory subunits ( $\beta$  and  $\gamma$ ) [1,5]. In mammals, there are two  $\alpha$ -subunit isoforms, together with two  $\beta$ - and three  $\gamma$ -isoforms, leading to 12 different possible AMPK complexes. The  $\gamma$ -subunit possesses four cystathionine- $\beta$ -synthase (CBS) domains, and these are involved in binding adenine nucleotides [5–7]. AMP binding to AMPK results in allosteric activation, which when assayed in the presence of physiological concentrations of ATP (5 mM) can reach up to 10-fold [8]. AMPK is activated 100–1000-fold following phosphorylation of the activation loop residue, Thr<sup>172</sup> (threonine 172 in the rat  $\alpha$ -subunit), by upstream kinases, including LKB1 (liver kinase B1) and calcium/calmodulin-dependent protein kinase kinase- $\beta$  [9–14]. A number of studies have reported that nucleotides regulate phosphorylation of AMPK by the upstream kinases, although it is not clear whether both AMP and ADP, or AMP alone, mediate this effect [8,15,16]. More

Received: 12 January 2017  
Revised: 15 March 2017  
Accepted: 16 March 2017

Accepted Manuscript online:  
16 March 2017  
Version of Record published:  
9 May 2017

recently, AMP has been shown to promote binding of AMPK and LKB1 to axin, providing a potential mechanism for increased AMPK phosphorylation by AMP [17]. In addition to these effects, both AMP and ADP protect against dephosphorylation of Thr<sup>172</sup> [7,18–20]. Two recent studies reported that the effects of nucleotides on AMPK activity depend on the isoform composition of the AMPK complexes [8,21].

Given its central role in regulating energy homeostasis, systemic activation of AMPK provides an attractive pharmacological target for combatting metabolic diseases, and many AMPK activators have been reported [22–24]. Most of the small-molecule AMPK activators that have been identified to date, such as A769662 and 991, bind to a single site formed between the  $\alpha$ - and  $\beta$ -subunits [24,25]. These direct activators protect against Thr<sup>172</sup> dephosphorylation and allosterically activate AMPK, although the mechanism by which they activate AMPK is distinct from the nucleotide-mediated effects on AMPK [24,25]. Similar to AMP, allosteric activation by synthetic small molecules depends on the isoform composition of the AMPK complex [21,24]. In addition, binding is significantly tighter with  $\beta$ 1- versus  $\beta$ 2-containing AMPK complexes, and recently, a highly selective  $\beta$ 1-activator was reported [22]. An important consideration in the development of AMPK activators as therapeutic drugs is the potential for deleterious effects of chronic AMPK activation *in vivo*. This is a particular concern with AMPK since dominantly inherited gain-of-function mutations in the  $\gamma$ 2-subunit of AMPK have been identified in humans [26]. These mutations lead to a cluster of cardiac abnormalities, including Wolff–Parkinson–White syndrome and cardiac hypertrophy associated with glycogen accumulation [27–29]. A recent study has also reported that humans carrying the R302Q mutation in  $\gamma$ 2 have increased adiposity and slightly raised fasting glucose levels compared with unaffected individuals [30]. This raises a concern for potential detrimental effects of chronic, systemic AMPK activation, although to date there have been no published studies for long-term *in vivo* effects of specific AMPK activators. Determining the effects of pharmacological activators on different AMPK complexes, and understanding whether there are any differences in their mechanism of activation, especially for  $\gamma$ 2-containing AMPK complexes, is an important factor in the design and development of potential drugs. Currently, there is little information regarding the effect of AMPK activators on  $\gamma$ 2-complexes, and in the recent work by Rajamohan et al. [21],  $\gamma$ 2-containing AMPK complexes were not included in the study.

In this study, we investigate the response of different  $\gamma$ -isoform AMPK complexes to pharmacological stimulation by 991, a potent direct activator of AMPK [24]. We show that in cell-based assays,  $\gamma$ 2-complexes are activated by 991 to a greater extent than either  $\gamma$ 1- or  $\gamma$ 3-complexes. This effect is mediated by the N-terminal region of  $\gamma$ 2 and is probably due to enhanced protection of Thr<sup>172</sup> from dephosphorylation. These findings suggest a role for the long N-terminal region of  $\gamma$ 2 in regulating AMPK activity that has not previously been recognised. Our results may also have important implications for the design and development of AMPK activators aimed at therapeutic intervention in human disease states.

## Experimental

### Materials and proteins

991 was synthesised as described previously [24]. PF06685249 [31] was a generous gift from Dr Russell Miller (Pfizer Global Research and Development). Recombinant PP2C $\alpha$  was purified as described previously [32].

### Cell culture

HEK293T cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM; Thermo) supplemented with 10% foetal bovine serum (Sigma–Aldrich). AMPK $\alpha$ ,  $\beta$  and  $\gamma$  constructs were cloned into pcDNA3 (Invitrogen) for transient expression, or pLPC (Addgene) with a puromycin selection cassette for stable transfection. For transfection experiments, cells were plated on 6 cm plates at 75% density and transfected using polyethylenimine (PEI; Polysciences) reagent at 2  $\mu$ g per construct with a PEI:DNA ratio of 3:1 (w/w). Thirty-six hours post-transfection, cells were transferred to serum-free DMEM for 2 h prior to treatment with 991 or PF249 (for 30 min). As a vehicle control, DMSO was added at the same concentration as for the 991-treated cells. Cells were washed three times with ice-cold PBS before the addition of lysis buffer: 50 mM HEPES (pH 7.4), 50 mM sodium fluoride, 5 mM sodium pyrophosphate, 1 mM ethylenediaminetetraacetic acid, 10% (v/v) glycerol, 1% (v/v) Triton X-100, 1 mM dithiothreitol, 0.1 mM phenylmethylsulfonyl fluoride, 4  $\mu$ g/ml trypsin inhibitor and 0.1 mM benzamidine. For stable transfections, subunits were cloned into pLPC vector and selected 48 h post-transfection with 3  $\mu$ g/ml puromycin (Thermo). Cell lines were maintained in puromycin and plated at 80% confluency the day before serum starvation and treatment as described above.

## CRISPR-mediated deletion of $\beta$ -subunit isoforms

HEK293T cells were transfected with plasmids containing Cas9 linked to GFP expression via a self-cleaving peptide and guide sequences targeting the first exon of  $\beta 1$  (GCTGGTATTGCCCATGATGG, GCCACCA TGCCGCTCCAGCG, GGGCTGTCCATCAGGATCTT, TTCCTCGGAGTGGAAGAGGT, TCAAGGTGCGA GCGGTGTGG) or  $\beta 2$  (CCCATGGCTGCAGCTCGTCG, ACCACCAGCGACCGGGTGTC, AGCGTGCAGCC TTGGCGCCG, ATGATCTTGTGCTCCTTCCC, CAGGGAGGCTGAACACGCTG) (Horizon Discovery, Cambridge, U.K.). Twenty-four hours post-transfection, cells were sorted based on GFP expression and individual colonies were analysed by western blotting.

## Western blotting

The protein concentration of cell lysates was determined by protein assay (Bio-Rad) prior to heating for 5 min at 95°C in 5 $\times$  SDS loading buffer. Proteins (15  $\mu$ g total) were resolved by SDS-PAGE on 10% polyacrylamide gels (National Diagnostics) and transferred to Immobilon-FL (Millipore) membranes at 4°C. Membranes were probed with primary antibodies at a 1:1000 dilution and incubated overnight at 4°C. The following antibodies were from Cell Signalling: mouse anti- $\alpha 1/2$  (#2793), rabbit anti- $\beta 1/2$  (#4150), rabbit anti- $\gamma 1$  (#4187), rabbit anti-His (#2365) and rabbit anti-pThr<sup>172</sup> (#2535). Rabbit anti-GAPDH was from Abcam (9485); mouse anti- $\alpha$ -tubulin (Sigma-Aldrich, V9131); mouse anti-FLAG M2 (Stratagene, 200472). After extensive washing, membranes were incubated for 30–60 min with LI-COR secondary antibodies diluted 1:10 000. Blots were imaged on an LI-COR Odyssey CLX. Peak area ratios for pThr<sup>172</sup> to total  $\alpha 1/2$  were obtained using Image Studio (LI-COR) to determine fold activation profiles. For capillary western blotting, cell lysates were diluted in HEPES lysis buffer to 0.6 mg/ml for experiments conducted on endogenous protein, or 0.1 mg/ml for experiments using transfected cells. Samples were prepared and analysed according to the manufacturer's instructions (ProteinSimple). Rabbit anti- $\beta 1/2$ , anti-pThr<sup>172</sup> and anti-pACC (Cell Signalling, 3661) were all used at 1:200 dilution in antibody dilution buffer (ProteinSimple).

## Immunoprecipitation of AMPK complexes

To obtain purified complexes for *in vitro* allosteric and protection assays,  $\beta 1\beta 2$  double knockout (dKO) cells were transfected as described above and treated with 5  $\mu$ M 991 (30 min) prior to lysis. FLAG-tagged complexes were immunoprecipitated by incubating lysate supernatant with FLAG M2 affinity gel (Sigma-Aldrich, A2220) overnight with gentle agitation. Following extensive washing with HEPES lysis buffer containing 300 mM NaCl, the resin was washed twice with 50 mM HEPES buffer (pH 7.4). Complexes were eluted with 2  $\times$  50  $\mu$ l of FLAG peptide (Sigma-Aldrich, F3290; 5  $\mu$ g/ml) in this buffer.

## AMPK assay

AMPK was assayed using the SAMS peptide assay as described previously [33]. Briefly, purified FLAG-tagged AMPK was incubated with 0.2 mM SAMS peptide, in the absence or presence of varying concentrations of 991 [in all cases, the final concentration of DMSO was kept at 0.5% (v/v)]. Incorporation of radiolabelled phosphate from ATP into SAMS peptide was measured by liquid scintillation counting. Isolated complexes were titrated to obtain a similar basal AMPK activity (measured in the absence of 991).

## Protection against dephosphorylation

Purified AMPK complexes were dephosphorylated by incubating at 37°C with 2.5  $\mu$ g/ml PP2C and 2.5 mM MgCl<sub>2</sub> in the absence or presence of 100 nM 991. Reactions were stopped by dilution (1:10) into ice-cold 50 mM HEPES buffer (pH 7.4), containing 50 mM NaF and 5 mM sodium pyrophosphate. AMPK activity was determined using the SAMS assay.

## Statistical analysis

Where appropriate, results were analysed using either Student's *t*-test or ANOVA using Bonferroni correction.

## Results

### Increased 991-stimulated Thr<sup>172</sup> phosphorylation in AMPK $\gamma$ 2 complexes compared with AMPK $\gamma$ 1


Two recent studies examined the role of AMPK subunit isoforms in regulation by nucleotides and pharmacological activators [8,21]. Despite these studies, however, the regulation of  $\gamma$ 2-containing AMPK complexes by pharmacological activators was not investigated. AMPK $\gamma$ 2 is widely expressed in mammalian tissues and mutations in *PRKAG2*, the gene encoding  $\gamma$ 2, cause a cluster of severe cardiac abnormalities in humans [27–29]. The lack of information regarding the regulation of AMPK $\gamma$ 2 complexes by pharmacological activators prompted us to investigate this further. We have been unable to express recombinant AMPK $\gamma$ 2 complexes in *Escherichia coli*, and hence we used mammalian expression for our study. We co-expressed  $\gamma$ 1 or  $\gamma$ 2 (isoform  $\gamma$ 2a, corresponding to the long form of  $\gamma$ 2 [34]) with either  $\alpha$ 1/ $\alpha$ 2 or  $\beta$ 1/ $\beta$ 2 in HEK293T cells and measured Thr<sup>172</sup> phosphorylation of the different AMPK complexes in response to activation by 991 (1  $\mu$ M), a potent pharmacological AMPK activator [24]. In all cases, there was a marked increase in 991-stimulated Thr<sup>172</sup> phosphorylation with the  $\gamma$ 2-containing AMPK complexes compared with the corresponding  $\gamma$ 1-containing complexes (Figure 1A,B). We carried out a similar experiment, but using varying concentrations of 991 (Figure 1C, D). In this experiment, we restricted our analysis to complexes containing the  $\beta$ 2-isoform, as this isoform is the predominant  $\beta$ -subunit isoform expressed in the heart [35]. Consistent with our initial experiment, 991-stimulated Thr<sup>172</sup> phosphorylation was greater in complexes containing the  $\gamma$ 2-isoform relative to the  $\gamma$ 1-isoform. There was also a trend for increased Thr<sup>172</sup> phosphorylation in the  $\gamma$ 2- versus  $\gamma$ 1-complexes in the absence of 991, although this was not statistically significant. The increase in 991-induced Thr<sup>172</sup> phosphorylation was similar for both  $\alpha$ 1 and  $\alpha$ 2 AMPK complexes, and since  $\alpha$ 2 is the predominant  $\alpha$ -isoform expressed in heart [36], we focussed on  $\alpha$ 2 $\beta$ 2 $\gamma$ 1- and  $\alpha$ 2 $\beta$ 2 $\gamma$ 2-complexes for our subsequent studies.

### CRISPR-generated $\beta$ 1/2 knockout HEK293T cell line

A potential complication with studying isoform-specific roles of AMPK subunits using mammalian expression systems is that the endogenous AMPK complexes could interfere with subsequent analyses. To circumvent this potential issue, we generated HEK293T cells lacking functional AMPK complexes. We used the CRISPR–Cas9 system to knockout expression of AMPK $\beta$ 1 and  $\beta$ 2. We chose to delete the  $\beta$ -subunits as these isoforms have been shown previously to play an important role in compound binding at the kinase domain–carbohydrate binding module (CBM) interface. In addition, there is a highly specific and sensitive antibody raised to a sequence that is conserved in both  $\beta$ 1 and  $\beta$ 2 that allows for detection of these isoforms (although the predicted masses of  $\beta$ 1 and  $\beta$ 2 are very similar, they are readily resolved by SDS–PAGE [35]). Cells were first transfected with guides targeting the first exon of  $\beta$ 1 and clones lacking expression of  $\beta$ 1 (as assessed by western blotting) were isolated. Clones lacking detectable  $\beta$ 1 expression were subsequently transfected with guides targeting the first exon of  $\beta$ 2, and cells lacking detectable  $\beta$ 2 expression were identified by blotting. In most cases, we observed clones that, although lacking expression of full-length  $\beta$ 1 and  $\beta$ 2, appeared to express truncated forms of the  $\beta$ -subunit that cross-reacted with the anti- $\beta$  antibody (Figure 2A). However, in some clones, there was no detectable cross-reactivity (e.g. clone 5 in Figure 2A). Clones lacking detectable  $\beta$ -subunit expression, which we term  $\beta$ 1 $\beta$ 2 dKO cells, were expanded and used for subsequent studies of AMPK regulation.

Total AMPK $\alpha$  and AMPK $\gamma$ 1 expression were barely detectable in the  $\beta$ 1 $\beta$ 2 dKO cells (Figure 2B,C). We have been unable to detect endogenous  $\gamma$ 2 or  $\gamma$ 3 expression in wild-type HEK293T cells using either commercial or in-house antibodies and hence cannot determine the effect of deletion of the  $\beta$ -subunit on their expression. However, a previous study reported that  $\gamma$ 1 is the predominant  $\gamma$ -isoform in HEK293 cells [37]. Moreover, phosphorylation of Thr<sup>172</sup> and ACC, a widely used downstream substrate for AMPK, were barely detectable in the  $\beta$ 1 $\beta$ 2 dKO cells, either in the absence or presence of 991 (Figure 2C). In contrast, phosphorylation of both Thr<sup>172</sup> and ACC was readily detected in wild-type cells and markedly increased following treatment with 991. Taken together, these findings demonstrate that CRISPR/Cas9-mediated deletion of  $\beta$ 1 and  $\beta$ 2 in HEK293T cells generates a cell line almost completely devoid of functional AMPKs.


Next, we generated cell lines in which we stably reintroduced either  $\beta$ 1 or  $\beta$ 2 into the  $\beta$ 1 $\beta$ 2 dKO cells. This provided us with cells with defined  $\beta$ -isoform expression, allowing us to investigate  $\beta$ -isoform-specific regulation of AMPK. Cells were treated in the absence or presence of increasing concentrations of two different AMPK activators, 991 or PF06685249 (hereafter referred to as PF249), a  $\beta$ 1-selective AMPK activator [31]. In both  $\beta$ 1- and  $\beta$ 2-expressing cells, 991 caused a dose-dependent increase in Thr<sup>172</sup> and ACC phosphorylation



**Figure 1. Activation of  $\gamma 1$  versus  $\gamma 2$  AMPK complexes by 991.**

HEK293T cells were co-transfected with cDNA encoding either  $\gamma 1$  or  $\gamma 2$  together with  $\alpha 1$  or  $\alpha 2$  and  $\beta 1$  or  $\beta 2$ . Thirty-six hours post-transfection, cells were serum-starved for 2 h and incubated with DMSO or 991 (1  $\mu\text{M}$ ) for 30 min prior to cell lysis. (A) Representative blots showing Thr<sup>172</sup> phosphorylation and  $\beta 1/\beta 2$  expression, determined by western blotting using an automated capillary-based electrophoresis system (Simon, Protein Simple), are shown. (B) Graphs showing quantification of Thr<sup>172</sup> phosphorylation relative to  $\beta$ -subunit expression. (C) Thr<sup>172</sup> phosphorylation and  $\beta 2$  expression were determined in response to varying concentrations of 991. (D) Fold change in Thr<sup>172</sup> phosphorylation:  $\beta$ -subunit expression relative to the corresponding  $\gamma 1$ -complex in the absence of 991 is shown. Results shown for (B) and (D) are mean  $\pm$  SEM for two independent experiments.

(Figure 3A). We previously reported that  $\beta 1$ -complexes bound 991 with  $\sim 10$ -fold higher affinity compared with  $\beta 2$ -complexes [24]. Consistent with this, we observed that phosphorylation of ACC is more sensitive to 991 in  $\beta 1$ -expressing cells relative to  $\beta 2$ -expressing cells. It is also clear that the 991-induced increase in ACC phosphorylation is markedly more pronounced than that in Thr<sup>172</sup> phosphorylation, particularly at low concentrations. This presumably reflects the fact that, in addition to increasing Thr<sup>172</sup> phosphorylation, 991 causes a significant allosteric activation of AMPK, which would result in more marked phosphorylation of downstream targets. Treatment with PF249, which is more than 1000-fold more potent towards  $\beta 1$ - compared with  $\beta 2$ -complexes [31], led to a robust increase in ACC phosphorylation, together with a weaker, but detectable, increase in Thr<sup>172</sup> phosphorylation in  $\beta 1$ -expressing cells. Importantly, however, PF249 had virtually no effect on Thr<sup>172</sup> or ACC phosphorylation in  $\beta 2$ -expressing cells (Figure 3B). There was a very slight increase in ACC phosphorylation in the  $\beta 2$ -expressing cells at the highest concentration of 991 (5  $\mu\text{M}$ ), but this was less than the phosphorylation of ACC observed at the lowest concentration (0.1  $\mu\text{M}$ ) in the  $\beta 1$ -expressing cells. These




**Figure 2. Generation of  $\beta 1\beta 2$  dKO cells.**

Clones of HEK293T cells were isolated by flow cytometry following transfection with plasmids containing Cas9 and guide RNAs targeting the  $\beta 1$ - and  $\beta 2$ -isoforms of AMPK. **(A)** A representative capillary western blot of  $\beta$  subunit expression in five different clones compared with wild-type HEK293T is shown. Lower molecular mass cross-reacting bands, potentially due to truncated  $\beta$ -subunit protein, are indicated. **(B)** Total AMPK $\alpha$  and **(C)** AMPK $\gamma 1$  expression in  $\beta 1\beta 2$  dKO cells relative to wild-type cells were determined by western blotting. **(D)** Wild-type and  $\beta 1\beta 2$  dKO cells were incubated with DMSO or 991 (1  $\mu$ M) for 30 min. Levels of ACC and Thr<sup>172</sup> phosphorylation, together with  $\beta$ -subunit expression, were determined. In each case, a representative blot with two independent samples is shown.

results confirm that AMPK signalling is restored by re-expression of the  $\beta$ -subunit in the  $\beta 1\beta 2$  dKO cells, and demonstrate the  $\beta$ -isoform selectivity of AMPK activators in human cells.

## $\gamma 2$ N-terminal region mediates the increased 991-induced Thr<sup>172</sup> phosphorylation

The  $\gamma$ -isoforms all contain a highly conserved C-terminal region harbouring the four CBS domains involved in nucleotide binding. In addition, the  $\gamma 2$ - and  $\gamma 3$ -isoforms contain long N-terminal extensions that are not present in the  $\gamma 1$ -isoform [38]. These N-terminal extensions show no obvious sequence conservation between isoforms, or significant homology with other proteins. In addition, variant forms of  $\gamma 2$ - and  $\gamma 3$ -transcripts have been identified that encode predicted proteins with truncated N-terminal regions [34,39]. At present, it is not known whether the N-terminal extensions of  $\gamma 2$  or  $\gamma 3$  play any specific functional role, or what role, if any, the different variants play *in vivo*. To investigate further the effect of the different  $\gamma$ -isoforms on AMPK regulation, we expressed  $\alpha 2\beta 2$  with either  $\gamma 1$ ,  $\gamma 2$  or  $\gamma 3$  in the  $\beta 1\beta 2$  dKO cells and compared Thr<sup>172</sup> in the different AMPK complexes in response to 991. As shown in Figure 4, 991 stimulated Thr<sup>172</sup> phosphorylation of all AMPK complexes. However, the increase in Thr<sup>172</sup> phosphorylation within the  $\gamma 2$ -complex was significantly greater than that in either the  $\gamma 1$ - or  $\gamma 3$ -complexes. To determine whether the enhanced response of  $\gamma 2$ -containing complexes to 991 was mediated by its N-terminal region, we synthesised constructs in which we replaced the N-terminal region of  $\gamma 1$  (corresponding to the first 31 amino acids of  $\gamma 1$ ) with sequences from  $\gamma 2$  (see Figure 5A). In one chimeric construct (C1), we replaced the  $\gamma 1$  sequence with the N-terminal sequence of the short form of  $\gamma 2$  (termed  $\gamma 2b$ ), and in a second construct (C2), we replaced it with the N-terminal sequence of the long form of  $\gamma 2$  (termed  $\gamma 2a$ ). It is important to note that despite numerous attempts, and using a variety of epitope tags at both the N- and C-termini, we have been unable to express the short form of  $\gamma 2$  (isoform  $\gamma 2b$ ) in mammalian cells. In addition, we were unable to express a chimeric construct in which we replaced the N-terminal sequence of  $\gamma 2a$  with the N-terminal region of  $\gamma 1$  (data not shown). We co-expressed the two chimeric  $\gamma$ -constructs with  $\alpha 2$  and  $\beta 2$  and confirmed their expression by western blotting with an anti-His antibody directed against a C-terminal hexa-histidine tag included in the constructs (Figure 5B). Thr<sup>172</sup>




**Figure 3. Activation of  $\beta$ 1- and  $\beta$ 2-AMPK complexes by direct small-molecule activators.**

AMPK $\beta$ 1 or AMPK $\beta$ 2 expressing stable cell lines were generated in the  $\beta$ 1 $\beta$ 2 dKO HEK293T cells. Cells were incubated with different concentrations of 991 (A) or PF249 (B). Levels of ACC and Thr<sup>172</sup> phosphorylation, together with  $\beta$ -subunit expression, were determined.

phosphorylation was markedly enhanced in the AMPK complex containing chimera 2 (which has the entire N-terminal region of  $\gamma$ 2a replacing the N-terminal region of  $\gamma$ 1) and was similar to Thr<sup>172</sup> phosphorylation in AMPK complexes containing native  $\gamma$ 2 (Figure 5C,D). In contrast, Thr<sup>172</sup> phosphorylation of AMPK containing chimera 1 (replacing the N-terminal region of  $\gamma$ 1 with the short N-terminal region of  $\gamma$ 2b) was the same as in AMPK containing native  $\gamma$ 1. These data strongly suggest that the enhanced response of  $\gamma$ 2-AMPK to 991 is mediated by the long N-terminal region of  $\gamma$ 2.

### Phosphorylation of Ser<sup>108</sup> in $\beta$ 2

Previous studies have shown that phosphorylation of serine 108 (Ser<sup>108</sup>) in the  $\beta$ 1-subunit significantly increases the binding affinity of AMPK to both 991 and A769662 [24], leading us to speculate whether Ser<sup>108</sup> phosphorylation might play a role in the enhanced activation of  $\gamma$ 2-containing AMPK complexes by 991. Moreover, since our experimental system focussed mainly on  $\beta$ 2-containing AMPK complexes, this also provided us with an opportunity to examine the role of Ser<sup>108</sup> phosphorylation in  $\beta$ 2, which has not been reported previously. We expressed  $\beta$ 2 harbouring mutation of Ser<sup>108</sup> to a non-phosphorylatable alanine  $\beta$ 2<sub>(S108A)</sub> with either  $\alpha$ 2 $\gamma$ 1 or  $\alpha$ 2 $\gamma$ 2 in  $\beta$ 1 $\beta$ 2 dKO cells. As shown in Figure 6, Thr<sup>172</sup> phosphorylation in response to treatment with 991 was similar in AMPK complexes containing either wild-type  $\beta$ 2 or  $\beta$ 2<sub>(S108A)</sub>. This contrasts with the effect of S108A mutation in  $\beta$ 1, where activation by 991 is significantly reduced in AMPK complexes harbouring  $\beta$ 1<sub>(S108A)</sub> compared with AMPK containing wild-type  $\beta$ 1 [24]. The lack of effect of the  $\beta$ 2<sub>(S108A)</sub> mutation on AMPK activation in cells is intriguing, as the region surrounding Ser<sup>108</sup> shows a high degree of amino acid sequence conservation between  $\beta$ 1 and  $\beta$ 2 [35]. We extended our study to determine the effect of the S108A mutation in  $\beta$ 2 on allosteric activation of AMPK by 991 in cell-free assays (Figure 7 and Table 1). AMPK complexes were expressed in HEK293T cells and isolated by affinity purification using FLAG resin. AMPK activity was determined in the absence or presence of varying concentrations of 991. Wild-type  $\alpha$ 2 $\beta$ 2 $\gamma$ 1- and



**Figure 4. Differential activation of AMPK $\gamma$  isoforms by 991.**

FLAG-tagged AMPK $\gamma$ 1,  $\gamma$ 2 and  $\gamma$ 3 constructs were transiently co-transfected with  $\alpha$ 2 and  $\beta$ 2 into  $\beta$ 1 $\beta$ 2 dKO cells. Cells were treated with or without 991 (1  $\mu$ M) for 30 min. (A) Thr<sup>172</sup> phosphorylation and total  $\alpha$  expression levels were determined by fluorescence-based western blotting. (B) Blots were quantified and the increase in Thr<sup>172</sup> phosphorylation in response to 991 relative to the DMSO control was determined as a fold change. Results are for three independent experiments, performed in duplicate, and are plotted as means  $\pm$  SEM. Statistically significant results, determined by one-way ANOVA, using Bonferroni correction, are indicated: \* $P$  < 0.05, \*\* $P$  < 0.01. There was no significant difference between  $\gamma$ 1- and  $\gamma$ 3-complexes. (C) Expression of the different  $\gamma$ -isoforms in the AMPK complex was confirmed by western blotting using an anti-FLAG antibody.

$\alpha 2\beta 2\gamma 2$ -complexes were allosterically activated  $\sim 5$ -fold and showed very similar dose–response curves with 991 (Figure 7A). There was no significant difference in the  $A_{0.5}$  for 991 between the  $\gamma 1$ - and  $\gamma 2$ -containing complexes (Table 1), indicating that 991 binds with similar affinity to both complexes. Consistent with the cell-based studies, there was no significant difference in allosteric activation of AMPK containing wild-type  $\beta 2$  versus  $\beta 2$  harbouring the S108A mutation (Figure 7B). In marked contrast, the S108A mutation in  $\beta 1$  caused a marked rightward shift in the dose–response curve for 991, similar to the effect we previously reported for AMPK $\beta 1$  complexes expressed in *E. coli* [24]. These findings suggest that, unlike  $\beta 1$ , phosphorylation of Ser<sup>108</sup> in  $\beta 2$  does not play a significant role in binding or activation of AMPK by 991. Moreover, in keeping with our previous study using bacterially expressed AMPK complexes, 991 was more potent and caused a greater level of allosteric activation (8-fold versus 5-fold), with mammalian expressed  $\beta 1$ -containing AMPK complexes compared with  $\beta 2$  complexes (Figure 7B and Table 1).


### Effect of the $\gamma$ -isoform on protection against Thr<sup>172</sup> dephosphorylation

The finding that the  $\gamma 1$  and  $\gamma 2$  AMPK complexes show similar allosteric activation by 991 indicates that there is no difference in binding between the complexes. This rules out the possibility that the increased 991-induced Thr<sup>172</sup> phosphorylation of the  $\gamma 2$ -complex is due to increased affinity for 991. We therefore decided to investigate whether the  $\gamma$ -isoforms might play a role in protection against Thr<sup>172</sup> dephosphorylation. Recombinant AMPK complexes were purified from HEK293T cells and the rate of dephosphorylation by PP2C was measured in the absence (Figure 8A) or presence (Figure 8B) of 991. Consistent with previous studies, 991 protected AMPK against Thr<sup>172</sup> dephosphorylation for both  $\gamma 1$ - and  $\gamma 2$ -complexes. Importantly, however, the rate of dephosphorylation of  $\gamma 2$ -containing complexes was slower than that of  $\gamma 1$ -complexes, both in the absence and presence of 991. At the 15 min time point, the percentage of AMPK activity remaining was  $24 \pm 5$  versus  $45 \pm 5\%$  (for the AMPK $\gamma 1$  complex in the absence or presence of 991, respectively;  $n = 5 \pm \text{SEM}$ ) and  $54 \pm 4$  versus  $70 \pm 3\%$  (for AMPK $\gamma 2$  complex in the absence or presence of 991, respectively;  $n = 5 \pm \text{SEM}$ ). These results demonstrate that  $\gamma 2$  is more resistant to dephosphorylation by PP2C in cell-free assays. This finding provides a potential mechanism for the increased Thr<sup>172</sup> phosphorylation in  $\gamma 2$ -complexes in response to 991 in cells. Furthermore, this finding may also account for the increased basal level of Thr<sup>172</sup> phosphorylation in  $\gamma 2$ -complexes relative to  $\gamma 1$ -complexes that was observed in HEK293T cells.

### Discussion

Our results show that AMPK complexes containing the  $\gamma 2$ -isoform have increased Thr<sup>172</sup> phosphorylation compared with either  $\gamma 1$ - or  $\gamma 3$ -complexes in response to 991 in human cells. This effect is mediated by the long N-terminal region present in  $\gamma 2$ , since a chimeric protein containing the C-terminal region of  $\gamma 1$  fused to the N-terminal region of  $\gamma 2$  mimicked the response of native  $\gamma 2$  to 991. Furthermore, we demonstrate that a potential mechanism for the enhanced phosphorylation of Thr<sup>172</sup> in  $\gamma 2$  AMPK is a decrease in the rate of dephosphorylation by protein phosphatases. This effect was seen in the absence and presence of 991, suggesting that it is an intrinsic feature of  $\gamma 2$ -containing complexes. This would explain the finding that the level of Thr<sup>172</sup> phosphorylation in  $\gamma 2$ -complexes is greater than that of  $\gamma 1$ - or  $\gamma 3$ -complexes in untreated cells. At present, it is not clear how the N-terminal region of  $\gamma 2$  could interfere with dephosphorylation of Thr<sup>172</sup>. The simplest explanation would be that the long N-terminal extension of  $\gamma 2$  sterically hinders access to Thr<sup>172</sup> by protein phosphatases. Another possibility would be that the N-terminal region of  $\gamma 2$  interacts with a distinct region of AMPK causing a conformational change within the AMPK complex which leads to increased protection against dephosphorylation. At present, however, there is no molecular structural information for AMPK  $\gamma 2$ -containing complexes to support either of these hypotheses. The N-terminal region of  $\gamma 2a$  (residues 1–241) shows no obvious sequence similarity to other proteins. Analysis using the PSIPRED web server [40], a secondary protein structure prediction package, reveals that residues 1–220 are predicted to have a high probability of being intrinsically disordered. If this were the case, it would raise many possibilities for how this region might play a role in regulating AMPK activity. For instance, the N-terminal region might bind to other proteins and/or undergo changes in post-translational modification, features commonly associated with intrinsically disordered proteins [41]. It is interesting to note that we have been unsuccessful in all our efforts to express AMPK complexes containing the long isoform of  $\gamma 2$  in bacteria. Furthermore, in a recent publication investigating different AMPK complexes expressed in bacteria, the  $\gamma 2$  complex was not studied [21], although no reason was stated for its omission. It is possible that stable expression of the AMPK complex containing the long isoform of  $\gamma 2$  requires another protein or cofactor, and/or post-translational modification, that is not present in bacterial cells. Further






**Figure 5. Increased activation by 991 is mediated by the N-terminal region of  $\gamma$ 2.**


(A) A schematic representation of the  $\gamma$ 1- and  $\gamma$ 2-isoforms is shown, together with the two chimeric  $\gamma$ -constructs [ $\gamma$ (C1) and  $\gamma$ (C2)] used in the present study. Different regions within the isoforms are shown colour-coded. The conserved core region, containing the four CBS repeats, is indicated in red for  $\gamma$ 1 and purple for  $\gamma$ 2. The N-terminal region of  $\gamma$ 1 is shown in green, the N-terminal sequence for  $\gamma$ 2b (the short isoform of  $\gamma$ 2) is shown in yellow and the long N-terminal extension in  $\gamma$ 2a (the long isoform of  $\gamma$ 2) is coloured in blue. All constructs have a C-terminal hexa-His tag shown in grey. (B) Expression of the two chimeric proteins was confirmed by western blotting using an anti-His antibody and is shown relative to  $\gamma$ 1-expression. Vinculin expression was used as a loading control. (C)  $\beta$ 1 $\beta$ 1 dKO cells were transfected with different AMPK complexes and treated with DMSO or 991 (1  $\mu$ M) for 30 min. Thr<sup>172</sup> phosphorylation and total  $\alpha$  expression levels were determined by fluorescence-based western blotting. Although the  $\alpha$ 2 $\beta$ 2 $\gamma$ 2-complex was analysed in the same experiment at the same time, it was performed on a separate blot and is therefore shown as a distinct panel. (D) Blots were quantified and the increase in Thr<sup>172</sup> phosphorylation in response to 991 relative to the DMSO control was determined as a fold change. Results are for three independent experiments, performed in duplicate, and are plotted as means  $\pm$  SEM. Statistically significant results, determined by one-way ANOVA, using Bonferroni correction, are indicated (\* $P$  < 0.05).

work is required to explore these possibilities. Similarly, we have been unable to express the short form of  $\gamma$ 2 ( $\gamma$ 2b), or a chimeric protein comprising the short N-terminal sequence present in  $\gamma$ 2b fused to the C-terminal core region of  $\gamma$ 1, in mammalian cells. It is possible, therefore, that the short N-terminal sequence in  $\gamma$ 2b destabilises the AMPK complex in mammalian cells. Two related studies reported protein expression of AMPK



**Figure 6. Mutation of Ser<sup>108</sup> in  $\beta$ 2 does not mediate activation by 991 in cells.**

$\beta$ 1 $\beta$ 2 dKO cells were transfected with  $\alpha$ 2 and either  $\gamma$ 1 or  $\gamma$ 2 with either wild-type  $\beta$ 2 or  $\beta$ 2 harbouring mutation of Ser<sup>108</sup> to alanine ( $\beta$ 2<sub>(S108A)</sub>). Cells were treated with DMSO or 991 (1  $\mu$ M) for 30 min and analysed for Thr<sup>172</sup> phosphorylation and total  $\alpha$  expression, and a representative western blot is shown in (A). (B) Blots were quantified and the increase in Thr<sup>172</sup> phosphorylation in response to 991 relative to the DMSO control was determined as a fold change. Results are for three independent experiments, performed in duplicate, and are plotted as means  $\pm$  SEM. Statistically significant results, determined by one-way ANOVA, using Bonferroni correction, are indicated (\*\*\* $P$  < 0.001). There was no statistically significant difference (NS) between wild-type  $\beta$ 2 and  $\beta$ 2<sub>(S108A)</sub> for either  $\gamma$ 1- or  $\gamma$ 2-complexes.



**Figure 7. Allosteric activation of AMPK $\gamma$ 1 and  $\gamma$ 2 complexes by 991.**

AMPK  $\alpha$ 2 $\beta$ 2 $\gamma$ 1- and  $\alpha$ 2 $\beta$ 2 $\gamma$ 2-complexes (harbouring a FLAG tag at the N-terminus of the  $\gamma$ -isoform) were expressed in  $\beta$ 1 $\beta$ 2 dKO cells and isolated by immunoprecipitation using FLAG affinity resin. After extensive washing, proteins were eluted using FLAG peptide. (A) AMPK activity was measured using the SAMS peptide assay in the presence of varying concentrations of 991 ( $n = 5$  independent experiments). (B) AMPK  $\alpha$ 2 $\beta$ 1 $\gamma$ 2,  $\alpha$ 2 $\beta$ 1<sub>(S108A)</sub> $\gamma$ 2,  $\alpha$ 2 $\beta$ 2 $\gamma$ 2 and  $\alpha$ 2 $\beta$ 2<sub>(S108A)</sub> $\gamma$ 2 were expressed in  $\beta$ 1 $\beta$ 2 dKO cells and isolated by immunoprecipitation using FLAG affinity resin (the  $\beta$ -constructs contain a C-terminal FLAG tag). AMPK activity for each of the complexes was measured using the SAMS peptide assay in the presence of varying concentrations of 991 ( $n = 3$  independent experiments). The data were fitted to the equation:  $v = b + \frac{[(s \times b) - b] \times [991]}{A_{0.5} + [991]}$ , where  $s$  is the relative stimulation,  $b$  is the basal activity and  $A_{0.5}$  is the concentration of 991 giving half-maximal stimulation, using the GraphPad Prism software. In all cases, results are plotted as the means  $\pm$  SEM and the theoretical curves were fitted using the curve-fitting programme.

complexes containing  $\gamma$ 2b in mouse heart [34,42], although the specificity of the anti-AMPK $\gamma$ 2 antibodies used in these studies was not demonstrated convincingly. More recently, evidence for the expression of the long form of  $\gamma$ 2 ( $\gamma$ 2a) in mouse liver and heart was reported, and in this case, the specificity of the antibody used was demonstrated using tissues from AMPK $\gamma$ 2 knockout mice [43]. Thus, while direct evidence for protein expression of  $\gamma$ 2b *in vivo* remains equivocal, it is clear that  $\gamma$ 2a protein is expressed *in vivo*.

A recent study investigated the effects of the different  $\gamma$  subunit isoforms on AMPK regulation by nucleotides [8]. In that study, similar to our current study, AMPK complexes were expressed in human cells. Our results complement and add to the previous study by investigating the effects of the different  $\gamma$ -isoforms on AMPK regulation by a pharmacological activator, 991. One potential problem with expressing AMPK in mammalian cells is the possibility that endogenous AMPK subunits might compromise interpretation of the results. To mitigate against this possibility, we generated HEK293T cells in which both the  $\beta$ 1- and  $\beta$ 2-isoforms were genetically deleted. We chose to delete the  $\beta$ -subunit isoforms as previous studies have shown that the  $\beta$ -subunit plays a fundamental role in binding small-molecule AMPK activators, and that there is a significant difference in binding between AMPK complexes containing either the  $\beta$ 1- or  $\beta$ 2-isoform [22,24]. The  $\beta$ 1 $\beta$ 2 dKO cells provide a human cell line that is readily amenable to transfection (transient or stable) that can be used to investigate the regulation of the  $\beta$ -isoforms, including mutant variants of these isoforms, by different compounds. In

**Table 1 Allosteric activation of AMPK complexes by 991**


AMPK complexes expressed in  $\beta 1\beta 2$  dKO cells were isolated by immunoprecipitation using FLAG affinity resin. After extensive washing, proteins were eluted using the FLAG peptide and AMPK activity was measured using the SAMS peptide assay in the presence of varying concentrations of 991. The data were fitted to the equation:  $v = b + \{[(s \times b) - b] \times [991]\} / (A_{0.5} + [991])$ , where  $s$  is the relative stimulation,  $b$  is the basal activity and  $A_{0.5}$  is the concentration of 991 giving half-maximal stimulation, using the GraphPad Prism software. In all cases, results are from at least three independent experiments and are plotted as the means  $\pm$  SEM.

Complex	$A_{0.5}$ 991 (nM)	Activation (fold)
$\alpha 2\beta 2\gamma 1$	147 $\pm$ 28	5.7 $\pm$ 0.3
$\alpha 2\beta 2\gamma 2$	97 $\pm$ 18	5.0 $\pm$ 0.2
$\alpha 2\beta 2_{(S108A)}\gamma 2$	170 $\pm$ 35	5.4 $\pm$ 0.3
$\alpha 2\beta 1\gamma 2$	12.7 $\pm$ 4	7.7 $\pm$ 0.4
$\alpha 2\beta 1_{(S108A)}\gamma 2$	462 $\pm$ 142	5.2 $\pm$ 0.5

addition, it is possible that a natural ligand exists that binds to the same site as the pharmacological activators, although no such ligand has been identified to date. The availability of cell models expressing different forms of the  $\beta$ -subunit, e.g. full-length protein versus a C-terminal fragment lacking the CBM required for compound binding, will also provide useful tools in searching for endogenous AMPK activators.

We show that total AMPK $\alpha$  expression,  $\gamma 1$ , pThr172 and pACC were almost completely undetectable in the  $\beta 1\beta 2$  dKO cells. A caveat to the present work, however, is that we have been unable to determine directly the effect of  $\beta$ -subunit deletion on endogenous  $\gamma 2$ - and  $\gamma 3$ -expression. This may reflect the finding that  $\gamma 1$  is the predominant  $\gamma$ -isoform expressed in HEK293 cells [37], and/or could be due to the lack of availability of sensitive antibodies against these isoforms. Nonetheless, the almost complete loss of functional AMPK provides an excellent background in which to express AMPK complexes of defined isoform composition. As part of our validation of the  $\beta 1\beta 2$  dKO cells, we stably reintroduced either  $\beta 1$ - or  $\beta 2$ -isoforms in order to generate cells expressing exclusively either  $\beta 1$  or  $\beta 2$  and treated them with two different pharmacological AMPK activators, with different  $\beta$ -isoform selectivity. The effect of the activators on phosphorylation of Thr<sup>172</sup> and ACC correlated extremely well with their  $\beta$ -isoform selectivity determined using purified AMPK complexes. These cells will provide a useful resource for future studies aimed at determining  $\beta$ -isoform selectivity of different AMPK activators in a cellular context.

An important finding that emerged from our study is that Ser<sup>108</sup> in  $\beta 2$  does not appear to play a role in AMPK activation by 991. This is in contrast with  $\beta 1$ , where phosphorylation of Ser<sup>108</sup> increases 991 binding by 50–100-fold, depending on the nature of the  $\alpha$ -isoform present in the complex [24]. In  $\beta 1$ , pSer<sup>108</sup> forms many electrostatic interactions with residues from the  $\alpha$ -subunit as well as Asn<sup>111</sup> in  $\beta 1$  [24]. The sequence



**Figure 8. Protection against dephosphorylation of AMPK $\gamma 1$  and  $\gamma 2$  complexes by 991.**

AMPK  $\alpha 2\beta 2\gamma 1$ - and  $\alpha 2\beta 2\gamma 2$ -complexes (harbouring a FLAG tag at the N-terminus of the  $\gamma$ -isoform) were expressed in  $\beta 1\beta 2$  dKO cells and purified using FLAG affinity resin. AMPK complexes were incubated with recombinant PP2C $\alpha$  in the absence (A) or presence of 100 nM 991 (B). At the indicated times, aliquots were removed and diluted in buffer containing protein phosphatase inhibitors, and AMPK activity was measured using the SAMS peptide assay. The amount of AMPK was adjusted so that an equal amount of AMPK activity was added at the start of the incubation ( $T = 0$ ). Results are plotted as the percentage of activity remaining relative to the starting activity at time 0. In both cases, results are from five independent experiments and are the means  $\pm$  SEM.

surrounding Ser<sup>108</sup> is well conserved between  $\beta 1$  and  $\beta 2$ , although in  $\beta 2$  residue 111 is an aspartate, rather than an asparagine present in  $\beta 1$  [35]. To date, however, there is no high-resolution structural information for AMPK $\beta 2$  complexes bound to 991 (or other related pharmacological activators), and hence the molecular basis for the difference in requirement for Ser<sup>108</sup> phosphorylation between the  $\beta$ -isoforms is not clear. In  $\beta 1$ , Ser<sup>108</sup> is capable of undergoing autophosphorylation [44], although it is not known whether there are other kinases capable of phosphorylating this site. As far as we are aware, there are no published data on Ser<sup>108</sup> phosphorylation of  $\beta 2$ , and hence further studies will be required to determine whether Ser<sup>108</sup> phosphorylation in  $\beta 2$  plays a role in the regulation of AMPK. Although the physiological significance of Ser<sup>108</sup> phosphorylation for either  $\beta 1$  or  $\beta 2$  remains enigmatic, the finding that it is not required for 991-induced activation of  $\beta 2$ -complexes may prove significant for future studies exploring pharmacological regulation of AMPK.

Pharmacological activation of AMPK has attracted significant attention as a potential target for the treatment of metabolic diseases. Many potent AMPK activators have recently been identified, although information regarding their effects in *in vivo* models is currently lacking. Understanding whether there are any substantive differences in pharmacological activation of different AMPK complexes would provide useful complementary information for designing and interpreting subsequent *in vivo* studies. A recent *in vitro* study investigated the effect of A769662, the first direct synthetic AMPK activator to be identified [45], on different AMPK complexes, but this did not include  $\gamma 2$ -containing complexes [21]. In that recent study, recombinant AMPK complexes purified from bacteria were used, and no cell-based studies were included. In our current study, we used human cells for expressing AMPK complexes and an important consequence of this approach is that we were able to express complexes containing all three  $\gamma$ -isoforms. Another potentially important advantage gained by using AMPK complexes expressed in human cells is that they are likely to undergo the same post-translational modifications that occur *in vivo* and would be expected to recapitulate more closely the native enzyme. One of the key reasons for carrying out the current study was to investigate whether the different  $\gamma$ -isoforms has an impact on pharmacological activation of AMPK. In addition to demonstrating that 991-induced Thr<sup>172</sup> phosphorylation is greatest in  $\gamma 2$ -containing complexes, we also found that the basal phosphorylation of AMPK $\gamma 2$  is increased relative to either  $\gamma 1$  or  $\gamma 3$ . Despite significant interest in AMPKs, the physiological relevance of the different AMPK $\gamma$  isoforms is poorly understood. There is evidence for differences in tissue expression of the different isoforms. For example, expression of the  $\gamma 3$ -isoform is restricted to skeletal muscle [36,39], which might suggest tissue-specific roles for certain AMPK complexes. However, direct experimental evidence to support this hypothesis is lacking. Does our finding that  $\gamma 2$ -containing AMPK complexes have higher basal activity than the other  $\gamma$ -isoforms have any physiological significance? One possibility is that in human heart, where  $\gamma 2$ -expression is highest [36], the increased basal activity of AMPK $\gamma 2$  complexes is required to maintain cardiac-specific functions. In this respect, it is interesting that a previous study reported that ADP was as effective as AMP in protecting  $\gamma 2$ -complexes against dephosphorylation, whereas AMP was much more effective compared with ADP in protecting either  $\gamma 1$ - or  $\gamma 3$ -complexes [8]. This differential regulation could perhaps be important during periods of altered cardiac energetics which result in changes in adenine nucleotide levels.

The finding that 991-induced Thr<sup>172</sup> phosphorylation is greatest in  $\gamma 2$ -containing complexes may have implications for the therapeutic potential of long-term pharmacological activation of AMPK. Gain-of-function mutations in  $\gamma 2$  lead to a cluster of cardiac abnormalities in humans. Clearly, a major concern in targeting AMPK for activation is the possibility of developing phenotypes that overlap with those resulting from the  $\gamma 2$  mutations. In this regard, the increased sensitivity of  $\gamma 2$ -containing complexes could add to this potential risk. Many strategies could be envisaged to mitigate against this potential adverse effect. For instance, the beneficial effects of AMPK activation on metabolism are likely to be mediated by non-cardiac tissue. It may be possible to design compounds with tissue-specific delivery, and hence prevent accumulation in the heart. A greater understanding of the isoform composition of AMPK complexes in human heart might also provide nuanced strategies for pharmacological targeting. If, for example,  $\gamma 2$  is largely associated with  $\beta 2$  in the heart, development of  $\beta 1$ -selective activators would greatly reduce activation of the  $\gamma 2$ -complexes. The identification of highly selective  $\beta 1$ -activators [22] is an encouraging step in this approach.

### Abbreviations

ACC, acetyl-CoA carboxylase; AMPK, AMP-activated protein kinase; ANOVA, analysis of variance; CBM, carbohydrate binding module; CBS, cystathionine- $\beta$ -synthase; CRISPR, clustered regularly interspaced short palindromic repeats; dKO, double knockout; DMEM, Dulbecco's modified Eagle's medium; DMSO, dimethyl sulphoxide; HEPES, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid; LKB1, liver kinase B1; PAGE,

polyacrylamide gel electrophoresis; PEI, polyethylenimine; PP, protein phosphatase; SAMS, the synthetic peptide HMRSAMSGHLHLVKRR; SDS, sodium dodecyl sulphate.

### Author Contribution

R.W., A.L. and N.N. carried out the experimental work. All authors helped conceive and design the experiments, and contributed to the writing of the manuscript.

### Funding

This work was funded by the Medical Research Council UK and AstraZeneca. R.W. received a PhD studentship from the Medical Research Council and A.L. was funded by a studentship from the British Heart Foundation.

### Acknowledgements

We are grateful to David Carmena for preparation of the recombinant protein phosphatase 2C used in the present study and Dr Russell Miller (Pfizer Global Research and Development) for providing us with PF06685249. We thank Dr James Elliot and Thomas Adejumo in the Flow Cytometry Facility, MRC London Institute of Medical Sciences.

### Competing Interests

The Authors declare that there are no competing interests associated with the manuscript.

### References

- 1 Carling, D., Thornton, C., Woods, A. and Sanders, M.J. (2012) AMP-activated protein kinase: new regulation, new roles? *Biochem. J.* **445**, 11–27 doi:10.1042/BJ20120546
- 2 Hardie, D.G. (2014) AMP-activated protein kinase: maintaining energy homeostasis at the cellular and whole-body levels. *Annu. Rev. Nutr.* **34**, 31–55 doi:10.1146/annurev-nutr-071812-161148
- 3 Hardie, D.G. and Ashford, M.L. (2014) AMPK: regulating energy balance at the cellular and whole body levels. *Physiology* **29**, 99–107 doi:10.1152/physiol.00050.2013
- 4 Steinberg, G.R. and Kemp, B.E. (2009) AMPK in health and disease. *Physiol. Rev.* **89**, 1025–1078 doi:10.1152/physrev.00011.2008
- 5 Carling, D., Mayer, F.V., Sanders, M.J. and Gambliin, S.J. (2011) AMP-activated protein kinase: nature's energy sensor. *Nat. Chem. Biol.* **7**, 512–518 doi:10.1038/nchembio.610
- 6 Xiao, B., Heath, R., Saiu, P., Leiper, F.C., Leone, P., Jing, C. et al. (2007) Structural basis for AMP binding to mammalian AMP-activated protein kinase. *Nature* **449**, 496–500 doi:10.1038/nature06161
- 7 Xiao, B., Sanders, M.J., Underwood, E., Heath, R., Mayer, F.V., Carmena, D. et al. (2011) Structure of mammalian AMPK and its regulation by ADP. *Nature* **472**, 230–233 doi:10.1038/nature09932
- 8 Ross, F.A., Jensen, T.E. and Hardie, D.G. (2016) Differential regulation by AMP and ADP of AMPK complexes containing different  $\gamma$  subunit isoforms. *Biochem. J.* **473**, 189–199 doi:10.1042/BJ20150910
- 9 Hawley, S.A., Boudeau, J., Reid, J.L., Mustard, K.J., Udd, L., Mäkelä, T.P. et al. (2003) Complexes between the LKB1 tumor suppressor, STRAD  $\alpha/\beta$  and MO25  $\alpha/\beta$  are upstream kinases in the AMP-activated protein kinase cascade. *J. Biol. Chem.* **278**, 28 doi:10.1186/1475-4924-2-28
- 10 Hawley, S.A., Pan, D.A., Mustard, K.J., Ross, L., Bain, J., Edelman, A.M. et al. (2005) Calmodulin-dependent protein kinase kinase- $\beta$  is an alternative upstream kinase for AMP-activated protein kinase. *Cell Metab.* **2**, 9–19 doi:10.1016/j.cmet.2005.05.009
- 11 Hurley, R.L., Anderson, K.A., Franzone, J.M., Kemp, B.E., Means, A.R. and Witters, L.A. (2005) The  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinases are AMP-activated protein kinase kinases. *J. Biol. Chem.* **280**, 29060–29066 doi:10.1074/jbc.M503824200
- 12 Shaw, R.J., Kosmatka, M., Bardeesy, N., Hurley, R.L., Witters, L.A., DePinho, R.A. et al. (2004) The tumor suppressor LKB1 kinase directly activates AMP-activated kinase and regulates apoptosis in response to energy stress. *Proc. Natl Acad. Sci. U.S.A.* **101**, 3329–3335 doi:10.1073/pnas.0308061100
- 13 Woods, A., Dickerson, K., Heath, R., Hong, S.-P., Momcilovic, M., Johnstone, S.R. et al. (2005)  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab.* **2**, 21–33 doi:10.1016/j.cmet.2005.06.005
- 14 Woods, A., Johnstone, S.R., Dickerson, K., Leiper, F.C., Fryer, L.G.D., Neumann, D. et al. (2003) LKB1 is the upstream kinase in the AMP-activated protein kinase cascade. *Curr. Biol.* **13**, 2004–2008 doi:10.1016/j.cub.2003.10.031
- 15 Gowans, G.J., Hawley, S.A., Ross, F.A. and Hardie, D.G. (2013) AMP is a true physiological regulator of AMP-activated protein kinase by both allosteric activation and enhancing net phosphorylation. *Cell Metab.* **18**, 556–566 doi:10.1016/j.cmet.2013.08.019
- 16 Oakhill, J.S., Stell, R., Chen, Z.-P., Scott, J.W., Ling, N., Tam, S. et al. (2011) AMPK is a direct adenylate charge-regulated protein kinase. *Science* **332**, 1433–1435 doi:10.1126/science.1200094
- 17 Zhang, Y.-L., Guo, H., Zhang, C.-S., Lin, S.-Y., Yin, Z., Peng, Y. et al. (2013) AMP as a low-energy charge signal autonomously initiates assembly of AXIN-AMPK-LKB1 complex for AMPK activation. *Cell Metab.* **18**, 546–555 doi:10.1016/j.cmet.2013.09.005
- 18 Davies, S.P., Helps, N.R., Cohen, P.T. and Hardie, D.G. (1995) 5'-AMP inhibits dephosphorylation, as well as promoting phosphorylation, of the AMP-activated protein kinase. Studies using bacterially expressed human protein phosphatase-2C alpha and native bovine protein phosphatase-2AC. *FEBS Lett.* **377**, 421–425 doi:10.1016/0014-5793(95)01368-7
- 19 Sanders, M.J., Grondin, P.O., Hegarty, B.D., Snowden, M.A. and Carling, D. (2007) Investigating the mechanism for AMP activation of the AMP-activated protein kinase cascade. *Biochem. J.* **403**, 139–148 doi:10.1042/BJ20061520

- 20 Suter, M., Riek, U., Tuerk, R., Schlattner, U., Wallimann, T. and Neumann, D. (2006) Dissecting the role of 5'-AMP for allosteric stimulation, activation, and deactivation of AMP-activated protein kinase. *J. Biol. Chem.* **281**, 32207–32216 doi:10.1074/jbc.M606357200
- 21 Rajamohan, F., Reyes, A.R., Frisbie, R.K., Hoth, L.R., Sahasrabudhe, P., Magyar, R. et al. (2016) Probing the enzyme kinetics, allosteric modulation and activation of  $\alpha$ 1- and  $\alpha$ 2-subunit-containing AMP-activated protein kinase (AMPK) heterotrimeric complexes by pharmacological and physiological activators. *Biochem. J.* **473**, 581–592 doi:10.1042/BJ20151051
- 22 Cameron, K.O., Kung, D.W., Kalgutkar, A.S., Kurumbail, R.G., Miller, R., Salatto, C.T. et al. (2016) Discovery and preclinical characterization of 6-chloro-5-[4-(1-hydroxycyclobutyl)phenyl]-1*H*-indole-3-carboxylic acid (PF-06409577), a direct activator of adenosine monophosphate-activated protein kinase (AMPK), for the potential treatment of diabetic nephropathy. *J. Med. Chem.* **59**, 8068–8081 doi:10.1021/acs.jmedchem.6b00866
- 23 Hardie, D.G., Ross, F.A. and Hawley, S.A. (2012) AMP-activated protein kinase: a target for drugs both ancient and modern. *Chem. Biol.* **19**, 1222–1236 doi:10.1016/j.chembiol.2012.08.019
- 24 Xiao, B., Sanders, M.J., Carmena, D., Bright, N.J., Haire, L.F., Underwood, E. et al. (2013) Structural basis of AMPK regulation by small molecule activators. *Nat. Commun.* **4**, 3017 PMID:24352254
- 25 Calabrese, M.F., Rajamohan, F., Harris, M.S., Caspers, N.L., Magyar, R., Withka, J.M. et al. (2014) Structural basis for AMPK activation: natural and synthetic ligands regulate kinase activity from opposite poles by different molecular mechanisms. *Structure* **22**, 1161–1172 doi:10.1016/j.str.2014.06.009
- 26 Arad, M., Seidman, C.E. and Seidman, J.G. (2007) AMP-activated protein kinase in the heart: role during health and disease. *Circ. Res.* **100**, 474–488 doi:10.1161/01.RES.0000258446.23525.37
- 27 Arad, M., Benson, D.W., Perez-Atayde, A.R., McKenna, W.J., Sparks, E.A., Kanter, R.J. et al. (2002) Constitutively active AMP kinase mutations cause glycogen storage disease mimicking hypertrophic cardiomyopathy. *J. Clin. Invest.* **109**, 357–362 doi:10.1172/JCI0214571
- 28 Blair, E., Redwood, C., Ashrafian, H., Ostman-Smith, I. and Watkins, H. (2001) Mutations in the  $\gamma$ 2 subunit of AMP-activated protein kinase cause familial hypertrophic cardiomyopathy: evidence for the central role of energy compromise in disease pathogenesis. *Hum. Mol. Genet.* **10**, 1215–1220 doi:10.1093/hmg/10.11.1215
- 29 Gollob, M.H., Green, M.S., Tang, A.S.-L., Gollob, T., Karibe, A., Hassan, A. et al. (2001) Identification of a gene responsible for familial Wolff-Parkinson-White syndrome. *New Engl. J. Med.* **344**, 1823–1831 doi:10.1056/NEJM200106143442403
- 30 Yavari, A., Stocker, C.J., Ghaffari, S., Wargent, E.T., Steeples, V., Czibik, G. et al. (2016) Chronic activation of  $\gamma$ 2 AMPK induces obesity and reduces  $\beta$  cell function. *Cell Metab.* **23**, 821–836 doi:10.1016/j.cmet.2016.04.003
- 31 Salatto, C.T., Miller, R.A., Cameron, K.O., Cokorinos, E., Reyes, A., Ward, J. et al. (2017) Selective activation of AMPK  $\beta$ 1-containing isoforms improves kidney function in a rat model of diabetic nephropathy. *J. Pharm. Exp. Ther.* **316**, 303–311 doi:10.1124/jpet.116.237925
- 32 Marley, A.E., Sullivan, J.E., Carling, D., Abbott, W.M., Smith, G.J., Taylor, I.W. et al. (1996) Biochemical characterization and deletion analysis of recombinant human protein phosphatase 2C $\alpha$ . *Biochem. J.* **320**, 801–806 doi:10.1042/bj3200801
- 33 Davies, S.P., Carling, D. and Hardie, D.G. (1989) Tissue distribution of the AMP-activated protein kinase, and lack of activation by cyclic-AMP-dependent protein kinase, studied using a specific and sensitive peptide assay. *Eur. J. Biochem.* **186**, 123–128 doi:10.1111/j.1432-1033.1989.tb15185.x
- 34 Pinter, K., Grignani, R.T., Czibik, G., Farza, H., Watkins, H. and Redwood, C. (2012) Embryonic expression of AMPK  $\gamma$  subunits and the identification of a novel  $\gamma$ 2 transcript variant in adult heart. *J. Mol. Cell Cardiol.* **53**, 342–349 doi:10.1016/j.yjmcc.2012.05.017
- 35 Thornton, C., Snowden, M.A. and Carling, D. (1998) Identification of a novel AMP-activated protein kinase  $\beta$  isoform which is highly expressed in skeletal muscle. *J. Biol. Chem.* **273**, 12443–12450 doi:10.1074/jbc.273.20.12443
- 36 Cheung, P.C.F., Salt, I.P., Davies, S.P., Hardie, D.G. and Carling, D. (2000) Characterization of AMP-activated protein kinase  $\gamma$ -subunit isoforms and their role in AMP binding. *Biochem. J.* **346**, 659–669 doi:10.1042/bj3460659
- 37 Hawley, S.A., Ross, F.A., Chevzoff, C., Green, K.A., Evans, A., Fogarty, S. et al. (2010) Use of cells expressing  $\gamma$  subunit variants to identify diverse mechanisms of AMPK activation. *Cell Metab.* **11**, 554–565 doi:10.1016/j.cmet.2010.04.001
- 38 Carling, D. (2004) The AMP-activated protein kinase cascade — a unifying system for energy control. *Trends Biochem. Sci.* **29**, 18–24 doi:10.1016/j.tibs.2003.11.005
- 39 Yu, H., Fujii, N., Hirshman, M.F., Pomerleau, J.M. and Goodyear, L.J. (2004) Cloning and characterization of mouse 5'-AMP-activated protein kinase  $\gamma$ 3 subunit. *Am. J. Physiol. Cell Physiol.* **286**, C283–C292 doi:10.1152/ajpcell.00319.2003
- 40 Buchan, D.W.A., Minneci, F., Nugent, T.C.O., Bryson, K. and Jones, D.T. (2013) Scalable web services for the PSIPRED protein analysis workbench. *Nucleic Acids Res.* **41**, W349–W357 doi:10.1093/nar/gkt381
- 41 Babu, M.M., van der Lee, R., de Groot, N.S. and Gsponer, J. (2011) Intrinsically disordered proteins: regulation and disease. *Curr. Opin. Struct. Biol.* **21**, 432–440 doi:10.1016/j.sbi.2011.03.011
- 42 Pinter, K., Grignani, R., Watkins, H. and Redwood, C. (2013) Localisation of AMPK  $\gamma$  subunits in cardiac and skeletal muscle. *J. Muscle Res. Cell Motil.* **34**, 369–378 doi:10.1007/s10974-013-9359-4
- 43 Bultot, L., Jensen, T.E., Lai, Y.C., Madsen, A.L., Collodet, C., Kviklyte, S. et al. (2016) Benzimidazole derivative small-molecule 991 enhances AMPK activity and glucose uptake induced by AICAR or contraction in skeletal muscle. *Am. J. Physiol.* **311**, E706–E719 PMID:27577855
- 44 Mitchelhill, K.I., Mitchell, B.J., House, C.M., Stapleton, D., Dyck, J., Gamble, J. et al. (1997) Posttranslational modifications of the 5'-AMP-activated protein kinase  $\beta$ 1 subunit. *J. Biol. Chem.* **272**, 24475–24479 doi:10.1074/jbc.272.39.24475
- 45 Cool, B., Zinker, B., Chiou, W., Kifle, L., Cao, N., Perham, M. et al. (2006) Identification and characterization of a small molecule AMPK activator that treats key components of type 2 diabetes and the metabolic syndrome. *Cell Metab.* **3**, 403–416 doi:10.1016/j.cmet.2006.05.005