

Interactions In Space For Archaeological Models*

T.S. Evans, R.J. Rivers

Theoretical Physics, Imperial College London,
London, SW7 2AZ, U.K.

C. Knappett

Department of Art Sidney Smith Hall, Room 6036 100 St George Street,
University of Toronto Toronto Ontario M5S 3G3

Abstract

In this article we examine a variety of quantitative models for describing archaeological networks, with particular emphasis on the maritime networks of the Aegean Middle Bronze Age. In particular, we discriminate between those gravitational networks that are most likely (maximum entropy) and most efficient (best cost/benefit outcomes).

1 Introduction

In archaeology, the primary sources of information are usually finds from specific sites. In order to understand the social, cultural and political context of these finds it is essential that we understand how sites relate. However, it often requires so much effort to obtain the physical information from a single site that there is a danger that we become ‘site-centric’. Despite our best attempts at deducing relationships from the artefacts found at them, there is often little direct information about how sites interact. Quantitative modelling provides one response to this challenge. Good quantitative modelling can be insensitive to poor data, make assumptions and biases clear and debatable, and can allow us to provide possible answers to questions that could not be asked any other way. In the best case, such answers can be checked later from the archaeological record.

One of the major problems that we attempt to address is to determine how the relationships between (archaeological) sites are conditioned by geographical space; to what extent does the exchange between them transcend their geographical constraints? As the basis for our modelling we have found the language of complex networks particularly useful as a network is *both* a set of vertices — the sites — *and* a set of edges, or links — the relationships between them. Socio-spatial networks do not arise and develop arbitrarily. Their maintenance incurs explicit and implicit ‘costs’, both in sustaining sites and in supporting links [5]. The common agency in the networks that we shall discuss below is the need to regulate or distribute these costs. We now encounter a conundrum. Despite these general features, the modelling of social networks requires very specific tailoring for individual case studies. This is contingent on the level of social organisation, the nature of the exchange, relevant distance scales, travel technology, etc. of the society being studied. Thus,

*Contribution to special issue of Advances in Complex Systems from the conference ‘Cultural Evolution in Spatially Structured Populations’, UCL September 2010. Imperial/TP/11/TSE/1

for exemplary purposes, we have found it useful to work with one particular data set, namely the Middle Bronze Age (MBA) to Late Bronze Age I (LB I) S.Aegean. Nonetheless, we shall try to be as general as possible within this framework.

2 Networks

As we have said, networks primarily consist of vertices (or nodes), representing agents or populations or resources, and edges (or links), which represent the exchange between them, from the physical trade of goods to the transmission of ideas or culture. The MBA Aegean has several distinct features that lends itself to a network analysis. There is a relatively clear temporal delineation both at the beginning of the MBA and at the end of LB I. From the start of the MBA (c.2000 BC), society on Crete, often dubbed ‘Minoan’, sees a series of innovations, such as monumental architecture at palatial centres, new craft technologies and more sustained long-distance exchange. This last trait is most likely enabled by the innovation of the sail that seems to have occurred around this time, supplementing what was previously a paddle-based maritime technology. The end of our period of study, LB I, c.1450 BC, sees the destruction of the Minoan palaces and a swing in power and influence to the Greek mainland; this marks a useful delineation for analytical purposes. The physical boundaries of our focus for study are also relatively well defined, though long range trade beyond the Aegean was essential e.g. in providing tin, and questions regarding the relationship to Egypt remain important. Nevertheless, as a first approximation, limiting our analysis to the Aegean region provides a good example of where modelling may be of particular use. There is a large amount of material from this era, with Akrotiri and Minoan palaces such as Knossos being rich sources. In this paper we will describe our network model, **ariadne**, and show how it is related to other approaches to archaeological modelling, using these MBA maritime exchange networks as a testing ground. In so doing we will go beyond the qualitative descriptions given in our previous work [12, 17].

2.1 Defining Our Vertices

One of the main ways in which the nature of society impacts on the construction of networks is in determining what constitutes vertices which, in principle, could vary in scale from individuals/households to communities to islands. Ironically this also reflects the fact that most network analysis is also site-centric (although graph theory does provide some tools to get around this [13, 11]). In general, island archipelagos are ideal for networks as the geography provides a natural definition for our vertices. Our working hypothesis is that the detailed behaviour of individuals can be subsumed in the behaviour of the ‘island’ community. In this regard, we have taken each vertex to correspond to an island, or a large part of an island or, in the case of N. Crete and other coastal areas, to an isolated centre of population. The latter sites effectively behave as islands because of the difficulty of land travel. This hypothesis is contingent on a maritime technology permitting long-distance travel that enables the island (or ‘island-like’ coastal site) to be thought of as the basic unit for ‘trade’ and has been discussed in detail elsewhere [12, 17]. However, once we

move away from individuals to communities, it is apparent that each vertex has two very different attributes; its resource base and the exploitation of these resources. The former, the carrying capacity of the vertex, is part of our *input*. The latter is represented by the ‘size’ of the community that inhabits it. That is part of our *output*.

In many models there is no discrimination between carrying capacity and population (i.e. one is taken proportional to the other) and, often, the carrying capacities of the sites are set equal. This may well be appropriate for some cases such as pre-urban civilisations where land was occupied in small villages of roughly equal size. The work of Broodbank [4] on the Early Bronze Age in the Cyclades is a good example of this approach. There the potential cultivable land of each island was assessed and the density of sites on habitable land was chosen to be uniform. However we are interested in an era when significant differences emerge between the size of different sites. These differences reflect more than just local differences in resources and we are looking to understand how interactions with other sites may have supported very large sites. Thus it is important that site sizes are variable in our model, both in terms of the fixed inputs given and in terms of the outputs, since we expect a complex interdependence of interactions and site sizes. We have chosen a set of major known islands or coastal sites as representing the most important locations for vertices. See Figure 1, with details in Table 1. The choice is made on the basis of the archaeological record. It is sufficient for our purposes in Table 1 to classify the carrying capacities of the sites as ‘small’ (S), ‘medium’ (M) or ‘large’ (L). What this means quantitatively will be explained later. In so doing we are assuming that other locations are peripheral to the dynamics of the whole system or at least their effect is well represented by the dominant site in their region.




Figure 1: Thirty nine important Aegean sites of the MBA (Middle Bronze Age).

1. Knossos (L)	14. Kea (M)	27. Mycenae (L)
2. Malia (L)	15. Karpathos (S)	28. Ayios Stephanos (L)
3. Phaistos (L)	16. Rhodes (L)	29. Lavrion (M)
4. Kommos (M)	17. Kos (M)	30. Kasos (S)
5. Ayia Triadha (L)	18. Miletus (L)	31. Kalymnos (S)
6. Palaikastro (L)	19. Iasos (M)	32. Myndus (M)
7. Zakros (M)	20. Samos (M)	33. Cesme (M)
8. Gournia (L)	21. Petras (L)	34. Akbuk (M)
9. Chania (L)	22. Rethymnon (L)	35. Menelaion (S)
10. Thera (M)	23. Paroikia (M)	36. Argos (M)
11. Phylakopi (M)	24. Amorgos (S)	37. Lerna (M)
12. Kastri (M)	25. Ios (S)	38. Asine (S)
13. Naxos (L)	26. Aegina (M)	39. Eleusis (M)

Table 1: Thirty-nine sites of the MBA Aegean. Note Knossos as site 1 and Akrotiri, subsequently destroyed in the eruption of Thera, as site 10. The S,M,L indicates a rough assignment of size — small, medium or large.

As for notation, we denote vertices using lowercase mid Latin indices $i, j, \dots$. The fixed carrying capacities of sites will be denoted S_i (habitable land available). In some models the total population of a site may differ from this as we will indicate.

2.2 Defining our Edges

The edges represent the interactions and can be seen within the framework of different types of space e.g. artefact space [28, 29, 32]. However we take the view that archaeology is rooted in geographical space and it is an unavoidable constraint on interactions.

This means that one of the most important inputs to our model will be a table of distances between our sites. These may be simple Euclidean (as-the-crow-flies) distances. A more sophisticated approach will use estimates of typical journey times. This will be sensitive to the technology available and could require information on currents, typical winds, slopes, and even security. For instance in our examples we have analysed the layout of islands in the Aegean by hand to take promontories into account, but not the effects of winds and currents. Likewise for land travel we picked routes by hand that reflected current three-dimensional geography, arguing that the accuracy of a full least cost path analysis is unnecessary given the level of approximation. Finally in the examples shown here, we choose to penalise land travel over sea travel by a ‘friction’ coefficient of 3.0. Results are largely insensitive to reasonable values of the coefficient (larger than unity), primarily affecting the width of Crete. We will not look at this issue in any depth here, rather we will take it that these effective distances between sites are given. The issue we address in detail is how to model the actual interactions, given a framework of sites and their separations.

For notation, if the attribute of an edge from i to j is A_{ij} then the matrix $\mathbf{A}$ of these elements is an adjacency matrix for the network. When A_{ij} defines the flow from site i to site j it will be denoted by F_{ij} . Note that unless otherwise stated,

models exclude self-loops excluded, that is $F_{ii} = 0$. The effective distance from site i to site j will be d_{ij} . For all examples here, this is symmetric $d_{ij} = d_{ji}$, although as discussed above this is not always appropriate.

3 Geographical Models

Our emphasis, and that of the other main models that we shall discuss, is based on network analysis, as we have already indicated. However, before going into details we stress that there are alternative approaches that work more with ‘zones of influence’ than exchange directly. Although vertices are preserved, networks themselves are abandoned, and we begin by discussing such models briefly.

3.1 Geographical Models without Networks

There are several examples of models for archaeology, based on significant sites such as we have defined, yet without any explicit edges defined and so no network. If we treat all sites as equal we can construct partitions (or partial partitions) of geographical space into something like ‘zones of influence’. This is a very old idea, reincarnated in its simplest form as simple Voronoi, or Thiessen, polygons, each of which contain all the points in space for which the single site at the centre of that polygon is the closest site. They can be used to indicate zones of control of each site, e.g. for Etruscan Cities using Euclidean distances [23]. The XTent model of Renfrew and Level [24] is a generalisation for cases with different site sizes. Here site i of size S_i is deemed to ‘dominate’ site j of size S_j if $\tan(\theta) > d_{ij}/(S_i - S_j)$ where d_{ij} is the distance between the two sites and $\tan(\theta)$ is a parameter of the model. In the XTent model one can work with just the sites (rather than all points in space needed for Voronoi diagrams) and we can represent the resulting hierarchy of sites as a directed network. An example of how this may be used with modern GIS techniques to gain a good estimate of actual walking time, rather than using simple Euclidean distance, has been given for Neopalatial Crete by Bevan [3]. However, such a network representation adds little and the model is not usually visualised in this way, e.g. see [24, 3].

3.2 Simple Geographical Networks

There are two simple ways to construct a network which captures non-trivial information about the global interactions between sites, both based on a thresholding of the distance matrix.

The first is a ‘maximum distance network’ (MDN) in which an edge from site i to site j is connected if the distance is less than some model parameter D . In the language of links, we might construe this as imposing a link likelihood $A_{ij} = \Theta(D - d_{ij})$ for the journey, where $\Theta(x)$ is the step function, taking value 1 for $d_{ij} < D$ and zero otherwise. MDN is used in many fields to establish distance scales, e.g. as a model of ad-hoc wireless networks formed between mobile devices [30]. A great deal of work exists on these types of models (particularly ‘random geometric graphs’ [22, 30]), but they appear not to have been widely used for models in archaeology.

See Fig. 2 for the Aegean when $D = 100km$ and $110km$. [The reader should impose the pattern of sites in Fig. 2 on the geographic template of Fig. 1, to identify the sites.] At this value the S. Aegean splits into four identifiable zones; Crete, the Cyclades, the Dodecanese and the Peloponnese. If we increase the distance to $D = 130km$, these regions connect [not shown]. These are relevant values for the MBA, for which the dominant marine technology is sail, with vessels routinely capable of travelling large distances in single journeys. As a marker we note that the distance from Knossos to Akrotiri is in excess of $100km$, which we believe was achievable in a single journey. Typically we take D to be $100km$ or somewhat greater.




Figure 2: On the left an example of an MDN (Maximum distance network) while the right hand network is a k -nearest-neighbour graph or PPA (Proximal Point Analysis) network. These are for the 39 MBA Aegean sites of Fig. 1, using estimated travel times where land travel has a friction coefficient of 3 compared to sea travel. In the MDN network, thick edges in black are present if the separation is less than $D = 100km$ (thin grey lines indicate edges between 100 and 110km). In the PPA network thick black lines indicate the case where each vertex is connected to its three nearest neighbours $k = 3$ (thin grey lines indicate connections to fourth and fifth nearest neighbours). In both cases the colour of vertices indicates the connectivity of the network defined by the thick black edges only.

The second thresholding method is more common in archaeology where it is known as PPA (Proximal Point Analysis) [31, 16, 15, 4, 7, 32] though it has received less attention in other fields, e.g. as ‘ k -nearest-neighbour graphs’ [21, 1]. Here each site is connected to its k nearest neighbours (k is a parameter of the model) to give a directed network, see Fig. 2, though the directions are usually ignored in the literature. An example is given in Fig. 2 for $k = 3$ nearest neighbours. It differs strongly from its MDN counterpart in that, by definition, sites on the extrema of the map that are separated from their neighbours will connect nonetheless, despite the distances, because the assumption is that they will connect somehow, independent of how easy this may be. As a result, there is a tendency for sites to be connected in ‘strings’.




Figure 3: A simple Gravity Model with edge weights $F_{ij} = S_i S_j f(d_{ij}/D)$ for the 39 MBA Aegean sites of Fig. 1, using estimated travel times where land travel has a friction coefficient of 3 compared to sea travel. The function $f(x)$ used is given in (1) with a distance scale of $D = 100\text{km}$. Vertices are proportional to the site size (0.33, 0.67 or 1.00 for small, medium and large sites). On the right all edges are shown but with their thickness proportional to the flow F_{ij} . The same network is shown on the right but now all links with $F_{ij} < 0.23$ are shown only as thin grey lines. In addition the colour of vertices indicates the connectivity of the network of strong ($F_{ij} > 0.23$) links.

The MDN can be thought of as imposing prohibitive costs on implementing long single journeys and PPA as imposing prohibitive costs on sustaining more than a few social interactions. The last simple model worth mentioning is the elementary ‘Gravity model’ [10, 20]. For this the flow F_{ij} from site i to site j is assumed to take the form $F_{ij} = S_i S_j f(d_{ij}/D)$ (with $F_{ii} = 0$) where the distance ‘potential’ $f(x)$ is a monotonic decreasing function, reflecting the ease of travel from i to j or, as we shall see, the effective cost of travel. In the context of the MBA Aegean we want short trips by sea to be relatively easy, or of low cost, while we want a strong cutoff or penalty for single trips of distance D or more. We stress that, unlike for the MDN, D is no longer an absolute cut-off, but, with $f(1) = 0.5$, sets the scale above which single journeys become increasingly difficult. D therefore takes somewhat smaller values than we would read off from the MDN.

We have chosen a generic form for $f(x)$ that is a smoothed out version of the MDN, namely

$$f(x) = [1 + x^{\beta_1}]^{-\beta_2}. \quad (1)$$

In our work we have used only $\beta_1 = 4.0$ and $\beta_2 = 1.0$. As long as there is a plateau for $x < 1$ followed by rapid falloff for $x > 1$ the outputs are similar. As we see in Fig.3, it produces a dense network, although many of the long distance links will be weak. In particular either of the previous two thresholding techniques could be used to create a network from the flows F_{ij} . In this sense this Gravity model can be seen as a generalisation of the previous two models to sites of different sizes.

In the first two models sites are of fixed and equal size. This, and subsequent,

Gravity Models avoid this limitation. In particular, by aggregating island communities into an island ‘centre of population’ and a similar conflation of inter-island exchange, they assume that, at the island level, the whole is the sum of the parts. By this coarse-graining, they minimise our need to have detailed local knowledge, in that sense making the best of poor data. However, the elementary model above fails to capture the rich variety of possible interactions — there is no feedback between interactions and site size. In arriving at the flow from i to j , we could change the location of all other sites and it would make no difference.

4 Optimal Geographical Networks

The models which are most fit for our purpose are ones in which the strength of the interactions reflect both the local geographical topology between two sites but also the wider regional structure in which these sites reside. In different ways the two model types that have been developed most fully are optimal models, either looking for the ‘most likely’ networks, all other things being equal, or the most ‘efficient’ networks.

The simplest example of the former is that of the *doubly constrained* Gravity Model (DCGM). Here the flow takes the same form as before but with coefficients determined self-consistently:-

$$F_{ij} = A_i O_i B_j I_j f\left(\frac{d_{ij}}{D}\right), \quad \frac{1}{A_i} = \sum_k B_k I_k f\left(\frac{d_{ik}}{D}\right), \quad \frac{1}{B_j} = \sum_k A_k O_k f\left(\frac{d_{kj}}{D}\right). \quad (2)$$

Although not immediately transparent, solutions of this form are in fact turning points with respect to the $N(N-1)$ variables F_{ij} (as $F_{ii} = 0$) of the Hamiltonian

$$H = \sum_{(i,j)} F_{ij} (\ln(F_{ij}) - 1) - \sum_i \alpha_i \left[O_i - \sum_j F_{ij} \right] - \sum_j \alpha'_j \left[I_j - \sum_i F_{ij} \right] - \beta \left[C - \sum_{ij} (F_{ij} c_{ij}) \right]. \quad (3)$$

The first term in (3) is the negative of the entropy of a network of flows F_{ij} . Maximising this entropy term produces the most likely distribution of the total exchange, $\sum_{i,j} F_{ij}$, amongst all the possible edges, in the absence of any further knowledge as to how exchange occurs. As all edges are treated equally by this entropy term, it is the remaining terms which produce a non-trivial solution.

The N parameters $\{\alpha_i\}$ are Lagrange multipliers which enforce constraints on the total outflow, $O_i = \sum_j F_{ij}$, from each site i . The $\{\alpha'_j\}$ do the same job for the total inflow of site j , $I_j = \sum_i F_{ij}$. Here O_i and I_j are input parameters of the model which we have taken proportional to the carrying capacity/population. The simplicity of the model allows for an algebraic solution for these Lagrange multipliers which allows us to replace the $\{\alpha, \alpha'_i\}$ with the normalisations $\{A_i\}$ and $\{B_j\}$ in (2). Enforcing these constraints ensures the flow along the edge from i to j depends directly on the flow from i to all other sites and upon the flow into j from all other sites. The largest contributions will be from sites near to i and/or to j so the

interactions do depend on the whole region. Changing the location of other sites in the region will alter the flow from i to j , the type of behaviour we are seeking.

The last Lagrange multiplier, β , is associated with the total ‘cost’ C for the pattern of flows. Here the cost per unit of flow is given by c_{ij} for a link from i to j . The total cost C could reflect many factors rather than monetary and so it is rarely known in a physical example. So rather than specify the total cost as a parameter of the model, it is normal to keep β as a parameter. Larger (smaller) β means higher (lower) costs. In practice β dependence becomes absorbed into one or more parameters into the travel, or distance, ‘potential’ $f(x)$ of (1) on making the identification $f(d_{ij}/D) = \exp(-\beta c_{ij})$. Thus larger D means cheaper costs and corresponds to larger β .

An exemplary network of the DCGM is given in Fig. 4. The drawback of this model is that we have to fix both input and output site sizes and these do not respond to the pattern of interactions which emerges. For example, in our MBA Aegean context, Knossos can never alter the volume of economic, cultural or social exchange from the values we gave it at the start. Further, because inflows and outflows are fixed, we get a repeat of the pattern in the PPA that remote sites are still strongly connected, even if distances are large.




Figure 4: Two different optimal Gravity models for the 39 MBA Aegean sites of Fig. 1, using estimated travel times where land travel has a friction coefficient of 3 compared to sea travel. In both cases the distance scale is $D = 100\text{km}$. Vertices are proportional to the site size (0.33, 0.67 or 1.00 for small, medium and large sites). On the left solutions for the doubly constrained Gravity model (DCGM) are shown. On the right is the Rihll and Wilson Gravity model (RWGM) (6) with $\gamma = 1.3$.

Some of these issues are addressed in the work of Rihll and Wilson [25, 26] who, instead, use a model originally devised to study the emergence of dominant retail centres [6, 34, 35, 8, 9], which we now summarise. The basic Rihll and Wilson Gravity model (RWGM) gives the flow F_{ij} from site i to site j as

$$F_{ij} = A_i O_i I_j^\gamma f(d_{ij}/D), \quad A_i^{-1} = \sum_k I_k^\gamma f(d_{ik}/D). \quad (4)$$

As before, O_i is the total outflow of site i , $O_i = \sum_j F_{ij}$ and I_j is the total inflow of site j , $I_j = \sum_i F_{ij}$. The difference from the DCGM is that, while outflows O_i are still input parameters of the theory (taken to be proportional to site carrying capacity), the inflows I_j are now outputs determined by the model. These are used by Rihll and Wilson to assign an importance to a site¹. Solutions to this nonlinear equation are in fact turning points of the Hamiltonian

$$H = \sum_{i,j} F_{ij} (\ln(F_{ij}) - 1) - \sum_i \alpha_i \left[O_i - \sum_j F_{ij} \right] - \beta \left[C - \sum_{i,j} (F_{ij} c_{ij}) \right] - \gamma \left[X - \sum_{i,j} F_{ij} (\ln(\sum_k F_{kj}) - 1) \right], \quad (5)$$

again understood as maximising entropy. As before, we have encoded β and the cost function c_{ij} in the distance potential f of (1). Likewise we retain the γ Lagrange multiplier in the solution (4) rather than specify the unknown X in the Hamiltonian (5). An important solution is where the flow into one or more sites is zero so that $I_j = F_{ij} = 0$ for a finite number of sites j , but for all values of index i .

In principle there are many ways to find these solutions. Rihll and Wilson generate a sequence of values $\{I_j(0), I_j(1), \dots, I_j(t), \dots\}$ where

$$I_j(t+1) = \sum_i A_i(t) O_i(I_j(t))^\gamma f(d_{ij}/D), \quad A_i^{-1}(t) = \sum_k I_k^\gamma(t) f(d_{ik}/D) \quad (6)$$

Provided the parameters are chosen suitably the limiting value of this sequence, $\lim_{t \rightarrow \infty} I_j(t)$, generates flows which optimise H of (5). Rihll and Wilson use an “egalitarian hypothesis” and set the fixed outputs and the initial inputs all equal $O_i = I_j(t=0) = 1$. This is deemed equivalent to the assumption that “all sites were approximately equal in size and importance at the beginning of the period under consideration”, since there is often little information available to make any other hypothesis.

In principle, t could just represent some computational time and have no physical significance. However it is sometimes interpreted as physical time [34] in which case this represents a whole further set of assumptions. In particular one can view the finite difference equation as the Euler method for solving the differential equation

$$\frac{dI_j}{dt} = \epsilon \left(\sum_i A_i O_i(I_j(t))^\gamma e^{(-\beta c_{ij})} - K I_j \right) \quad (7)$$

where K and ϵ are new constants.

An example of the RWGM is given in Fig. 4 for $\gamma > 1$, which generates hubs. Taking values $\gamma < 1$ recreates networks closer in structure to those of the DCGM (not shown). As with the DCGM, the enforced outflows ensure that however remote a site may be, it will always be connected. As a result, although we have not been able to show this in the limited Figures in the text, these models give networks whose important links are largely insensitive to our choice of D , within reason (e.g. varying D from 80km to 130km).

Whatever choice we make, one possibility is to exclude such peripheral sites, as we have for our chosen MBA sites and for the Iron Age Greek mainland sites chosen by Rihll and Wilson.

¹In a second self-consistent version, Rihll and Wilson set $O_i = I_i$. The model parameters now include an initial value for I_i for each site. We will not consider this variant further.

5 ariadne

We conclude with a discussion of the model, named **ariadne**, that we have developed over a period of time for examining MBA maritime networks in the S. Aegean [12, 17]. We wanted site sizes to be informed, but not absolutely constrained, by the local resources, with physical geography placing similar limitations on interactions. Rather than look for the most likely networks, as we have been doing with gravity models, we are looking for the most efficient networks from the viewpoint of the costs and benefits that the network demands and provides.

In our model the input parameters are the fixed local site resources, S_i , and the site locations plus four independent parameters characterising the component parts of the ‘Hamiltonian’ or ‘social potential’ H , understood as representing the costs *minus* the benefits in sustaining the network. As outputs the model produces the flows F_{ij} between different sites i and j but now also produces self-loop terms $F_{ii} \geq 0$ to represent parts of the population not involved in interactions. The values are found by minimising the following Hamiltonian² H (i.e. maximising benefits with respect to costs):-

$$H = -\kappa \sum_i 4W_i(1 - W_i/S_i) - \lambda \sum_{i,j} F_{ij}f(d_{ij}/D)W_j + J \sum_i W_i + \mu \sum_{i,j} F_{ij}. \quad (8)$$

where $W_i = \sum_j F_{ij}$ represents a variable total population at site i . Note that in this model W_i is not the same as the total output $O_i = \sum_{j|j \neq i} F_{ij} = W_i - F_{ii}$ if some of the population at site i stays at home, $F_{ii} > 0$.

The first term of H is proportional to κ is a measure of the benefit of local productivity. It is minimised by setting $W_i = S_i/2$, only over-exploitation $W_i > S_i$ produces positive energy (negative benefits). This term is similar in form to $W_i[\ln(W_i) - 1]$ which would be present if we were maximising the entropy associated with the arrangement of the total ‘population’, $\sum_j W_j$, amongst the sites. This is in contrast to the optimal Gravity models where it is the entropy associated with the distribution of people amongst the *edges* which is maximised.

The second term, proportional to λ , provides benefits of exchange, as tempered by geography through the function f of (1). Links over distances much longer than the parameter D produce relatively little benefit to reflect the low probability of such direct links being maintained. The benefit of a link is also deemed to be in proportion to the target site’s total size W_j . As the source site’s size is related to the flow from i to j , F_{ij} , this term is reminiscent of the product of source and target sites sizes that characterise gravity models as in (2). However our product is in the Hamiltonian, not in the solution for the flows. Our nonlinear dependence on the site sizes in this term is a key distinction between our models and the optimised gravity models presented earlier. Thus, an increase in interactions from i to j will produce a positive feedback and increasing site j ’s size will produce even greater benefits, encouraging in turn more growth in the flow between the two sites, possibly (but not necessarily) requiring an increase in the size of site i . The first term proportional to κ

²In previous papers [12, 17, 27] we worked in terms of fractional values for total site size v_i and for flows e_{ij} with $W_i = S_i v_i$ and $F_{ij} = S_i v_i e_{ij}$. The Hamiltonian is then $H = -\kappa \sum_i 4S_i v_i(1 - v_i) - \lambda \sum_{i,j} (S_i v_i) e_{ij} f(d_{ij}/D)(S_j v_j) + J \sum_i S_i v_i + \mu \sum_{i,j} S_i v_i e_{ij}$.

prevents this process from running away but it does allow the volume of interactions to grow if the dynamics favours it³.

The last two terms, proportional to J and μ are the costs in sustaining the population and the total flow. The J term is equivalent to setting the total output rather than individual site outputs in Gravity models, so that $J = \sum_i \alpha_i$ in the optimal Gravity models of (3) and (5). The term proportional to μ is different only if $F_{ii} \neq 0$ for some sites i , emphasising that the full potential for interactions need not be met in our model — some ‘people’ can stay at home and need not ‘trade’ if there is insufficient benefit. This is another improvement over gravity models as in **ariadne** a remote site i separated by several D from the nearest neighbour will probably not interact, $F_{ij} = 0$ if $i \neq j$, and will probably maintain a population of about $S_i/2$ based on the benefits of local resources encoded through the κ term.

We also impose some further constraints on our variables. Clearly we demand that $W_i \geq 0$ and $F_{ij} \geq 0$. More importantly we also impose a short range cutoff such that, for sites separated by less than a certain minimum distance, $d_{ij} > d_{\min}$, we set $F_{ij} = 0$. Technically, if we impose this cutoff then we can split any site into two pieces separated by less than $d_{\min}$, and these two small sites behave like a single unified site. In this way we have an explicit scale for our coarse graining. We suggest that this scale be set by the distance of travel in a day by land, say 10km. In our case it only effects three sites on the Southern coasts of Crete.

We note that the scale of the Hamiltonian is irrelevant so we usually choose $\kappa = 1.0$ leaving us with the three Hamiltonian parameters λ, J, μ (the last two may be of any sign) plus the distance scales D and $d_{\min}$. The final parameters are the fixed site resources S_i along with their separations d_{ij} . However, unlike for the previous models, the networks are now more sensitive to the values of D chosen. For D noticeably less than 100km the networks becomes fragmented, whereas for D more than 130km, say, the networks have a tendency to become very dense. This is a direct reflection of the typical distance scale $\approx 130km$ (not quite identified with D , see earlier) for which the sites form a connected whole. We can invert the problem to say that the existence of a strong maritime network requires a marine technology whose single journeys match the distances required to produce a network. As we noted earlier, this is plausible, given the 100km plus of the distance from Knossos to Akrotiri.

To produce our networks, we find a set of values for our N^2 parameters F_{ij} which produce an approximate minimum for H of (8) using a Monte Carlo method. Unlike the RWGM, **ariadne** is not strictly deterministic, and only discriminates between comparably optimal solutions statistically. This introduces a stochastic element which we feel is appropriate given the uncertainties in modelling such a complex system. Of course we do not know the values for our input parameters, so our approach is to look at how the networks change as we change our parameters. Some examples are given in Fig. 5. For instance reducing the benefits of interaction, reducing λ , destroys the longer distance weak links that are maintaining the global connectivity. This also produces a general reduction in site sizes despite the maintenance of more localised networks. Likewise, the variation in city sizes, a feature

³While one could imagine other dependencies on the target site size, say $(W_j)^\gamma$ with some new model parameter γ , we have not found such an additional parameter necessary.

which emerges in this period and exemplified by the size of Knossos, can be produced by emphasising the non-linear terms in our Hamiltonian (also an attractive feature of the RWGM). Finally, though we produce equilibrium networks, we can tackle questions about time evolution. Slow evolution can be simulated by comparing different values for our parameters, mimicking adiabatic changes in physical thermodynamic problems. However we can also look at ‘quenches’, as in removing Akrotiri after the eruption of Thera [18].

6 Conclusions

Despite the well-established nature of many of the network modelling approaches discussed here, a detailed statistical comparison remains to be given. The main purpose of this article has been to start this process by considering the relationship among popular archaeological models which take geography as their primary driving force. Some of their key properties are summarised in table 2.

Model Type	(Fig.)	Network Type			Site Sizes	
		W	D	Other	Outflows	Inflows
Voronoi		none			Fixed	Equal
XTent		UW	D	Trees	Fixed & Different	
MDN	(2)	UW	UD		Fixed	Equal
PPA	(2)	UW	D	$k_{\text{out}} > 0$	Fixed	Equal
Simple GM	(3)	W	UD		Fixed Different	Fixed Different
DCGM	(4)	W	D	$k_{\text{in}}, k_{\text{out}} > 0$	Fixed Different	Fixed Different
RWGM	(4)	W	D	$k_{\text{out}} > 0$	Fixed Different	Variable
ariadne	(5)	W	D		Variable	Variable

Table 2: Summary of some of the features of different models of interactions for geographically embedded systems. (U)W = (un)weighted, (U)D = (un)directed. $k > 0$ ($k_{\text{out}} > 0$) indicates that a model forces all sites to have at least one (outgoing) edge, however isolated it may be.

Our emphasis has been on the last two models, the Rihll and Wilson Gravity Model (RWGM) [25, 26] and our own response to these issues, the **ariadne** model [12, 17]. Both of these use optimisation as a key principle, drawing on the wealth of experience from statistical physics. However it is interesting to highlight what we consider to be positive aspects of our **ariadne** model, and how these compare with the other models discussed here:-

- **ariadne** and the gravity models give weights to interactions so that we can have both strong and weak links in the nomenclature of Granovetter [14].
- **ariadne** gives the most likely arrangement of ‘population’ over distinct sites, whereas Gravity models optimise entropy of links.
- **ariadne** has no absolute constraint on individual site sizes. In contrast DCGM constrains both inflow and outflow while RWGM does not constrain inflow at sites.

- The more isolated a site is in **ariadne**, the less it will participate. However in gravity models and PPA, the constraints force even the most remote sites to be integrated into the system.
- More generally, both the RWGM and **ariadne** have to be understood statistically. From the viewpoint of ensemble theory RWGM is a microcanonical description of network flow, whereas **ariadne** provides a grand canonical description.

It can be argued that different models are capturing different types of phenomena. For instance, the XTent and RWGM (for $\gamma > 1$) are defining zones of control, which sites dominate their neighbours, rather than defining interaction patterns. In practice, the strengths of models only become apparent in specific applications to the archaeological record. Whereas PPA has been used for EBA Cyclades with rowing technology [4] we have used **ariadne** to describe the MBA S.Aegean with its sailing technology [12, 17, 27, 18]. The maritime networks here are characterised by an ability to make single journeys on a scale that is the typical length for which the S.Aegean is largely connected. For networks requiring travel on a scale of several days journey (e.g. the East Mediterranean in the LBA) the RWGM may be appropriate. We shall consider this elsewhere.

There are many properties that we have not addressed. In particular, the strongly non-linear behaviour of RWGM and **ariadne** mean that the resulting networks have a propensity for instability. For our model these are given in more detail elsewhere [12, 17, 27, 18] to which we refer the reader.

References

- [1] Balister, P., Bollobás, B., Sarkar, A., and Walters, M., Connectivity of random k -nearest-neighbour graphs, *Adv. in Appl. Probab.* **37** (2005) 1–24.
- [2] Bevan, A., Computational models for understanding movement and territory, in *Sistemas de Información Geográfica y Análisis Arquelógico del Territorio. V Simposio Internacional de Arqueología de Mérida, Anejos de Archivo Español de Arqueología.*, eds. Mayoral, V. and Celestino, S. (2008).
- [3] Bevan, A., Political geography and palatial Crete, *J.Mediterranean Arch.* **23** (2010).
- [4] Broodbank, C., *An Island Archaeology of the Early Cyclades* (CUP, 2000).
- [5] Barthélémy, M., Spatial Networks, *Physics Reports* **499** (2011) 1–101 .
- [6] Clarke, M. and Wilson, A. G., The dynamics of urban spatial structure: the progress of a research programme, *Trans. Inst. Br. Geogr. New Ser.* **10** (1985) 427–451.
- [7] Collar, A., Network theory and religious innovation, *Mediterranean Historical Review* **22** (2007) 149–162.

- [8] Dearden, J. and Wilson, A., Exploring urban retail phase transitions, Technical report, UCL (2009).
- [9] Dearden, J., and Wilson, A., Urban retail phase transitions 1: an analysis system, Technical report, UCL (2010).
- [10] Erlander, S., and Stewart, N.F., *The Gravity Model in Transportation Analysis* (VSP, 1990).
- [11] Evans, T.S., Clique graphs and overlapping communities, *J.Stat.Mech* (2010) P12037.
- [12] Evans, T., Knappett, C., and Rivers, R., Using statistical physics to understand relational space: A case study from mediterranean prehistory, in *Complexity Perspectives on Innovation and Social Change*, eds. Lane, D., Pumain, D., van der Leeuw, S., and West, G., *Methodos Series*, Vol. 7, chapter 17 (Springer, 2009), pp. 451–479.
- [13] Evans, T.S., and Lambiotte, R.. Line Graphs, Link Partitions and Overlapping Communities, *Phys.Rev.E* **80** (2009) 016105.
- [14] Granovetter, M. S., The Strength of Weak Ties, *Am.J.Sociology* **78** (1973) 1360.
- [15] Hage, P. and Harary, F. *Exchange in Oceania: a graph theoretic analysis*. Oxford, Clarendon Press 1991.
- [16] Irwin, G. 1983. Chieftainship, kula and trade in Massim prehistory, in Leach, J.W. and Leach, E. (eds.) *The Kula: new perspectives on Massim exchange*: 29–72. Cambridge: Cambridge University Press.
- [17] Knappett, C., Evans, T., and Rivers, R., Modelling maritime interaction in the aegean bronze age, *Antiquity* **82** (2008) 1009–1024.
- [18] Knappett, C., Evans, T., and Rivers, R., The Theran eruption and Minoan palatial collapse: new interpretations gained from modelling the maritime network, *Antiquity* (in press) (2011).
- [19] Kohler, T. A., Gumerman, G. J., and Reynolds, R. G., Simulating ancient societies, *Scientific American* (2005).
- [20] Ortúzar, J. de D., and Willumsen, L.G. *Modelling Transport*, (Wiley, 1994)
- [21] Poduri, S. and Sukhatme, G. S., Constrained coverage for mobile sensor networks, in *IEEE International Conference on Robotics and Automation* (2004), pp. 165–172.
- [22] Penrose, M., *Random Geometric Graphs* (OUP, 2003).
- [23] Renfrew, C., Trade as action at a distance, in *Ancient Civilization and Trade*, eds. Sabloff, J. and Lamberg-Karlovsky, C. (University of New Mexico Press, Albuquerque), 1975), pp. 3–59.

- [24] Renfrew, C. and Level, E., Exploring dominance: predicting polities from centres, in *Transformations: Mathematical Approaches to Culture Change*, eds. Renfrew, A. and Cooke, K. (Academic Press, London, 1979), pp. 145–67.
- [25] Rihll, T. E. and Wilson, A. G., Spatial interaction and structural models in historical analysis: Some possibilities and an example, *Histoire & Mesure* **2** (1987) 5–32.
- [26] Rihll, T. and Wilson, A., Modelling settlement structures in ancient greece: new approaches to the polis, in *City and country in the ancient world*, eds. Rich, J. and Wallace-Hadrill, A., chapter 3 (1991).
- [27] Rivers, R., Knappett, C., and Evans, T., Network models and archaeological spaces, in *Computational Approaches to Archaeological Spaces*, eds. A. Bevan and Lake, M. (Left Coast Press, 2011).
- [28] Sindbæk, S. M., Networks and nodal points: the emergence of towns in early Viking age Scandinavia, *Antiquity* **81** (2007) 119132.
- [29] Sindbæk, S. M., The small world of the Vikings: Networks in early medieval communication and exchange, *Norwegian Archaeological Review* **40** (2007) 59–74.
- [30] Srinivasa, S. and Haenggi, M., Distance distributions in finite uniformly random networks: Theory and applications, *IEEE Transactions on Vehicular Technology* **59** (2010) 940–949.
- [31] Terrell, J., Human biogeography in the solomon islands, *Fieldiana, Anthropology* **68** (1977).
- [32] Terrell, J. E., Language and material culture on the Sepik coast of Papua New Guinea: Using social network analysis to simulate, graph, identify, and analyze social and cultural boundaries between communities, *The Journal of Island and Coastal Archaeology* **5** (2010) 3–32.
- [33] Wilkinson, T. J., Gibson, M., Christiansen, J. H., Widell, M., Schloen, D., Kouchoukos, N., Woods, C., Sanders, J., Simunich, K.-L., Altaweel, M., Ur, J. A., Hritz, C., Lauinger, J., Paulette, T., and Tenney, J., Modeling settlement systems in a dynamic environment: Case studies from mesopotamia, in *Model Based Archaeology*, eds. Kohler, T. and van der Leeuw, S. E. (Santa Fe, New Mexico: School for Advanced Research Press, 2007), pp. 175–208, ISSN/ISBN: 1-930618-87-9.
- [34] Wilson, A., Boltzmann, Lotka and Volterra and spatial structural evolution: an integrated methodology for some dynamical systems, *J.R.Soc.Interface* **5** (2008) 865–871.
- [35] Wilson, A. and Dearden, J., Phase transitions and path dependence in urban evolution, *Journal of Geographical Systems* (2010) 1–16.

A Supplementary Material

The following tables contain all the information needed to generate the examples in this paper. Simple replacements of text, e.g. using a text editor, should be able to convert the $\text{\LaTeX}$ source into other suitable formats.

Number	Name	Size	Lat	Long
1	Knossos	1	35.3	25.16
2	Malia	1	35.29	25.49
3	Phaistos	1	35.05	24.81
4	Kommos	0.67	35.02	24.76
5	A.Triadha	1	35.06	24.79
6	P-kastro	1	35.2	26.28
7	Zakros	0.67	35.1	26.26
8	Gournia	1	35.11	25.79
9	Chania	1	35.52	24.02
10	Akrotiri	0.67	36.35	25.4
11	Phylakopi	0.67	36.73	24.42
12	Kastri	0.67	36.22	23.06
13	Naxos	1	37.11	25.38
14	Kea	0.67	37.67	24.33
15	Karpathos	0.33	35.42	27.15
16	Rhodes	1	36.42	28.16
17	Kos	0.67	36.88	27.28
18	Miletus	1	37.84	27.24
19	Iasos	0.67	37.28	27.42
20	Samos	0.67	37.66	26.87
21	Petras	1	35.2	26.12
22	Rethymno	1	35.35	24.53
23	Paroikia	0.67	37.08	25.15
24	Amorgos	0.33	36.82	25.15
25	Ios	0.33	36.73	25.29
26	Aegina	0.67	37.75	23.42
27	Mycenae	1	37.73	22.76
28	A.Stephanos	1	36.8	22.58
29	Lavrion	0.67	37.7	24.05
30	Kasos	0.33	35.42	26.91
31	Kalymnos	0.33	36.98	27.02
32	Myndus	0.67	37.05	27.23
33	Cesme	0.67	38.32	26.3
34	Akbuk	0.67	37.41	27.41
35	Menelaion	0.33	37.11	22.37
36	Argos	0.67	37.63	22.73
37	Lerna	0.67	37.5	22.73
38	Asine	0.33	37.56	22.86
39	Eleusis	0.67	38.04	23.54

Table 3: List of the positions and sizes of the 39 MBA Aegean sites.



Figure 5: Exemplary networks in **ariadne** for the 39 MBA Aegean sites of Fig. 1. The distance scale used is $D = 100\text{km}$ with distances based on estimated travel times in which land travel has a friction coefficient of 3 compared to sea travel. Vertices are proportional to the site size (0.33, 0.67 or 1.00 for small, medium and large sites). Top right has $J = -1.00$, $\mu = 0.500$, $\kappa = 1.00$ and $\lambda = 4.00$. Top left has $\lambda = 2.50$ but other values unchanged. This shows the effect of reducing the benefits of interaction on the network. The lower right network has $J = -0.975$, $\mu = 0.500$, $\kappa = 0.90$ and $\lambda = 4.00$. This choice ensures top right and bottom left networks have Hamiltonians with the linear W_i coefficient (this is effectively $-4\kappa + J + \mu$ as here $F_{ii} \approx 0$) but the coefficient of the quadratic term in W_i is lower in the lower network encouraging larger variations in site sizes. The lower left network has exactly the same parameter values of the top right but site 10, Akrotiri on Thera, has been removed to reflect the post-eruption geography.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
1	0	47	180	180	180	135	143	100	130	128	185	228	215	300	205	315	288	335	330	316	110	80	215	200	168	325	393	300	300	175	265	283	365	320	398	366	341	330	347
2	47	0	270	270	270	90	100	54	145	118	185	243	204	294	163	273	260	305	295	300	65	95	204	183	160	330	401	312	308	130	235	251	349	293	410	375	353	341	352
3	180	270	0	0	0	183	173	153	213	293	350	244	380	465	262	408	408	472	461	475	215	135	380	365	333	490	558	308	465	238	400	415	547	457	406	404	383	373	431
4	180	270	0	0	0	183	173	153	213	293	350	244	380	465	262	408	408	472	461	475	215	135	380	365	333	490	558	308	465	238	400	415	547	457	406	404	383	373	431
5	180	270	0	0	0	183	173	153	213	293	350	244	380	465	262	408	408	472	461	475	215	135	380	365	333	490	558	308	465	238	400	415	547	457	406	404	383	373	431
6	135	90	183	183	183	0	17	70	230	155	243	330	232	340	90	215	218	277	270	279	42	180	233	205	195	395	471	395	358	61	212	226	360	268	488	436	414	404	417
7	143	100	173	173	173	17	0	83	246	169	256	339	244	359	96	228	233	288	283	291	48	195	244	217	208	400	484	409	364	68	225	238	375	281	507	447	425	415	427
8	100	54	153	153	153	70	83	0	196	142	219	293	224	327	143	258	252	308	300	303	39	144	225	203	186	368	447	360	332	108	240	252	373	295	458	413	391	380	384
9	130	145	213	213	213	230	246	196	0	156	147	116	214	242	310	390	356	370	385	366	211	60	212	225	177	256	304	189	243	271	328	342	375	385	287	277	254	244	282
10	128	118	293	293	293	155	169	142	156	0	91	211	91	185	200	245	203	216	230	205	144	135	86	79	47	253	334	272	201	169	170	184	245	226	370	303	281	270	252
11	185	185	350	350	350	243	256	219	147	91	0	147	89	107	291	334	279	267	289	242	228	158	84	120	70	148	235	198	112	265	247	261	238	275	295	198	176	165	165
12	228	243	244	244	244	330	339	293	116	211	147	0	233	202	391	457	411	412	434	398	303	161	229	264	208	187	218	87	194	359	376	390	372	429	185	184	162	124	212
13	215	204	380	380	380	232	244	224	214	91	89	233	0	117	234	293	188	185	207	152	226	209	13	67	53	198	305	289	140	247	170	184	162	195	387	270	248	237	202
14	300	294	465	465	465	340	359	327	242	185	107	202	117	0	377	378	291	266	302	235	324	260	119	175	145	90	223	247	28	347	262	281	195	286	340	190	168	157	91
15	205	163	262	262	262	90	96	143	310	200	291	391	234	377	0	147	175	260	250	265	108	253	263	229	232	433	523	455	397	30	193	196	353	251	548	487	465	454	460
16	315	273	408	408	408	215	228	258	390	245	334	457	293	378	147	0	106	189	179	202	228	351	287	244	265	462	559	518	402	162	133	123	298	170	611	528	506	495	482
17	288	260	408	408	408	218	233	252	356	203	279	411	188	291	175	106	0	78	66	94	227	329	197	140	191	367	477	446	314	175	35	17	191	63	539	470	448	437	365
18	335	305	472	472	472	277	288	308	370	216	267	412	185	266	260	189	78	0	70	37	282	346	188	148	215	356	471	459	297	247	69	59	146	50	552	447	425	414	348
19	330	295	461	461	461	270	283	300	385	230	289	434	207	302	250	179	66	70	0	88	276	352	215	176	230	386	501	490	332	240	65	49	197	41	583	453	431	420	384
20	316	300	475	475	475	279	291	303	366	205	242	398	152	235	265	202	94	37	88	0	283	332	163	132	194	335	443	442	275	252	81	76	117	70	535	407	385	374	336
21	110	65	215	215	215	42	48	39	211	144	228	303	226	324	108	228	227	282	276	283	0	157	223	197	185	375	452	372	337	76	216	230	353	272	465	424	402	391	388
22	80	95	135	135	135	180	195	144	60	135	158	161	209	260	253	351	329	346	352	332	157	0	201	206	164	281	341	235	263	220	296	308	368	349	328	313	290	280	309
23	215	204	380	380	380	233	244	225	212	86	84	229	13	119	263	287	197	188	215	163	223	201	0	62	44	210	310	282	146	239	188	203	168	206	375	253	231	220	185
24	200	183	365	365	365	205	217	203	225	79	120	264	67	175	229	244	140	148	176	132	197	206	62	0	67	254	349	315	203	199	124	141	175	164	408	257	235	224	197
25	168	160	333	333	333	195	208	186	177	47	70	208	53	145	232	265	191	215	230	194	185	164	44	67	0	207	302	266	159	204	177	192	211	229	359	259	249	250	212
26	325	330	490	490	490	395	400	368	256	253	148	187	198	90	433	462	367	356	386	335	375	281	210	254	207	0	170	237	77	410	356	371	289	383	330	164	142	131	38
27	393	401	558	558	558	471	484	447	304	334	235	218	305	223	523	559	477	471	501	443	452	341	310	349	302	170	0	268	209	493	467	481	410	492	330	30	60	75	190
28	300	312	308	308	308	395	409	360	189	272	198	87	289	247	455	518	446	459	490	442	372	235	282	315	266	237	268	0	241	424	429	445	423	472	93	233	209	212	246
29	300	308	465	465	465	358	364	332	243	201	112	194	140	28	397	402	314	297	332	275	337	263	146	203	159	77	209	241	0	369	290	309	214	314	334	174	152	141	77
30	175	130	238	238	238	61	68	108	271	169	265	359	247	347	30	162	175	247	240	252	76	220	239	199	204	410	493	424	369	0	179	194	332	236	517	468	446	435	418
31	265	235	400	400	400	212	225	240	328	170	247	376	170	262	193	133	35	69	65	81	216	296	188	124	177	356	467	429	290	179	0	19	175	58	522	426	404	393	353
32	283	251	415	415	415	226	238	252	342	184	261	390	184	281	196	123	17	59	49	76	230	308	203	141	192	371	481	445	309	194	19	0	175	47	538	438	416	405	366
33	365	349	547	547	547	360	375	373	375	245	238	372	162	195	353	298	191	146	197	117	353	368	168	175	211	289	410	423	214	332	175	175	0	184	516	376	354	343	295
34	320	293	457	457	457	268	281	295	385	226	275	429	195	286	251	170	63	50	41	70	272	349	206	164	229	383	492	472	314	236	58	47	184	0	565	459	437	425	387
35	398	410	406	406	406	488	507	458	287	370	295	185	387	340	548	611	539	552	583	535	465	328	375	408	359	330	330	93	334	517	522	538	516	565	0	300	270	327	339
36	366	375	404	404	404	436	447	413	277	303	198	184	270	190	487	528	470	447	453	407	424	313	253	257	259	164	30	233	174	468	426	438	376	459	300	0	30	54	220
37	341	353	383	383	383	414	425	391	254	281	176	162	248	168	465	506	448	425	431	385	402	290	231	235	249	142	60	209	152	446	404	416	354	437	270	30	0	15	158
38	330	341	373	373	373	404	415	380	244	270	165	124	237	157	454	495	437	414	420	374	391	280	220	224	250	131	75	212	141	435	393	405	343	425	327				