The Quarterly Journal of Mathematics Advance Access published on October 4, 2018
Quart. J. Math. 70 (2019), 305–318; doi:10.1093/qmath/hay040

L^P-CAFFARELLI–KOHN–NIRENBERG TYPE INEQUALITIES ON HOMOGENEOUS GROUPS

by TOHRU OZAWA[†]

(*Department of Applied Physics, Waseda University, Tokyo 169-8555, Japan*)

MICHAEL RUZHANSKY[‡]
(*Department of Mathematics, Imperial College London, 180 Queen's Gate, London SW7 2AZ, UK*)

and DURVUDKHAN SURAGAN[§]
(*Department of Mathematics, School of Science and Technology, Nazarbayev University, 53 Kabanbay Batyr Ave, Astana 010000, Kazakhstan*)

[Received 22 November 2016]

Abstract

We prove L^p-Caffarelli–Kohn–Nirenberg type inequalities on homogeneous groups, which is one of most general subclasses of nilpotent Lie groups, all with sharp constants. We also discuss some of their consequences. Already in the Abelian cases of isotropic or anisotropic $\mathbb{R}^n$, our results provide new conclusions in view of the arbitrariness of the choice of the not necessarily Euclidean quasi-norm.

1. Introduction

Consider the following weighted Hardy–Sobolev type inequalities due to Caffarelli–Kohn–Nirenberg [5]: for all $f \in C_0^\infty(\mathbb{R}^n)$, it holds

$$\left(\int_{\mathbb{R}^n} \|x\|^{-p\beta}|f|^p \,\mathrm{d}x\right)^{\frac{2}{p}} \le C_{\alpha,\beta}\int_{\mathbb{R}^n} \|x\|^{-2\alpha}|\nabla f|^2 \,\mathrm{d}x, \tag{1.1}$$

where, for $n \ge 3$

$$-\infty < \alpha < \frac{n-2}{2}, \quad \alpha \le \beta \le \alpha + 1, \quad \text{and} \quad p = \frac{2n}{n-2+2(\beta-\alpha)},$$

and, for $n = 2$

[†]E-mail: txozawa@waseda.jp

[‡]Corresponding author. E-mail: m.ruzhansky@imperial.ac.uk

[§]E-mail: durvudkhan.suragan@nu.edu.kz

© The Author(s) 2018. Published by Oxford University Press. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

$$-\infty < \alpha < 0, \quad \alpha < \beta \leq \alpha + 1, \quad \text{and } p = \frac{2}{\beta - \alpha},$$

and where $\|x\| = \sqrt{x_1^2 + \cdots + x_n^2}$ is the Euclidean norm. Nowadays, there is a lot of literature on Caffarelli–Kohn–Nirenberg type inequalities and their applications. In the case $p = 2$ (see for example [25]), for all $f \in C_0^\infty(\mathbb{R}^n)$, one has

$$\int_{\mathbb{R}^n} \|x\|^{-2(\alpha+1)} |f|^2 \, dx \leq \widetilde{C}_\alpha \int_{\mathbb{R}^n} \|x\|^{-2\alpha} |\nabla f|^2 \, dx, \tag{1.2}$$

with any $n \geq 2$ and $-\infty < \alpha < 0$, which in turn can be presented for any $f \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$ as

$$\left\| \frac{1}{\|x\|^{\alpha+1}} |f| \right\|_{L^2(\mathbb{R}^n)} \leq C_\alpha \left\| \frac{1}{\|x\|^\alpha} |\nabla f| \right\|_{L^2(\mathbb{R}^n)}, \tag{1.3}$$

all $\alpha \in \mathbb{R}$.

Motivating the development of the analysis associated to homogeneous groups in [12], Folland and Stein raised an important question of determining which elements of the classical harmonic analysis do depend only on the group and the dilation structures. The natural setting for this kind of problem is that of homogeneous groups, in particular, including the cases of anisotropic structures on $\mathbb{R}^n$. In this paper, we show that the Caffarelli–Kohn–Nirenberg inequality continues to hold in the setting of homogeneous groups. In particular, it has to also hold on anisotropic $\mathbb{R}^n$, with a number of different consequences. Moreover, the quasi-norm $|\cdot|$ does not need to be Euclidean, but can be an arbitrary homogeneous quasi-norm on $\mathbb{R}^n$.

Recently, a homogeneous group version of the inequality (1.3) was obtained in the work [21], that is, it was proved that if $\mathbb{G}$ is a homogeneous group of homogeneous dimension Q, then, for all $f \in C_0^\infty(\mathbb{G} \setminus \{0\})$ and, for every homogeneous quasi-norm $|\cdot|$ on $\mathbb{G}$, we have

$$\frac{|Q - 2 - 2\alpha|}{2} \left\| \frac{f}{|x|^{\alpha+1}} \right\|_{L^2(\mathbb{G})} \leq \left\| \frac{1}{|x|^\alpha} \mathcal{R} f \right\|_{L^2(\mathbb{G})}, \quad \forall \, \alpha \in \mathbb{R}, \tag{1.4}$$

where $\mathcal{R}$ is defined by (2.4). Note that if $\alpha \neq \frac{Q-2}{2}$, then the constant in (1.4) is sharp for any homogeneous quasi-norm $|\cdot|$ on $\mathbb{G}$. In the abelian case $\mathbb{G} = (\mathbb{R}^n, +)$, we have $Q = n$, $e(x) = x = (x_1, \ldots, x_n)$, so for each $\alpha \in \mathbb{R}$ with $\alpha \neq \frac{n-2}{2}$ and for any homogeneous quasi-norm $|\cdot|$ on $\mathbb{R}^n$, the inequality (1.4) implies a new inequality with the optimal constant:

$$\frac{|n - 2 - 2\alpha|}{2} \left\| \frac{f}{|x|^{\alpha+1}} \right\|_{L^2(\mathbb{R}^n)} \leq \left\| \frac{1}{|x|^\alpha} \frac{x}{|x|} \cdot \nabla f \right\|_{L^2(\mathbb{R}^n)}. \tag{1.5}$$

In turn, by using Schwarz's inequality with the standard Euclidean distance $\|x\| = \sqrt{x_1^2 + \cdots + x_n^2}$, this implies the L^2 Caffarelli–Kohn–Nirenberg inequality [5] for $\mathbb{G} \equiv \mathbb{R}^n$ with the optimal constant:

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

$$\frac{|n-2-2\alpha|}{2}\left\|\frac{f}{\|x\|^{\alpha+1}}\right\|_{L^2(\mathbb{R}^n)}\leq\left\|\frac{1}{\|x\|^{\alpha}}\nabla f\right\|_{L^2(\mathbb{R}^n)},\quad\forall\,\alpha\in\mathbb{R},\tag{1.6}$$

for all $f\in C_0^\infty(\mathbb{R}^n\setminus\{0\})$. Here optimality of the constant $\frac{|n-2-2\alpha|}{2}$ was proved in [6, Theorem 1.1. (ii)]. In addition, we can also note that the analysis in these type inequalities and improvements of their remainder terms has a long history, initiated by Brezis and Nirenberg in [3] and then in [1] for Sobolev inequalities, and in [4] for Hardy inequalities, see also [2], with many subsequent works in this subject.

The L^2-inequality (1.3) in the anisotropic setting as well as in the more abstract setting of homogeneous groups was analyzed in [23] by using an explicit formula for the remainder that is available in the case of L^2-spaces. Such a remainder formula fails in the scale of L^p-spaces for $p\neq 2$ and, therefore, in this paper, we approach these inequalities by a different method.

Thus, the main aim of this paper is to extend the above inequality (1.4) to the general L^p case for all $1<p<\infty$ by a different approach. Here, the first result of this paper is: for each $f\in C_0^\infty(\mathbb{G}\setminus\{0\}), 1<p<\infty$, and any homogeneous quasi-norm $|\cdot|$ on $\mathbb{G}$, we have

$$\frac{|Q-\gamma|}{p}\left\|\frac{f}{|x|^{\frac{\gamma}{p}}}\right\|_{L^p(\mathbb{G})}^{p}\leq\left\|\frac{1}{|x|^{\alpha}}\mathcal{R}f\right\|_{L^p(\mathbb{G})}\left\|\frac{f}{|x|^{\frac{\beta}{p-1}}}\right\|_{L^p(\mathbb{G})}^{p-1},\quad\forall\,\alpha,\quad\beta\in\mathbb{R},\tag{1.7}$$

where $\gamma=\alpha+\beta+1$, and the constant $\dfrac{|Q-\gamma|}{p}$ is sharp if $\gamma\neq Q$. Here $\mathcal{R}$ is the radial derivative operator on $\mathbb{G}$ defined by (2.4).

All above inequalities are generalizations of the classical Hardy inequality, which takes the form

$$\left\|\frac{f(x)}{\|x\|}\right\|_{L^p(\mathbb{R}^n)}\leq\frac{p}{n-p}\,\|\nabla f\|_{L^p(\mathbb{R}^n)},\quad n\geq 2,\quad 1\leq p<n,\tag{1.8}$$

where ∇ is the standard gradient in $\mathbb{R}^n$, $f\in C_0^\infty(\mathbb{R}^n\setminus\{0\}), \|x\|=\sqrt{x_1^2+\cdots+x_n^2}$ and the constant $\frac{p}{n-p}$ is known to be sharp.

We refer to a recent interesting paper of Hoffmann-Ostenhof and Laptev [14] on this subject for inequalities with weights, to [13] for many-particle versions, to Ekholm–Kovařík–Laptev [10] for p-Laplacian interpretations, and to many further references therein and otherwise. We also refer to more recent preprints [18–23] and references therein for the story behind Hardy-type inequalities on nilpotent Lie groups. Moreover, we also note that we do not work with the 'horizontal' gradients since there may not be any for general homogeneous groups. If the group is stratified, also the horizontal versions are possible, and we refer to [24] for this subject.

Before giving preliminaries for stating our results, let us mention another observation that the Hardy inequality (1.8) can be sharpened to the inequality

$$\left\|\frac{f(x)}{\|x\|}\right\|_{L^p(\mathbb{R}^n)}\leq\frac{p}{n-p}\,\|\mathcal{R}f\|_{L^p(\mathbb{R}^n)},\quad n\geq 2,\quad 1\leq p<n.\tag{1.9}$$

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

Here $\mathcal{R}$ is the radial derivative operator on $\mathbb{G}$ defined by (2.4). It is clear that (1.9) implies (1.8) since the function $\frac{x}{\|x\|}$ is bounded. The remainder terms for (1.9) have been analyzed by Ioku–Ishiwata–Ozawa [**15**], see also Machihara–Ozawa–Wadade [**17**].

One of the results in [**21**] was that if $\mathbb{G}$ is any homogeneous group and $|\cdot|$ is a homogeneous quasi-norm on $\mathbb{G}$, as an analogue of (1.9), we obtain the following generalized L^p-Hardy inequality:

$$\left\| \frac{f}{|x|} \right\|_{L^p(\mathbb{G})} \leq \frac{p}{Q-p} \|\mathcal{R}f\|_{L^p(\mathbb{G})}, \quad 1 < p < Q, \tag{1.10}$$

for all $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$. Here $\mathcal{R}$ is the radial derivative operator on $\mathbb{G}$ defined by (2.4). Let A be the n-diagonal matrix

$$A = \operatorname{diag}(\nu_1, \ldots, \nu_n), \tag{1.11}$$

where ν_k is the homogeneous degree of X_k, and

$$Q = \operatorname{Tr} A = \nu_1 + \cdots + \nu_n$$

is the homogeneous dimension of $\mathbb{G}$. We note that the exponential mapping $\exp_{\mathbb{G}}:\mathfrak{g} \to \mathbb{G}$ is a global diffeomorphism and the vector $e(x) = (e_1(x), \ldots, e_n(x))$ is the decomposition of its inverse $\exp_{\mathbb{G}}^{-1}$ with respect to the basis $\{X_1, \ldots, X_n\}$, namely $e(x)$ is determined by

$$\exp_{\mathbb{G}}^{-1}(x) = e(x) \cdot \nabla \equiv \sum_{j=1}^{n} e_j(x) X_j.$$

For $p = n$ or $p = Q$, the inequalities (1.8) and (1.10) fail for any constant. The critical versions of (1.9) with $p = n$ were investigated by Ioku–Ishiwata–Ozawa [**16**]. Their generalizations as well as a number of other critical (logarithmic) Hardy inequalities on homogeneous groups were obtained in recent works [**18**, **21**, **22**]. Here, we only mention a related family of logarithmic Hardy inequalities

$$\sup_{R>0} \left\| \frac{f - f_R}{|x|^{\frac{Q}{p}} \log \frac{R}{|x|}} \right\|_{L^p(\mathbb{G})} \leq \frac{p}{p-1} \left\| \frac{1}{|x|^{\frac{Q}{p}-1}} \mathcal{R}f \right\|_{L^p(\mathbb{G})}, \tag{1.12}$$

for all $1 < p < \infty$, where $f_R = f(R\frac{x}{|x|})$. We refer to [**18**] for further explanations and extensions but only mention here that for $p = Q$, the inequality (1.12) gives a critical case of Hardy's inequalities (1.10).

In Section 2, we give some necessary tools on homogeneous groups and fix the notation. In Section 3, we present L^p-Caffarelli–Kohn–Nirenberg type inequalities on the homogeneous group $\mathbb{G}$ and then discuss their consequences and proofs. In Section 4, we discuss higher order cases.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

2. Preliminaries

In this standard preliminary section, we very shortly recall some basics details of homogeneous groups. The general analysis on homogeneous groups was developed by Folland and Stein in their book [12], and we also refer for more recent developments to the monograph [11] by Véronique Fischer and the second named author.

It is known that a dilation family of a Lie algebra $\mathfrak{g}$ has the following representation:

$$D_\lambda = \operatorname{Exp}(A \ln \lambda) = \sum_{k=0}^{\infty} \frac{1}{k!} (\ln(\lambda) A)^k ,$$

where A is a diagonalizable positive linear operator on $\mathfrak{g}$, and each family of linear mappings D_λ is a morphism of $\mathfrak{g}$, that is a linear mapping from $\mathfrak{g}$ to itself with the property

$$\forall X, Y \in \mathfrak{g}, \quad \lambda > 0, \quad [D_\lambda X, D_\lambda Y] = D_\lambda [X, Y],$$

where $[X, Y] := XY - YX$ is the Lie bracket. Shortly, a *homogeneous group* is a connected and simply connected Lie group whose Lie algebra is equipped with dilations. It induces the dilation structure on $\mathbb{G}$ which we continue to denote by $D_\lambda(x)$ or simply by λx. We denote by

$$Q := \operatorname{Tr} A$$

the homogeneous dimension of $\mathbb{G}$. We also recall that the standard Lebesque measure on $\mathbb{R}^N$ is the Haar measure for $\mathbb{G}$ (see, for example [11, Proposition 1.6.6]).

Let us fix a basis $\{X_1, \ldots, X_n\}$ of the Lie algebra $\mathfrak{g}$ of the homogeneous group $\mathbb{G}$ such that

$$AX_k = \nu_k X_k$$

for each $1 \leq k \leq n$, so that A can be taken to be

$$A = \operatorname{diag}(\nu_1, \ldots, \nu_n). \tag{2.1}$$

Then each X_k is homogeneous of degree ν_k and also

$$Q = \nu_1 + \cdots + \nu_n, \tag{2.2}$$

which is called a homogeneous dimension of $\mathbb{G}$. Since homogeneous groups are nilpotent, the exponential mapping $\exp_{\mathbb{G}}: \mathfrak{g} \to \mathbb{G}$ is a global diffeomorphism. In addition, the decomposition of $\exp_{\mathbb{G}}^{-1}(x)$ in the Lie algebra $\mathfrak{g}$ defines the vector

$$e(x) = (e_1(x), \ldots, e_n(x))$$

by the formula

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

$$\exp_{\mathbb{G}}^{-1}(x) = e(x) \cdot \nabla \equiv \sum_{j=1}^{n} e_j(x) X_j,$$

where $\nabla = (X_1, \ldots, X_n)$. Alternatively, this means the equality

$$x = \exp_{\mathbb{G}}(e_1(x)X_1 + \cdots + e_n(x)X_n).$$

Using homogeneity, we have

$$rx := D_r(x) = \exp_{\mathbb{G}}(r^{\nu_1} e_1(x)X_1 + \cdots + r^{\nu_n} e_n(x)X_n),$$

that is,

$$e(rx) = (r^{\nu_1} e_1(x), \ldots, r^{\nu_n} e_n(x)).$$

Thus, since $r > 0$ is arbitrary, without loss of generality taking $|x| = 1$, we obtain

$$\frac{d}{\mathrm{d}|rx|} f(rx) = \frac{d}{\mathrm{d}r} f(\exp_{\mathbb{G}}(r^{\nu_1} e_1(x)X_1 + \cdots + r^{\nu_n} e_n(x)X_n)). \tag{2.3}$$

Denoting by

$$\mathcal{R} := \frac{d}{\mathrm{d}r}, \tag{2.4}$$

for all $x \in \mathbb{G}$, this gives the equality

$$\frac{d}{\mathrm{d}|x|} f(x) = \mathcal{R} f(x), \tag{2.5}$$

for each homogeneous quasi-norm $|x|$ on a homogeneous group $\mathbb{G}$. That is, the operator $\mathcal{R}$ plays the role of the radial derivative on $\mathbb{G}$. It follows from (2.4) that $\mathcal{R}$ is homogeneous of order -1. It is known that every homogeneous group $\mathbb{G}$ admits a homogeneous quasi-norm $|\cdot|$. The $|\cdot|$-ball centred at $x \in \mathbb{G}$ with radius $R > 0$ can be defined by

$$B(x, R) := \{y \in \mathbb{G}\colon |x^{-1}y| < R\}.$$

The following polar decomposition was established in [**12**] (see also [**11**, Section 3.1.7]).

PROPOSITION 2.1 *Let $\mathbb{G}$ be a homogeneous group equipped with a homogeneous quasi-norm $|\cdot|$. Then there is a (unique) positive Borel measure σ on the unit quasi sphere*

$$\mathfrak{S} := \{x \in \mathbb{G}\colon\ |x| = 1\}, \tag{2.6}$$

such that for all $f \in L^1(\mathbb{G})$, we have

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

$$\int_{\mathbb{G}} f(x)\,\mathrm{d}x = \int_0^\infty \int_{\mathfrak{S}} f(ry)\, r^{Q-1}\,\mathrm{d}\sigma(y)\,\mathrm{d}r. \tag{2.7}$$

3. L^p-Caffarelli–Kohn-Nirenberg type inequalities and consequences

In this section and in the sequel, we adopt all the notation introduced in Section 2 concerning homogeneous groups and the operator $\mathcal{R}$. We formulate the following L^p-Caffarelli–Kohn–Nirenberg type inequalities on the homogeneous group $\mathbb{G}$ and then discuss their consequences and proofs.

THEOREM 3.1 *Let $\mathbb{G}$ be a homogeneous group of homogeneous dimension Q and let $\alpha,\ \beta \in \mathbb{R}$. Then for all complex-valued functions $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$, $1 < p < \infty$ and any homogeneous quasi-norm $|\cdot|$ on $\mathbb{G}$, we have*

$$\frac{|Q-\gamma|}{p}\left\|\frac{f}{|x|^{\frac{\gamma}{p}}}\right\|^p_{L^p(\mathbb{G})} \le \left\|\frac{1}{|x|^\alpha}\mathcal{R}f\right\|_{L^p(\mathbb{G})}\left\|\frac{f}{|x|^{\frac{\beta}{p-1}}}\right\|^{p-1}_{L^p(\mathbb{G})}, \tag{3.1}$$

where $\gamma = \alpha+\beta+1$. If $\gamma \neq Q$, then the constant $\dfrac{|Q-\gamma|}{p}$ is sharp.

In the abelian case $\mathbb{G} = (\mathbb{R}^n,\ +)$, we have $Q = n$, $e(x) = x = (x_1,\ldots,x_n)$, so for any homogeneous quasi-norm $|\cdot|$ on $\mathbb{R}^n$ (3.1) implies a new inequality with the optimal constant

$$\frac{|n-\gamma|}{p}\left\|\frac{f}{|x|^{\frac{\gamma}{p}}}\right\|^p_{L^p(\mathbb{R}^n)} \le \left\|\frac{1}{|x|^\alpha}\frac{df}{d|x|}\right\|_{L^p(\mathbb{R}^n)}\left\|\frac{f}{|x|^{\frac{\beta}{p-1}}}\right\|^{p-1}_{L^p(\mathbb{R}^n)}, \tag{3.2}$$

which in turn, by using Schwarz's inequality with the standard Euclidean distance $\|x\| = \sqrt{x_1^2+\cdots+x_n^2}$, implies the L^p-Caffarelli–Kohn–Nirenberg type inequality (see [8, 9]) for $\mathbb{G} \equiv \mathbb{R}^n$ with the sharp constant

$$\frac{|n-\gamma|}{p}\left\|\frac{f}{\|x\|^{\frac{\gamma}{p}}}\right\|^p_{L^p(\mathbb{R}^n)} \le \left\|\frac{1}{\|x\|^\alpha}\nabla f\right\|_{L^p(\mathbb{R}^n)}\left\|\frac{f}{\|x\|^{\frac{\beta}{p-1}}}\right\|^{p-1}_{L^p(\mathbb{R}^n)}, \tag{3.3}$$

for all $f \in C_0^\infty(\mathbb{R}^n\setminus\{0\})$.

When $\alpha = 0,\ \beta = p-1$ and $1 < p < Q$, the inequality (3.1) gives the homogeneous group version of L^p-Hardy inequality

$$\left\|\frac{1}{|x|}f\right\|_{L^p(\mathbb{G})} \le \frac{p}{Q-p}\,\|\mathcal{R}f\|_{L^p(\mathbb{G})}, \tag{3.4}$$

again with $\frac{p}{Q-p}$ being the best constant (see [18–24] for weighted, critical, higher order cases, horizontal cases and their applications in different settings). Note that in comparison with stratified (Carnot) group versions, here the constant is best for any homogeneous quasi-norm $|\cdot|$.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

In the abelian case $\mathbb{G} = (\mathbb{R}^n, +)$, $n \geq 3$, we have $Q = n$, $e(x) = x = (x_1, \ldots, x_n)$, so for any quasi-norm $|\cdot|$ on $\mathbb{R}^n$, (3.4) implies the new inequality

$$\left\| \frac{f}{|x|} \right\|_{L^p(\mathbb{R}^n)} \leq \frac{p}{n-p} \left\| \frac{df}{d|x|} \right\|_{L^p(\mathbb{R}^n)}. \tag{3.5}$$

In turn, by using Schwarz's inequality with the standard Euclidean distance $\|x\| = \sqrt{x_1^2 + \cdots + x_n^2}$, it implies the classical Hardy inequality for $\mathbb{G} \equiv \mathbb{R}^n$

$$\left\| \frac{f}{\|x\|} \right\|_{L^p(\mathbb{R}^n)} \leq \frac{p}{n-p} \|\nabla f\|_{L^p(\mathbb{R}^n)},$$

for all $f \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$. When $|x| \equiv \|x\|$ is the Euclidean distance the remainder terms for (3.5), that is, the exact formulae of the difference between the right-hand side and the left-hand side of the inequality have been analyzed by Ioku–Ishiwata–Ozawa [**15**], see also Machihara–Ozawa–Wadade [**17**] as well as [**16**].

The inequality (3.4) also implies the following Heisenberg–Pauli–Weyl type uncertainly principle on homogeneous groups (see for example [**7**, **19**, **20**] for versions of abelian and stratified groups): for each $f \in C_0^\infty(\mathbb{G} \setminus \{0\})$ and any homogeneous quasi-norm $|\cdot|$ on $\mathbb{G}$, using Hölder's inequality and (3.4), we have

$$\begin{aligned} \|f\|_{L^2(\mathbb{G})}^2 &\leq \left\| \frac{1}{|x|} f \right\|_{L^p(\mathbb{G})} \||x|f\|_{L^{\frac{p}{p-1}}(\mathbb{G})} \\ &\leq \frac{p}{Q-p} \|\mathcal{R}f\|_{L^p(\mathbb{G})} \||x|f\|_{L^{\frac{p}{p-1}}(\mathbb{G})}, \quad 1 < p < Q, \end{aligned} \tag{3.6}$$

that is,

$$\|f\|_{L^2(\mathbb{G})}^2 \leq \frac{p}{Q-p} \|\mathcal{R}f\|_{L^p(\mathbb{G})} \||x|f\|_{L^{\frac{p}{p-1}}(\mathbb{G})}, \quad 1 < p < Q. \tag{3.7}$$

In the abelian case $\mathbb{G} = (\mathbb{R}^n, +)$, taking $Q = n$ and $e(x) = x$, we obtain that (3.7) with $p = 2$ implies the uncertainly principle with any quasi-norm $|x|$

$$\left(\int_{\mathbb{R}^n} |u(x)|^2 \, \mathrm{d}x \right)^2 \leq \left(\frac{2}{n-2} \right)^2 \int_{\mathbb{R}^n} \left| \frac{\mathrm{d}u(x)}{\mathrm{d}|x|} \right|^2 \mathrm{d}x \int_{\mathbb{R}^n} |x|^2 |u(x)|^2 \, \mathrm{d}x. \tag{3.8}$$

In turn, it implies the classical uncertainty principle for $\mathbb{G} \equiv \mathbb{R}^n$ with the standard Euclidean distance $\|x\|$

$$\left(\int_{\mathbb{R}^n} |u(x)|^2 \, \mathrm{d}x \right)^2 \leq \left(\frac{2}{n-2} \right)^2 \int_{\mathbb{R}^n} |\nabla u(x)|^2 \, \mathrm{d}x \int_{\mathbb{R}^n} \|x\|^2 |u(x)|^2 \, \mathrm{d}x,$$

which is the Heisenberg–Pauli–Weyl uncertainly principle on $\mathbb{R}^n$.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

On the other hand, directly from the inequality (3.1), we can obtain a number of Heisenberg–Pauli–Weyl type uncertainly inequities which have various consequences and applications. For example, when $\alpha p = \alpha + \beta + 1$, we have

$$\frac{|Q - \alpha p|}{p} \left\| \frac{f}{|x|^\alpha} \right\|_{L^p(\mathbb{G})}^p \leq \left\| \frac{\mathcal{R} f}{|x|^\alpha} \right\|_{L^p(\mathbb{G})} \left\| |x|^{\frac{1}{p-1}} \frac{f}{|x|^\alpha} \right\|_{L^p(\mathbb{G})}^{p-1}, \tag{3.9}$$

and, on the other hand, if $0 = \alpha + \beta + 1$ and $\alpha = -p$, then

$$\frac{Q}{p} \|f\|_{L^p(\mathbb{G})}^p \leq \| |x|^p \mathcal{R} f \|_{L^p(\mathbb{G})} \left\| \frac{f}{|x|} \right\|_{L^p(\mathbb{G})}^{p-1}, \tag{3.10}$$

all with sharp constants.

Proof of Theorem 3.1. We may assume that $\gamma \neq Q$ since for $\gamma = Q$, there is nothing to prove. Introducing polar coordinates $(r, y) = \left(|x|, \frac{x}{|x|} \right) \in (0, \infty) \times \mathfrak{S}$ on $\mathbb{G}$, where $\mathfrak{S}$ is the pseudo-sphere in (2.6), and using Proposition 2.1, one calculates

$$\begin{aligned}
\int_{\mathbb{G}} \frac{|f(x)|^p}{|x|^\gamma} \, \mathrm{d}x &= \int_0^\infty \int_{\mathfrak{S}} \frac{|f(ry)|^p}{r^\gamma} r^{Q-1} \, \mathrm{d}\sigma(y) \, \mathrm{d}r \\
&= \frac{1}{Q - \gamma} \int_0^\infty \int_{\mathfrak{S}} |f(ry)|^p \frac{\mathrm{d}r^{Q-\gamma}}{\mathrm{d}r} \, \mathrm{d}\sigma(y) \, \mathrm{d}r \\
&= -\frac{1}{Q - \gamma} \operatorname{Re} \int_0^\infty \int_{\mathfrak{S}} p f(ry) |f(ry)|^{p-2} \left(\overline{\frac{\mathrm{d}f(ry)}{\mathrm{d}r}} \right) \frac{1}{r^{\gamma-1}} r^{Q-1} \, \mathrm{d}\sigma(y) \, \mathrm{d}r \\
&= -\frac{p}{Q - \gamma} \operatorname{Re} \int_{\mathbb{G}} f(x) \frac{|f(x)|^{p-2}}{|x|^{\gamma-1}} \left(\overline{\frac{d}{\mathrm{d}|x|} f(x)} \right) \mathrm{d}x \\
&\leq \left| \frac{p}{Q - \gamma} \right| \int_{\mathbb{G}} \frac{|f(x)|^{p-1}}{|x|^{\gamma-1}} \left| \frac{d}{d|x|} f(x) \right| \mathrm{d}x \\
&= \left| \frac{p}{Q - \gamma} \right| \int_{\mathbb{G}} \frac{|f(x)|^{p-1}}{|x|^{\alpha+\beta}} |\mathcal{R} f(x)| \, \mathrm{d}x \\
&\leq \left| \frac{p}{Q - \gamma} \right| \left(\int_{\mathbb{G}} \frac{|\mathcal{R} f(x)|^p}{|x|^{\alpha p}} \, \mathrm{d}x \right)^{\frac{1}{p}} \left(\int_{\mathbb{G}} \frac{|f(x)|^p}{|x|^{\frac{\beta p}{p-1}}} \, \mathrm{d}x \right)^{\frac{p-1}{p}},
\end{aligned}$$

where we have used Hölder's inequality. Thus, we arrive at

$$\left| \frac{Q - \gamma}{p} \right| \int_{\mathbb{G}} \frac{|f(x)|^p}{|x|^\gamma} \, \mathrm{d}x \leq \left(\int_{\mathbb{G}} \frac{|\mathcal{R} f(x)|^p}{|x|^{\alpha p}} \, \mathrm{d}x \right)^{\frac{1}{p}} \left(\int_{\mathbb{G}} \frac{|f(x)|^p}{|x|^{\frac{\beta p}{p-1}}} \, \mathrm{d}x \right)^{\frac{p-1}{p}}. \tag{3.11}$$

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

Now let us show the sharpness of the constant. We need to examine the equality condition in above Hölder's inequality as in the Euclidean case (see [9]). Consider the function

$$g(x) = \begin{cases} e^{-\frac{C}{\lambda}|x|^{\lambda}}, & \lambda := \alpha - \dfrac{\beta}{p-1} + 1 \neq 0, \\ \dfrac{1}{|x|^{C}}, & \alpha - \dfrac{\beta}{p-1} + 1 = 0, \end{cases} \tag{3.12}$$

where $C = \left|\frac{Q-\gamma}{p}\right|$ and $\gamma \neq Q$. Then, it can be checked that

$$\left|\frac{p}{Q-\gamma}\right|^{p} \frac{|\mathcal{R}g(x)|^{p}}{|x|^{\alpha p}} = \frac{|g(x)|^{p}}{|x|^{\frac{\beta p}{p-1}}}, \tag{3.13}$$

which satisfies the equality condition in Hölder's inequality. This shows that the constant $C = \left|\frac{Q-\gamma}{p}\right|$ is sharp. □

4. Higher order cases

In this section, we shortly discuss that by iterating the established L^p-Caffarelli–Kohn–Nirenberg type inequalities, one can get inequalities of higher order. To start let us consider in (3.1), the case

$$\beta = \gamma\left(1 - \frac{1}{p}\right),$$

that is, taking $\beta = (\alpha+1)(p-1)$, the inequality (3.1) implies that $\gamma = p(\alpha+1)$ and

$$\left\| \frac{f}{|x|^{\alpha+1}} \right\|_{L^p(\mathbb{G})} \leq \frac{p}{|Q - p(\alpha+1)|} \left\| \frac{1}{|x|^{\alpha}} \mathcal{R} f \right\|_{L^p(\mathbb{G})}, \quad 1 < p < \infty, \tag{4.1}$$

for any $f \in C_0^{\infty}(\mathbb{G}\setminus\{0\})$ and all $\alpha \in \mathbb{R}$ with $\alpha \neq \frac{Q}{p} - 1$.

Now putting $\mathcal{R}f$ instead of f and $\alpha - 1$ instead of α in (4.1), we consequently have

$$\left\| \frac{\mathcal{R}f}{|x|^{\alpha}} \right\|_{L^p(\mathbb{G})} \leq \frac{p}{|Q - p\alpha|} \left\| \frac{1}{|x|^{\alpha-1}} \mathcal{R}^2 f \right\|_{L^p(\mathbb{G})},$$

for $\alpha \neq \frac{Q}{p}$. Combining it with (4.1), we get

$$\left\| \frac{f}{|x|^{\alpha+1}} \right\|_{L^p(\mathbb{G})} \leq \frac{p}{|Q - p(\alpha+1)|} \frac{p}{|Q - p\alpha|} \left\| \frac{1}{|x|^{\alpha-1}} \mathcal{R}^2 f \right\|_{L^p(\mathbb{G})}, \tag{4.2}$$

for each $\alpha \in \mathbb{R}$ such that $\alpha \neq \frac{Q}{p} - 1$ and $\alpha \neq \frac{Q}{p}$. This iteration process gives

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

$$\left\| \frac{f}{|x|^{\theta+1}} \right\|_{L^p(\mathbb{G})} \leq A_{\theta,k} \left\| \frac{1}{|x|^{\theta+1-k}} \mathcal{R}^k f \right\|_{L^p(\mathbb{G})}, \quad 1 < p < \infty, \tag{4.3}$$

for any $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$ and all $\theta \in \mathbb{R}$ such that $\prod_{j=0}^{k-1} |Q - p(\theta+1-j)| \neq 0$, and

$$A_{\theta,k} := p^k \left[\prod_{j=0}^{k-1} |Q - p(\theta+1-j)| \right]^{-1}.$$

Similarly, we have

$$\left\| \frac{\mathcal{R}f}{|x|^{\vartheta+1}} \right\|_{L^p(\mathbb{G})} \leq A_{\vartheta,m} \left\| \frac{1}{|x|^{\vartheta+1-m}} \mathcal{R}^{m+1} f \right\|_{L^p(\mathbb{G})}, \quad 1 < p < \infty, \tag{4.4}$$

for any $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$ and all $\vartheta \in \mathbb{R}$ such that $\prod_{j=0}^{m-1} |Q - p(\vartheta+1-j)| \neq 0$, and

$$A_{\vartheta,m} := p^m \left[\prod_{j=0}^{m-1} |Q - p(\vartheta+1-j)| \right]^{-1}.$$

Now putting $\vartheta + 1 = \alpha$ and $\theta + 1 = \frac{\beta}{p-1}$ into (4.4) and (4.3), respectively, from (3.1), we obtain

PROPOSITION 4.1 *Let $1 < p < \infty$. For any $k, m \in \mathbb{N}$, we have*

$$\frac{|Q-\gamma|}{p} \left\| \frac{f}{|x|^{\frac{\gamma}{p}}} \right\|_{L^p(\mathbb{G})}^p \leq \widetilde{A}_{\alpha,m} \widetilde{A}_{\beta,k} \left\| \frac{1}{|x|^{\alpha-m}} \mathcal{R}^{m+1} f \right\|_{L^p(\mathbb{G})} \left\| \frac{1}{|x|^{\frac{\beta}{p-1}-k}} \mathcal{R}^k f \right\|_{L^p(\mathbb{G})}^{p-1}, \tag{4.5}$$

for any complex-valued function $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$, $\gamma = \alpha + \beta + 1$, and $\alpha \in \mathbb{R}$ such that $\prod_{j=0}^{m-1} |Q - p(\alpha - j)| \neq 0$, and

$$\widetilde{A}_{\alpha,m} := p^m \left[\prod_{j=0}^{m-1} |Q - p(\alpha - j)| \right]^{-1},$$

as well as $\beta \in \mathbb{R}$ such that $\prod_{j=0}^{k-1} \left| Q - p\left(\frac{\beta}{p-1} - j \right) \right| \neq 0$, and

$$\widetilde{A}_{\beta,k} := p^{k(p-1)} \left[\prod_{j=0}^{k-1} \left| Q - p\left(\frac{\beta}{p-1} - j \right) \right| \right]^{-(p-1)}.$$

We also highlight the case $p = 2$. In this case, an interesting feature is that when we have the exact formula for the remainder which yields the sharpness of the constants as well. We first recall the following estimate and formula.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

THEOREM 4.2 ([21]). *Let* $Q \geq 3$, $\alpha \in \mathbb{R}$ *and* $k \in \mathbb{N}$ *be such that we have* $\prod_{j=0}^{k-1} \left| \frac{Q-2}{2} - (\alpha + j) \right| \neq 0$. *Then for all complex-valued functions* $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$, *we have*

$$\left\| \frac{f}{|x|^{k+\alpha}} \right\|_{L^2(\mathbb{G})} \leq \left[\prod_{j=0}^{k-1} \left| \frac{Q-2}{2} - (\alpha + j) \right| \right]^{-1} \left\| \frac{1}{|x|^\alpha} \mathcal{R}^k f \right\|_{L^2(\mathbb{G})}, \tag{4.6}$$

where the constant above is sharp, and is attained if and only if $f = 0$.

Moreover, for all $k \in \mathbb{N}$ *and* $\alpha \in \mathbb{R}$, *the following equality holds*:

$$\begin{aligned} \left\| \frac{1}{|x|^\alpha} \mathcal{R}^k f \right\|_{L^2(\mathbb{G})}^2 &= \left[\prod_{j=0}^{k-1} \left(\frac{Q-2}{2} - (\alpha + j) \right)^2 \right] \left\| \frac{f}{|x|^{k+\alpha}} \right\|_{L^2(\mathbb{G})}^2 \\ &\quad + \sum_{l=1}^{k-1} \left[\prod_{j=0}^{l-1} \left(\frac{Q-2}{2} - (\alpha + j) \right)^2 \right] \left\| \frac{1}{|x|^{l+\alpha}} \mathcal{R}^{k-l} f \right. \\ &\quad \left. + \frac{Q - 2(l+1+\alpha)}{2|x|^{l+1+\alpha}} \mathcal{R}^{k-l-1} f \right\|_{L^2(\mathbb{G})}^2 \\ &\quad + \left\| \frac{1}{|x|^\alpha} \mathcal{R}^k f + \frac{Q-2-2\alpha}{2|x|^{1+\alpha}} \mathcal{R}^{k-1} f \right\|_{L^2(\mathbb{G})}^2. \end{aligned} \tag{4.7}$$

When $p = 2$, Theorem 3.1 can be restated that for each $f \in C_0^\infty(\mathbb{G}\setminus\{0\})$, and any homogeneous quasi-norm $|\cdot|$ on $\mathbb{G}$, we have

$$\frac{|Q-\gamma|}{2} \left\| \frac{f}{|x|^{\frac{\gamma}{2}}} \right\|_{L^2(\mathbb{G})}^2 \leq \left\| \frac{1}{|x|^\alpha} \mathcal{R} f \right\|_{L^2(\mathbb{G})} \left\| \frac{f}{|x|^\beta} \right\|_{L^2(\mathbb{G})}, \quad \forall\, \alpha,\ \beta \in \mathbb{R}, \tag{4.8}$$

where $\gamma = \alpha + \beta + 1$. Combining (4.8) with (4.6) (or (4.7)), one can obtain a number of inequalities with sharp constants, for example

$$\frac{|Q-\gamma|}{2} \left\| \frac{f}{|x|^{\frac{\gamma}{2}}} \right\|_{L^2(\mathbb{G})}^2 \leq C_j(\beta, k) \left\| \frac{1}{|x|^\alpha} \mathcal{R} f \right\|_{L^2(\mathbb{G})} \left\| \frac{1}{|x|^{\beta-k}} \mathcal{R}^k f \right\|_{L^2(\mathbb{G})}, \tag{4.9}$$

for $\gamma = \alpha + \beta + 1$ and all $\alpha,\ \beta \in \mathbb{R}$ and $k \in \mathbb{N}$, such that,

$$C_j(\beta, k) := \left[\prod_{j=0}^{k-1} \left| \frac{Q-2}{2} - (\beta - k + j) \right| \right]^{-1} \neq 0,$$

as well as

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

$$\frac{|Q-\gamma|}{2}\left\|\frac{f}{|x|^{\frac{\gamma}{2}}}\right\|^2_{L^2(\mathbb{G})} \leq C_j(\alpha,k)\left\|\frac{1}{|x|^{\alpha-k}}\mathcal{R}^{k+1}f\right\|_{L^2(\mathbb{G})}\left\|\frac{f}{|x|^{\beta}}\right\|_{L^2(\mathbb{G})}, \tag{4.10}$$

for $\gamma=\alpha+\beta+1$ and all $\alpha,\ \beta\in\mathbb{R}$ and $k\in\mathbb{N}$, such that,

$$C_j(\alpha,k) := \left[\prod_{j=0}^{k-1}\left|\frac{Q-2}{2}-(\alpha+k+j)\right|\right]^{-1} \neq 0.$$

It follows from (4.7) that these constants $C_j(\beta,k)$ and $C_j(\alpha,k)$ in (4.9) and (4.10) are sharp.

Funding

The second and third authors were supported in parts by the EPSRC Grants EP/K039407/1 and EP/R003025/1, and by the Leverhulme Grants RPG-2014-02 and RPG-2017-151. No new data were collected or generated during the course of research.

References

1. H. Brezis and E. Lieb, Sobolev inequalities with remainder terms, *J. Funct. Anal.* **62** (1985), 73–86.
2. H. Brezis and M. Marcus, Hardy's inequalities revisited, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **25** (1997), 217–237.
3. H. Brézis and L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Commun. Pure Appl. Math.* **36** (1983), 437–477.
4. H. Brezis and J. Vázquez, Blow-up solutions of some nonlinear elliptic problems, *Rev. Mat. Univ. Complut. Madrid* **10** (1997), 443–469.
5. L. Caffarelli, R. Kohn and L. Nirenberg, First order interpolation inequalities with weights, *Compos. Math.* **53** (1984), 259–275.
6. F. Catrina and Z. Q. Wang, On the Caffarelli–Kohn–Nirenberg inequalities: sharp constants, existence (and non existence), and symmetry of extremals functions, *Commun. Pure Appl. Math.* **54** (2001), 229–258.
7. P. Ciatti, M. G. Cowling and F. Ricci, Hardy and uncertainty inequalities on stratified Lie groups, *Adv. Math.* **227** (2015), 365–387.
8. D. G. Costa, Some new and short proofs for a class of Caffarelli–Kohn–Nirenberg type inequalities, *J. Math. Anal. Appl.* **337** (2008), 311–317.
9. Y. Di, L. Jiang, S. Shen and Y. Jin, A note on a class of Hardy–Rellich type inequalities, *J. Inequal. Appl.* **84** (2013), 1–6.
10. T. Ekholm, H. Kovařík and A. Laptev, Hardy inequalities for p-Laplacians with Robin boundary conditions, *Nonlinear Anal.* **128** (2015), 365–379.
11. V. Fischer and M. Ruzhansky, Quantization on Nilpotent Lie Groups*, Vol 314,* Progress in Mathematics, Birkhäuser, Basel, 2016.
12. G. B. Folland and E. M. Stein, Hardy Spaces on Homogeneous Groups*, Vol. 28,* Mathematical Notes, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1982.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019

13. M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof, A. Laptev and J. Tidblom, Many-particle Hardy inequalities, *J. Lond. Math. Soc. (2)* **77** (2008), 99–114.
14. T. Hoffmann-Ostenhof and A. Laptev, Hardy inequalities with homogeneous weights, *J. Funct. Anal.* **268** (2015), 3278–3289.
15. N. Ioku, M. Ishiwata and T. Ozawa, Hardy type inequalities in L^p with sharp remainders, *J. Inequal. Appl.* **2017** (2017), 5.
16. N. Ioku, M. Ishiwata and T. Ozawa, Sharp remainder of a critical Hardy inequality, *Arch. Math. (Basel)* **106** (2016), 65–71.
17. S. Machihara, T. Ozawa and H. Wadade, Remarks on the Hardy type inequalities with remainder terms in the framework of equalities. *To appear in Adv. Studies Pure Math*., 2018.
18. M. Ruzhansky and D. Suragan, Critical Hardy inequalities. arXiv: 1602.04809, 2016.
19. M. Ruzhansky and D. Suragan, Layer potentials, Kac's problem, and refined Hardy inequality on homogeneous Carnot groups, *Adv. Math.* **308** (2017), 483–528.
20. M. Ruzhansky and D. Suragan, Local Hardy and Rellich inequalities for sums of squares of vector fields, *Adv. Diff. Equ.* **22** (2017), 505–540.
21. M. Ruzhansky and D. Suragan, Hardy and Rellich inequalities, identities, and sharp remainders on homogeneous groups, *Adv. Math.* **317** (2017), 799–822.
22. M. Ruzhansky and D. Suragan, Uncertainty relations on nilpotent Lie groups, *Proc. R. Soc. A* **473** (2017), 20170082.
23. M. Ruzhansky and D. Suragan, Anisotropic L^2-weighted Hardy and L^2-Caffarelli–Kohn–Nirenberg inequalities, *Commun. Contemp. Math.* **19** (2017), 1750014.
24. M. Ruzhansky and D. Suragan, On horizontal Hardy, Rellich, Caffarelli–Kohn–Nirenberg and p-sub-Laplacian inequalities on stratified groups, *J. Differ. Equ.* **262** (2017), 1799–1821.
25. Z. Q. Wang and M. Willem, Caffarelli–Kohn–Nirenberg inequalities with remainder terms, *J. Funct. Anal.* **203** (2003), 550–568.

Downloaded from https://academic.oup.com/qjmath/article-abstract/70/1/305/5115498/ by Imperial College London Library user on 08 October 2019