

L^p -Nuclear Pseudo-Differential Operators on $\mathbb{Z}$ and $\mathbb{S}^1$

Julio Delgado

Departamento de Matemáticas
Universidad del Valle
Calle 13 100-00 Cali
Colombia

E-Mail: julio.delgado@correounivalle.edu.co

M. W. Wong

Department of Mathematics and Statistics
York University
4700 Keele Street
Toronto, Ontario M3J 1P3
Canada

E-Mail: mwwong@mathstat.yorku.ca

Abstract Conditions for pseudo-differential operators from $L^{p_1}(\mathbb{Z})$ into $L^{p_2}(\mathbb{Z})$ and from $L^{p_1}(\mathbb{S}^1)$ into $L^{p_2}(\mathbb{S}^1)$ to be nuclear are presented for $1 \leq p_1, p_2 < \infty$. In the cases when $p_1 = p_2$, the trace formulas are given.

2010 Mathematics Subject Classification: Primary 47G30; Secondary 47G10

Keywords: pseudo-differential operators, nuclear operators, 2/3-nuclear operators, traces, Lidskii's formula, eigenvalues

1 Introduction

Although all results in this paper are presented for $\mathbb{Z}$ and $\mathbb{S}^1$ only, extensions to the lattice $\mathbb{Z}^n$ and the torus $\mathbb{T}^n$ are valid.

Let $\mathbb{Z}$ be the set of all integers and let σ be a measurable function on $\mathbb{Z} \times \mathbb{S}^1$. Then for every sequence in $L^p(\mathbb{Z})$, $1 \leq p < \infty$, we define the sequence $T_\sigma a$ formally by

$$(T_\sigma a)(n) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-in\theta} \sigma(n, \theta) (\mathcal{F}_{\mathbb{Z}} a)(\theta) d\theta, \quad n \in \mathbb{Z},$$

where $\mathcal{F}_{\mathbb{Z}} a$ is the Fourier transform of a given by

$$(\mathcal{F}_{\mathbb{Z}} a)(\theta) = \sum_{n=-\infty}^{\infty} a(n) e^{in\theta}, \quad \theta \in [-\pi, \pi].$$

Let $\mathbb{S}^1$ be the unit circle centered at the origin and let τ be a measurable function on $\mathbb{S}^1 \times \mathbb{Z}$. Then for all f in $L^p(\mathbb{S}^1)$, $1 \leq p < \infty$, we define the function $T_\tau f$ on $\mathbb{S}^1$ formally by

$$(T_\tau f)(\theta) = \sum_{n=-\infty}^{\infty} e^{in\theta} \tau(\theta, n) \hat{f}(n), \quad \theta \in [-\pi, \pi],$$

where $\hat{f}$, sometimes denoted by $\mathcal{F}_{\mathbb{S}^1} f$, is the Fourier transform of f defined by

$$\hat{f}(n) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-in\theta} f(\theta) d\theta, \quad n \in \mathbb{Z}.$$

Sufficient conditions on σ are given in [10] to guarantee that the pseudo-differential operator $T_\sigma : L^p(\mathbb{Z}) \rightarrow L^p(\mathbb{Z})$, $1 \leq p < \infty$, is a bounded linear operator. Under additional mild conditions on σ , the bounded linear operator $T_\sigma : L^p(\mathbb{Z}) \rightarrow L^p(\mathbb{Z})$, $1 \leq p < \infty$, turns out to be compact.

Sufficient conditions on τ can be found in [11] in order to ensure that the pseudo-differential operator $T_\tau : L^p(\mathbb{S}^1) \rightarrow L^p(\mathbb{S}^1)$ is a bounded linear operator for $1 \leq p < \infty$.

In the case when $p = 2$, a simple necessary and sufficient condition for $T_\tau : L^2(\mathbb{S}^1) \rightarrow L^2(\mathbb{S}^1)$ and $T_\sigma : L^2(\mathbb{Z}) \rightarrow L^2(\mathbb{Z})$ to be Hilbert–Schmidt is that, respectively, $\tau \in L^2(\mathbb{S}^1 \times \mathbb{Z})$ and $\sigma \in L^2(\mathbb{Z} \times \mathbb{S}^1)$.

The results hitherto described can be found in the book [14]. Related results can be found in [12, 13].

The aim of this paper is to give a systematic investigation when the pseudo-differential operators $T_\sigma : L^{p_1}(\mathbb{Z}) \rightarrow L^{p_2}(\mathbb{Z})$ and $T_\tau : L^{p_1}(\mathbb{S}^1) \rightarrow L^{p_2}(\mathbb{S}^1)$ are nuclear for $1 \leq p_1, p_2 < \infty$. And if they are nuclear, we find formulas for the traces when $p_1 = p_2$.

The main tools that we use are the following results in [3, 4], which are the L^p -analogs of the L^2 -results in [1, 2].

Theorem 1.1 *Let (X_1, μ_1) and (X_2, μ_2) be σ -finite measure spaces. A bounded linear operator $A : L^{p_1}(X_1, \mu_1) \rightarrow L^{p_2}(X_2, \mu_2)$, $1 \leq p_1, p_2 < \infty$, is nuclear if and only if there exist sequences $\{g_n\}_{n=1}^\infty$ in $L^{p_2}(X_2, \mu_2)$ and $\{h_n\}_{n=1}^\infty$ in $L^{p_1'}(X_1, \mu_1)$ such that*

$$\sum_{n=1}^{\infty} \|g_n\|_{L^{p_2}(X_2, \mu_2)} \|h_n\|_{L^{p_1'}(X_1, \mu_1)} < \infty$$

and for all f in $L^{p_1}(X_1, \mu_1)$,

$$(Af)(x) = \int_{X_1} \left(\sum_{n=1}^{\infty} g_n(x) h_n(y) \right) f(y) d\mu_1(y)$$

for almost all x in X_1 .

The function k defined by

$$k(x, y) = \sum_{n=1}^{\infty} g_n(x) h_n(y), \quad x \in X_1, y \in X_2,$$

is known as a kernel of the nuclear operator $A : L^{p_1}(X_1, \mu_1) \rightarrow L^{p_2}(X_2, \mu_2)$.

Theorem 1.2 *Let (X, μ) be a σ -finite measure space. If $A : L^p(X, \mu) \rightarrow L^p(X, \mu)$, $1 \leq p < \infty$, is a nuclear operator, then the trace $\text{tr}(A)$ of $A : L^p(X, \mu) \rightarrow L^p(X, \mu)$ is given by*

$$\text{tr}(A) = \int_X k(x, x) d\mu(x),$$

where k is the kernel induced by Theorem 1.1.

Remark 1.3 Since k is defined only up to a set of measure zero on $X \times X$, the function $k(x, x)$ on the diagonal of $X \times X$ is not well-defined, resulting in a multitude of traces for a given nuclear operator. This caveat, however, does not come up in the case of the counting measure on $\mathbb{Z}$.

The following theorem of Grothendieck [6, 7] gives a subclass of nuclear operators known as 2/3-nuclear operators.

Theorem 1.4 *Let (X, μ) be a σ -finite measure space and let $1 \leq p < \infty$. If there exist sequences $\{g_n\}_{n=1}^\infty$ in $L^p(X, \mu)$ and $\{h_n\}_{n=1}^\infty$ in $L^{p'}(X, \mu)$ such that*

$$\sum_{n=1}^{\infty} \|g_n\|_{L^p(X, \mu)}^{2/3} \|h_n\|_{L^{p'}(X, \mu)}^{2/3} < \infty$$

and for all f in $L^p(X, \mu)$,

$$(Af)(x) = \int_X \left(\sum_{n=1}^{\infty} g_n(x) h_n(y) \right) f(y) d\mu(y)$$

for almost all x in X , then $A : L^p(X, \mu) \rightarrow L^p(X, \mu)$ is a nuclear operator. If $\{\lambda_j\}_{j=1}^\infty$ is the set of all eigenvalues of $A : L^p(X, \mu) \rightarrow L^p(X, \mu)$ with multiplicities counted, then

$$\mathrm{tr}(A) = \sum_{j=1}^{\infty} \lambda_j, \tag{1.1}$$

where the series is absolutely convergent.

Remark 1.5 It is important to point out that for 2/3-nuclear operators, Lidskii's formula (1.1) to the effect that the trace is the sum of all the eigenvalues with multiplicities counted is valid [5, 9, 8]. The book [5] contains a lucid exposition of nuclear operators on Banach spaces and Lidskii's formula.

In Section 2 we give results on the nuclearity of pseudo-differential operators from $L^{p_1}(\mathbb{Z})$ into $L^{p_2}(\mathbb{Z})$, where $1 \leq p_1, p_2 < \infty$, and the trace formulas when $p_1 = p_2$. Analogous results for the unit circle $\mathbb{S}^1$ are given in Section 3.

2 Pseudo-Differential Operators on $\mathbb{Z}$

To give conditions for nuclearity, we define for all n in $\mathbb{Z}$, the function e_n on $\mathbb{Z}$ by

$$e_n(m) = \delta_{nm},$$

where δ_{nm} is the Kronecker delta. Then we have the following theorem.

Theorem 2.1 *Let $k : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{C}$ be a function such that*

$$\sum_{n=-\infty}^{\infty} \left(\sum_{m=-\infty}^{\infty} |k(n, m)|^{p_2} \right)^{1/p_2} < \infty.$$

Let $A : L^{p_1}(\mathbb{Z}) \rightarrow L^{p_2}(\mathbb{Z})$, $1 \leq p_1, p_2 < \infty$, be the linear operator defined by

$$e_m(Ae_n) = k(n, m), \quad n, m \in \mathbb{Z},$$

where e_m is considered to be an element in $L^{p_2'}(\mathbb{Z})$. Then $A : L^{p_1}(\mathbb{Z}) \rightarrow L^{p_2}(\mathbb{Z})$ is a nuclear operator. Moreover if $p_1 = p_2$, then

$$\text{tr}(A) = \sum_{n=-\infty}^{\infty} k(n, n).$$

Proof For all a in $L^{p_1}(\mathbb{Z})$, we have

$$a = \sum_{n=-\infty}^{\infty} e_n(a) e_n$$

and

$$Aa = \sum_{n=-\infty}^{\infty} e_n(a) Ae_n.$$

So, by Minkowski's inequality,

$$\begin{aligned} \|Aa\|_{L^{p_2}(\mathbb{Z})} &\leq \sum_{n=-\infty}^{\infty} \|e_n(a) Ae_n\|_{L^{p_2}(\mathbb{Z})} \\ &\leq \sup_{n \in \mathbb{Z}} |a_n| \sum_{n=-\infty}^{\infty} \|Ae_n\|_{L^{p_2}(\mathbb{Z})} \\ &= \sup_{n \in \mathbb{Z}} |a_n| \sum_{n=-\infty}^{\infty} \left(\sum_{m=-\infty}^{\infty} |k(n, m)|^{p_2} \right)^{1/p_2} < \infty. \end{aligned}$$

So, $A : L^{p_1}(\mathbb{Z}) \rightarrow L^{p_2}(\mathbb{Z})$ is a bounded linear operator. Since

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \|Ae_n\|_{L^{p_2}(\mathbb{Z})} \|e_n\|_{L^{p_1}(\mathbb{Z})} &= \sum_{n=-\infty}^{\infty} \|Ae_n\|_{L^{p_2}(\mathbb{Z})} \\ &= \sum_{n=-\infty}^{\infty} \left(\sum_{m=-\infty}^{\infty} |k(n, m)|^{p_2} \right)^{1/p_2} < \infty, \end{aligned}$$

it follows from Theorem 1.1 that $A : L^{p_1}(\mathbb{Z}) \rightarrow L^{p_2}(\mathbb{Z})$ is a nuclear operator, and by Theorem 1.2,

$$\operatorname{tr}(A) = \sum_{n=-\infty}^{\infty} k(n, n).$$

□

The preceding theorem and Theorem 1.4 together imply the following theorem.

Theorem 2.2 *Let $k : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{C}$ be a function such that*

$$\sum_{n=-\infty}^{\infty} \left(\sum_{m=-\infty}^{\infty} |k(n, m)|^{p_2} \right)^{2/(3p_2)} < \infty.$$

Then the linear operator $A : L^{p_1}(\mathbb{Z}) \rightarrow L^{p_2}(\mathbb{Z})$ is a $2/3$ -nuclear operator.

We can now come back to pseudo-differential operators on $\mathbb{Z}$.

Theorem 2.3 *Let σ be a measurable function on $\mathbb{Z} \times \mathbb{S}^1$ such that we can find a function c in $L^1(\mathbb{Z})$ and a function w in $L^p(\mathbb{Z})$, $1 < p < \infty$, for which*

$$|(\mathcal{F}_{\mathbb{S}^1}\sigma)(n, m)| \leq |c(n)| |w(m)|, \quad n, m \in \mathbb{Z}.$$

Then the pseudo-differential operator $T_\sigma : L^p(\mathbb{Z}) \rightarrow L^p(\mathbb{Z})$ is a nuclear operator. Moreover,

$$\operatorname{tr}(T_\sigma) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \int_{-\pi}^{\pi} \sigma(n, \theta) d\theta.$$

Proof Let $k : \mathbb{Z} \times \mathbb{Z}$ be the function defined by

$$k(n, m) = (\mathcal{F}_{\mathbb{S}^1} \sigma)(n, n - m), \quad n, m \in \mathbb{Z}.$$

Then

$$\sum_{n=-\infty}^{\infty} \left(\sum_{m=-\infty}^{\infty} |(\mathcal{F}_{\mathbb{S}^1} \sigma)(n, m)|^p \right)^{1/p} = \sum_{n=-\infty}^{\infty} |c(n)| \left(\sum_{m=-\infty}^{\infty} |w(m)|^p \right)^{1/p} < \infty.$$

So, by Theorem 2.1, $T_\sigma : L^p(\mathbb{Z}) \rightarrow L^p(\mathbb{Z})$ is a nuclear operator and

$$\text{tr}(T_\sigma) = \sum_{n=-\infty}^{\infty} (\mathcal{F}_{\mathbb{S}^1} \sigma)(n, 0) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \int_{-\pi}^{\pi} \sigma(n, \theta) d\theta.$$

□

If we use Theorem 2.2 instead of Theorem 2.1, then we can have Lidskii's formula as well.

Theorem 2.4 *Let σ be a measurable function on $\mathbb{Z} \times \mathbb{S}^1$ such that we can find a function c in $L^{2/3}(\mathbb{Z})$ and a function w in $L^p(\mathbb{Z})$, $1 \leq p < \infty$, for which*

$$|(\mathcal{F}_{\mathbb{S}^1} \sigma)(n, m)| \leq |c(n)| |w(m)|, \quad n, m \in \mathbb{Z}.$$

Then the pseudo-differential operator $T_\sigma : L^p(\mathbb{Z}) \rightarrow L^p(\mathbb{Z})$ is a $2/3$ -nuclear operator. Moreover,

$$\text{tr}(T_\sigma) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \int_{-\pi}^{\pi} \sigma(n, \theta) d\theta,$$

which is Lidskii's formula for the sum of all eigenvalues of $T_\sigma : L^p(\mathbb{Z}) \rightarrow L^p(\mathbb{Z})$ with multiplicities counted.

Proof Again, let $k : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{C}$ be the function defined by

$$k(n, m) = (\mathcal{F}_{\mathbb{S}^1} \sigma)(n, n - m), \quad n, m \in \mathbb{Z}.$$

Then

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \left(\sum_{m=-\infty}^{\infty} |(\mathcal{F}_{\mathbb{S}^1} \sigma)(n, m)|^p \right)^{2/(3p)} &= \sum_{n=-\infty}^{\infty} |c(n)|^{2/3} \left(\sum_{m=-\infty}^{\infty} |w(m)|^p \right)^{2/(3p)} \\ &< \infty. \end{aligned}$$

So, by Theorem 2.2, the proof is complete. □

3 Pseudo-Differential Operators on $\mathbb{S}^1$

We begin this section with the kernel representation of pseudo-differential operators on $\mathbb{S}^1$.

For all f in $L^p(\mathbb{S}^1)$, we get for all θ in $[-\pi, \pi]$,

$$\begin{aligned} (T_\tau f)(\theta) &= \sum_{n=-\infty}^{\infty} e^{in\theta} \tau(\theta, n) \hat{f}(n) \\ &= \sum_{n=-\infty}^{\infty} e^{in\theta} \tau(\theta, n) \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-in\phi} f(\phi) d\phi \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{n=-\infty}^{\infty} e^{in(\theta-\phi)} \tau(\theta, n) \right) f(\phi) d\phi \\ &= \int_{-\pi}^{\pi} k(\theta, \phi) f(\phi) d\phi, \end{aligned}$$

where

$$k(\theta, \phi) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{in(\theta-\phi)} \tau(\theta, n), \quad \theta, \phi \in [-\pi, \pi]. \quad (3.1)$$

A sufficient condition on the symbol τ for the corresponding pseudo-differential operator $T_\tau : L^{p_1}(\mathbb{S}^1) \rightarrow L^{p_2}(\mathbb{S}^1)$, $1 \leq p_1, p_2 < \infty$, to be nuclear is given in the following theorem.

Theorem 3.1 *Let τ be a measurable function on $\mathbb{S}^1 \times \mathbb{Z}$ such that*

$$\sum_{n=-\infty}^{\infty} \|\tau(\cdot, n)\|_{L^{p_2}(\mathbb{S}^1)} < \infty.$$

Then $T_\tau : L^{p_1}(\mathbb{S}^1) \rightarrow L^{p_2}(\mathbb{S}^1)$ is a nuclear operator.

Proof Let $\{g_n\}_{n=-\infty}^{\infty}$ and $\{h_n\}_{n=-\infty}^{\infty}$ be sequences given by

$$g_n(\theta) = e^{in\theta} \tau(\theta, n), \quad \theta \in [-\pi, \pi], n \in \mathbb{Z},$$

and

$$h_n(\phi) = \frac{1}{2\pi} e^{-in\phi}, \quad \phi \in [-\pi, \pi], n \in \mathbb{Z}.$$

Then for all n in $\mathbb{Z}$,

$$g_n \in L^{p_2}(\mathbb{S}^1)$$

and

$$h_n \in L^{p'_1}(\mathbb{S}^1).$$

So,

$$\sum_{n=-\infty}^{\infty} \|g_n\|_{L^{p_2}(\mathbb{S}^1)} \|h_n\|_{L^{p'_1}(\mathbb{S}^1)} = \sum_{n=-\infty}^{\infty} \|\tau(\cdot, n)\|_{L^{p_2}(\mathbb{S}^1)} < \infty.$$

Thus, by Theorem 1.1, the pseudo-differential operator $T_\tau : L^{p_1}(\mathbb{S}^1) \rightarrow L^{p_2}(\mathbb{S}^1)$ has kernel k given by (3.1) is nuclear. $\square$

The same proof of Theorem gives the following theorem.

Theorem 3.2 *Let τ be a measurable function on $\mathbb{S}^1 \times \mathbb{Z}$ such that*

$$\sum_{n=-\infty}^{\infty} \|\tau(\cdot, n)\|_{L^p(\mathbb{S}^1)}^{2/3} < \infty.$$

Then the pseudo-differential operator $T_\tau : L^p(\mathbb{S}^1) \rightarrow L^p(\mathbb{S}^1)$ is 2/3-nuclear. Moreover,

$$\text{tr}(T_\tau) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \int_{-\pi}^{\pi} \tau(\theta, n) d\theta,$$

which is also equal to the sum of all eigenvalues of $T_\tau : L^p(\mathbb{S}^1) \rightarrow L^p(\mathbb{S}^1)$ with multiplicities counted.

Example 3.3 In polar coordinates, the Laplacian $\Delta_{\mathbb{S}^1}$ on $\mathbb{S}^1$ is given by

$$\Delta_{\mathbb{S}^1} = -\frac{\partial^2}{\partial \theta^2}.$$

The one-parameter semigroup generated by $\Delta_{\mathbb{S}^1}$ is $e^{t\Delta_{\mathbb{S}^1}}$, $t > 0$. In fact, $\Delta_{\mathbb{S}^1} : L^p(\mathbb{S}^1) \rightarrow L^p(\mathbb{S}^1)$, $1 \leq p < \infty$, is given by

$$\begin{aligned} (e^{\Delta_{\mathbb{S}^1}} f)(\theta) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{n=-\infty}^{\infty} e^{-n^2 t} e^{in(\theta-\phi)} f(\phi) d\phi \\ &= \sum_{n=-\infty}^{\infty} e^{in\theta} e^{-n^2 t} \hat{f}(n), \quad \theta \in [-\pi, \pi], \end{aligned}$$

for all f in $L^p(\mathbb{S}^1)$. Thus, for $t > 0$, $e^{t\Delta_S} : L^p(\mathbb{S}^1) \rightarrow L^p(\mathbb{S}^1)$ is a pseudo-differential operator on $\mathbb{S}^1$ with symbol τ_t given by

$$\tau_t(\theta, n) = e^{-n^2 t}, \quad \theta \in [-\pi, \pi], n \in \mathbb{Z},$$

and with k_t given by

$$k_t(\theta, \phi) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{in(\theta-\phi)} e^{-n^2 t}, \quad \theta, \phi \in [-\pi, \pi].$$

Now, for $t > 0$, let $\{g_n\}_{n=-\infty}^{\infty}$ and $\{h_n^t\}_{n=-\infty}^{\infty}$ be sequences in, respectively, $L^p(\mathbb{S}^1)$ and $L^{p'}(\mathbb{S}^1)$ given by

$$g_n(\phi) = e^{in\phi}, \quad \phi \in [-\pi, \pi],$$

and

$$h_n^t(\phi) = e^{-n^2 t}, \quad \phi \in [-\pi, \pi].$$

Thus, for $t > 0$,

$$\sum_{n=-\infty}^{\infty} \|g_n\|_{L^p(\mathbb{S}^1)}^{2/3} \|h_n^t\|_{L^{p'}(\mathbb{S}^1)}^{2/3} < \infty.$$

So, by Theorem 1.4, $e^{t\Delta_{\mathbb{S}^1}} : L^p(\mathbb{S}^1) \rightarrow L^p(\mathbb{S}^1)$ is nuclear and

$$\mathrm{tr}(e^{t\Delta_{\mathbb{S}^1}}) = \int_{-\pi}^{\pi} k_t(\phi, \phi) d\phi = \sum_{n=-\infty}^{\infty} e^{-n^2 t},$$

thus confirming Lidskii's theorem.

Acknowledgment This research has been supported by vic. inv. Universidad del Valle Grant No. CI 7840 and the Natural Science and Engineering Research Council of Canada. The authors are grateful to the referee for pointing out a mistake in the first version of the paper and several suggestions leading to an improvement of the paper.

References

- [1] C. Brislawn, Kernels of trace class operators, *Proc. Amer. Math. Soc.* **104** (1988), 1181–1190.
- [2] C. Brislawn, Traceable integral kernels on countably generated measure spaces, *Pacific J. Math.* **150** (1991), 229–240.
- [3] J. Delgado, The trace of nuclear operators on $L^p(\mu)$ for σ -finite Borel measures on second countable spaces, *Integr. Equ. Oper. Theory* **68** (2010), 61–74.
- [4] J. Delgado, A trace formula for nuclear operators on L^p , in *Pseudo-Differential Operators: Complex Analysis and Partial Differential Equations*, Operator Theory: Advances and Applications **205**, Birkhäuser, 2010, 181–193.
- [5] I. Gohberg, S. Goldberg and N. Krupnik, *Traces and Determinants of Linear Operators*, Birkhäuser, 2001.
- [6] A. Grothendieck, La theorie de Fredholm, *Bull. Soc. Math. France* **84** (1956), 319–384.
- [7] A. Grothendieck, *Produits Tensoriels Topologiques et Espaces Nucléaires*, Memoirs Amer. Math. Soc. **16**, 1966.
- [8] P. D. Lax, *Functional Analysis*, Wiley, 2002.
- [9] V. B. Lidskii, Nonself-adjoint operators with trace, *Dokl. Akad. Nauk SSR* **125** (1959), 485–487; *Amer. Math. Soc. Translations* **47** (1961), 43–46.
- [10] S. Molahajloo, Pseudo-differential operators on $\mathbb{Z}$, in *Pseudo-Differential Operators: Complex Analysis and Partial Differential Equations*, Operator Theory: Advances and Applications **205**, Birkhäuser, 2010, 213–221.
- [11] S. Molahajloo and M. W. Wong, Pseudo-differential operators on $\mathbb{S}^1$, in *New Developments in Pseudo-Differential Operators*, Operator Theory: Advances and Applications **189**, Birkhäuser, 2008, 297–306.

- [12] S. Molahajloo and M. W. Wong, Ellipticity, Fredholmness and spectral invariance of pseudo-differential operators on $\mathbb{S}^1$, *J. Pseudo-Differ. Oper. Appl.* **1** (2010), 183–205.
- [13] C. A. Rodriguez, L^p -estimates for pseudo-differential operators on $\mathbb{Z}^n$, *J. Pseudo-Differ. Oper. Appl.* **2** (2011), 367–375.
- [14] M. W. Wong, *Discrete Fourier Analysis*, Birkhäuser, 2011.