



Precision Measurement of the Top Quark Mass in Lepton + Jets Final States

V. M. Abazov,³¹ B. Abbott,⁶⁷ B. S. Acharya,²⁵ M. Adams,⁴⁶ T. Adams,⁴⁴ J. P. Agnew,⁴¹ G. D. Alexeev,³¹ G. Alkhazov,³⁵ A. Alton,^{56,a} A. Askew,⁴⁴ S. Atkins,⁵⁴ K. Augsten,⁷ C. Avila,⁵ F. Badaud,¹⁰ L. Bagby,⁴⁵ B. Baldin,⁴⁵ D. V. Bandurin,⁷³ S. Banerjee,²⁵ E. Barberis,⁵⁵ P. Baringer,⁵³ J. F. Bartlett,⁴⁵ U. Bassler,¹⁵ V. Bazterra,⁴⁶ A. Bean,⁵³ M. Begalli,² L. Bellantoni,⁴⁵ S. B. Beri,²³ G. Bernardi,¹⁴ R. Bernhard,¹⁹ I. Bertram,³⁹ M. Besançon,¹⁵ R. Beuselinck,⁴⁰ P. C. Bhat,⁴⁵ S. Bhatia,⁵⁸ V. Bhatnagar,²³ G. Blazey,⁴⁷ S. Blessing,⁴⁴ K. Bloom,⁵⁹ A. Boehnlein,⁴⁵ D. Boline,⁶⁴ E. E. Boos,³³ G. Borissov,³⁹ M. Borysova,^{38,l} A. Brandt,⁷⁰ O. Brandt,²⁰ R. Brock,⁵⁷ A. Bross,⁴⁵ D. Brown,¹⁴ X. B. Bu,⁴⁵ M. Buehler,⁴⁵ V. Buescher,²¹ V. Bunichev,³³ S. Burdin,^{39,b} C. P. Buszello,³⁷ E. Camacho-Pérez,²⁸ B. C. K. Casey,⁴⁵ H. Castilla-Valdez,²⁸ S. Caughron,⁵⁷ S. Chakrabarti,⁶⁴ K. M. Chan,⁵¹ A. Chandra,⁷² E. Chapon,¹⁵ G. Chen,⁵³ S. W. Cho,²⁷ S. Choi,²⁷ B. Choudhary,²⁴ S. Cihangir,⁴⁵ D. Claes,⁵⁹ J. Clutter,⁵³ M. Cooke,^{45,k} W. E. Cooper,⁴⁵ M. Corcoran,⁷² F. Couderc,¹⁵ M.-C. Cousinou,¹² D. Cutts,⁶⁹ A. Das,⁴² G. Davies,⁴⁰ S. J. de Jong,^{29,30} E. De La Cruz-Burelo,²⁸ F. Déliot,¹⁵ R. Demina,⁶³ D. Denisov,⁴⁵ S. P. Denisov,³⁴ S. Desai,⁴⁵ C. Deterre,^{20,c} K. DeVaughan,⁵⁹ H. T. Diehl,⁴⁵ M. Diesburg,⁴⁵ P. F. Ding,⁴¹ A. Dominguez,⁵⁹ A. Dubey,²⁴ L. V. Dudko,³³ A. Duperrin,¹² S. Dutt,²³ M. Eads,⁴⁷ D. Edmunds,⁵⁷ J. Ellison,⁴³ V. D. Elvira,⁴⁵ Y. Enari,¹⁴ H. Evans,⁴⁹ V. N. Evdokimov,³⁴ A. Fauré,¹⁵ L. Feng,⁴⁷ T. Ferbel,⁶³ F. Fiedler,²¹ F. Filthaut,^{29,30} W. Fisher,⁵⁷ H. E. Fisk,⁴⁵ M. Fortner,⁴⁷ H. Fox,³⁹ S. Fuess,⁴⁵ P. H. Garbincius,⁴⁵ A. Garcia-Bellido,⁶³ J. A. García-González,²⁸ V. Gavrilov,³² W. Geng,^{12,57} C. E. Gerber,⁴⁶ Y. Gershtein,⁶⁰ G. Ginther,^{45,63} O. Gogota,³⁸ G. Golovanov,³¹ P. D. Grannis,⁶⁴ S. Greder,¹⁶ H. Greenlee,⁴⁵ G. Grenier,¹⁷ Ph. Gris,¹⁰ J.-F. Grivaz,¹³ A. Grohsjean,^{15,c} S. Grünendahl,⁴⁵ M. W. Grünewald,²⁶ T. Guillemain,¹³ G. Gutierrez,⁴⁵ P. Gutierrez,⁶⁷ J. Haley,⁶⁸ L. Han,⁴ K. Harder,⁴¹ A. Harel,⁶³ J. M. Hauptman,⁵² J. Hays,⁴⁰ T. Head,⁴¹ T. Hebbeker,¹⁸ D. Hedin,⁴⁷ H. Hegab,⁶⁸ A. P. Heinson,⁴³ U. Heintz,⁶⁹ C. Hensel,¹ I. Heredia-De La Cruz,^{28,d} K. Herner,⁴⁵ G. Hesketh,^{41,f} M. D. Hildreth,⁵¹ R. Hirosky,⁷³ T. Hoang,⁴⁴ J. D. Hobbs,⁶⁴ B. Hoeneisen,⁹ J. Hogan,⁷² M. Hohlfeld,²¹ J. L. Holzbauer,⁵⁸ I. Howley,⁷⁰ Z. Hubacek,^{7,15} V. Hynek,⁷ I. Iashvili,⁶² Y. Ilchenko,⁷¹ R. Illingworth,⁴⁵ A. S. Ito,⁴⁵ S. Jabeen,^{45,m} M. Jaffré,¹³ A. Jayasinghe,⁶⁷ M. S. Jeong,²⁷ R. Jesik,⁴⁰ P. Jiang,⁴ K. Johns,⁴² E. Johnson,⁵⁷ M. Johnson,⁴⁵ A. Jonckheere,⁴⁵ P. Jonsson,⁴⁰ J. Joshi,⁴³ A. W. Jung,⁴⁵ A. Juste,³⁶ E. Kajfasz,¹² D. Karmanov,³³ I. Katsanos,⁵⁹ R. Kehoe,⁷¹ S. Kermiche,¹² N. Khalatyan,⁴⁵ A. Khanov,⁶⁸ A. Kharchilava,⁶² Y. N. Kharzheev,³¹ I. Kiselevich,³² J. M. Kohli,²³ A. V. Kozelov,³⁴ J. Kraus,⁵⁸ A. Kumar,⁶² A. Kupco,⁸ T. Kurča,¹⁷ V. A. Kuzmin,³³ S. Lammers,⁴⁹ P. Lebrun,¹⁷ H. S. Lee,²⁷ S. W. Lee,⁵² W. M. Lee,⁴⁵ X. Lei,⁴² J. Lellouch,¹⁴ D. Li,¹⁴ H. Li,⁷³ L. Li,⁴³ Q. Z. Li,⁴⁵ J. K. Lim,²⁷ D. Lincoln,⁴⁵ J. Linnemann,⁵⁷ V. V. Lipaev,³⁴ R. Lipton,⁴⁵ H. Liu,⁷¹ Y. Liu,⁴ A. Lobodenko,³⁵ M. Lokajicek,⁸ R. Lopes de Sa,⁶⁴ R. Luna-Garcia,^{28,g} A. L. Lyon,⁴⁵ A. K. A. Maciel,¹ R. Madar,¹⁹ R. Magaña-Villalba,²⁸ S. Malik,⁵⁹ V. L. Malyshev,³¹ J. Mansour,²⁰ J. Martínez-Ortega,²⁸ R. McCarthy,⁶⁴ C. L. McGivern,⁴¹ M. M. Meijer,^{29,30} A. Melnitchouk,⁴⁵ D. Menezes,⁴⁷ P. G. Mercadante,³ M. Merkin,³³ A. Meyer,¹⁸ J. Meyer,^{20,i} F. Miconi,¹⁶ N. K. Mondal,²⁵ M. Mulhearn,⁷³ E. Nagy,¹² M. Narain,⁶⁹ R. Nayyar,⁴² H. A. Neal,⁵⁶ J. P. Negret,⁵ P. Neustroev,³⁵ H. T. Nguyen,⁷³ T. Nunnemann,²² J. Orduna,⁷² N. Osman,¹² J. Osta,⁵¹ A. Pal,⁷⁰ N. Parashar,⁵⁰ V. Parihar,⁶⁹ S. K. Park,²⁷ R. Partridge,^{69,e} N. Parua,⁴⁹ A. Patwa,^{65,j} B. Penning,⁴⁵ M. Perfilov,³³ Y. Peters,⁴¹ K. Petridis,⁴¹ G. Petrillo,⁶³ P. Pétróff,¹³ M.-A. Pleier,⁶⁵ V. M. Podstavkov,⁴⁵ A. V. Popov,³⁴ M. Prewitt,⁷² D. Price,⁴¹ N. Prokopenko,³⁴ J. Qian,⁵⁶ A. Quadt,²⁰ B. Quinn,⁵⁸ P. N. Ratoff,³⁹ I. Razumov,³⁴ I. Ripp-Baudot,¹⁶ F. Rizatdinova,⁶⁸ M. Rominsky,⁴⁵ A. Ross,³⁹ C. Royon,¹⁵ P. Rubinov,⁴⁵ R. Ruchti,⁵¹ G. Sajot,¹¹ A. Sánchez-Hernández,²⁸ M. P. Sanders,²² A. S. Santos,^{1,h} G. Savage,⁴⁵ M. Savitskyi,³⁸ L. Sawyer,⁵⁴ T. Scanlon,⁴⁰ R. D. Schamberger,⁶⁴ Y. Scheglov,³⁵ H. Schellman,⁴⁸ C. Schwanenberger,⁴¹ R. Schwienhorst,⁵⁷ J. Sekaric,⁵³ H. Severini,⁶⁷ E. Shabalina,²⁰ V. Shary,¹⁵ S. Shaw,⁵⁷ A. A. Shchukin,³⁴ V. Simak,⁷ P. Skubic,⁶⁷ P. Slattery,⁶³ D. Smirnov,⁵¹ G. R. Snow,⁵⁹ J. Snow,⁶⁶ S. Snyder,⁶⁵ S. Söldner-Rembold,⁴¹ L. Sonnenschein,¹⁸ K. Soustruznik,⁶ J. Stark,¹¹ D. A. Stoyanova,³⁴ M. Strauss,⁶⁷ L. Suter,⁴¹ P. Svoisky,⁶⁷ M. Titov,¹⁵ V. V. Tokmenin,³¹ Y.-T. Tsai,⁶³ D. Tsybychev,⁶⁴ B. Tuchming,¹⁵ C. Tully,⁶¹ L. Uvarov,³⁵ S. Uvarov,³⁵ S. Uzunyan,⁴⁷ R. Van Kooten,⁴⁹ W. M. van Leeuwen,²⁹ N. Varelas,⁴⁶ E. W. Varnes,⁴² I. A. Vasilyev,³⁴ A. Y. Verkheev,³¹ L. S. Vertogradov,³¹ M. Verzocchi,⁴⁵ M. Vesterinen,⁴¹ D. Vilanova,¹⁵ P. Vokac,⁷ H. D. Wahl,⁴⁴ M. H. L. S. Wang,⁴⁵ J. Warchol,⁵¹ G. Watts,⁷⁴ M. Wayne,⁵¹ J. Weichert,²¹ L. Welty-Rieger,⁴⁸ M. R. J. Williams,⁴⁹ G. W. Wilson,⁵³ M. Wobisch,⁵⁴ D. R. Wood,⁵⁵ T. R. Wyatt,⁴¹ Y. Xie,⁴⁵ R. Yamada,⁴⁵ S. Yang,⁴ T. Yasuda,⁴⁵ Y. A. Yatsunenkov,³¹ W. Ye,⁶⁴ Z. Ye,⁴⁵ H. Yin,⁴⁵ K. Yip,⁶⁵ S. W. Youn,⁴⁵ J. M. Yu,⁵⁶ J. Zennaro,⁶² T. G. Zhao,⁴¹ B. Zhou,⁵⁶ J. Zhu,⁵⁶ M. Zielinski,⁶³ D. Zieminska,⁴⁹ and L. Zivkovic¹⁴

(D0 Collaboration)

- ¹LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
- ²Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil
- ³Universidade Federal do ABC, Santo André, Brazil
- ⁴University of Science and Technology of China, Hefei, People's Republic of China
- ⁵Universidad de los Andes, Bogotá, Colombia
- ⁶Faculty of Mathematics and Physics, Center for Particle Physics, Charles University, Prague, Czech Republic
- ⁷Czech Technical University in Prague, Prague, Czech Republic
- ⁸Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
- ⁹Universidad San Francisco de Quito, Quito, Ecuador
- ¹⁰LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France
- ¹¹LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France
- ¹²CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France
- ¹³LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France
- ¹⁴LPNHE, Universités Paris VI and VII, CNRS/IN2P3, Paris, France
- ¹⁵CEA, Irfu, SPP, Saclay, France
- ¹⁶IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France
- ¹⁷IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France
- ¹⁸III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany
- ¹⁹Physikalisches Institut, Universität Freiburg, Freiburg, Germany
- ²⁰II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen, Germany
- ²¹Institut für Physik, Universität Mainz, Mainz, Germany
- ²²Ludwig-Maximilians-Universität München, München, Germany
- ²³Panjab University, Chandigarh, India
- ²⁴Delhi University, Delhi, India
- ²⁵Tata Institute of Fundamental Research, Mumbai, India
- ²⁶University College Dublin, Dublin, Ireland
- ²⁷Korea Detector Laboratory, Korea University, Seoul, Korea
- ²⁸CINVESTAV, Mexico City, Mexico
- ²⁹Nikhef, Science Park, Amsterdam, The Netherlands
- ³⁰Radboud University Nijmegen, Nijmegen, The Netherlands
- ³¹Joint Institute for Nuclear Research, Dubna, Russia
- ³²Institute for Theoretical and Experimental Physics, Moscow, Russia
- ³³Moscow State University, Moscow, Russia
- ³⁴Institute for High Energy Physics, Protvino, Russia
- ³⁵Petersburg Nuclear Physics Institute, St. Petersburg, Russia
- ³⁶Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Física d'Altes Energies (IFAE), Barcelona, Spain
- ³⁷Uppsala University, Uppsala, Sweden
- ³⁸Taras Shevchenko National University of Kyiv, Kiev, Ukraine
- ³⁹Lancaster University, Lancaster LA1 4YB, United Kingdom
- ⁴⁰Imperial College London, London SW7 2AZ, United Kingdom
- ⁴¹The University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴²University of Arizona, Tucson, Arizona 85721, USA
- ⁴³University of California Riverside, Riverside, California 92521, USA
- ⁴⁴Florida State University, Tallahassee, Florida 32306, USA
- ⁴⁵Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
- ⁴⁶University of Illinois at Chicago, Chicago, Illinois 60607, USA
- ⁴⁷Northern Illinois University, DeKalb, Illinois 60115, USA
- ⁴⁸Northwestern University, Evanston, Illinois 60208, USA
- ⁴⁹Indiana University, Bloomington, Indiana 47405, USA
- ⁵⁰Purdue University Calumet, Hammond, Indiana 46323, USA
- ⁵¹University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵²Iowa State University, Ames, Iowa 50011, USA
- ⁵³University of Kansas, Lawrence, Kansas 66045, USA
- ⁵⁴Louisiana Tech University, Ruston, Louisiana 71272, USA
- ⁵⁵Northeastern University, Boston, Massachusetts 02115, USA
- ⁵⁶University of Michigan, Ann Arbor, Michigan 48109, USA
- ⁵⁷Michigan State University, East Lansing, Michigan 48824, USA
- ⁵⁸University of Mississippi, University, Mississippi 38677, USA
- ⁵⁹University of Nebraska, Lincoln, Nebraska 68588, USA
- ⁶⁰Rutgers University, Piscataway, New Jersey 08855, USA

⁶¹Princeton University, Princeton, New Jersey 08544, USA⁶²State University of New York, Buffalo, New York 14260, USA⁶³University of Rochester, Rochester, New York 14627, USA⁶⁴State University of New York, Stony Brook, New York 11794, USA⁶⁵Brookhaven National Laboratory, Upton, New York 11973, USA⁶⁶Langston University, Langston, Oklahoma 73050, USA⁶⁷University of Oklahoma, Norman, Oklahoma 73019, USA⁶⁸Oklahoma State University, Stillwater, Oklahoma 74078, USA⁶⁹Brown University, Providence, Rhode Island 02912, USA⁷⁰University of Texas, Arlington, Texas 76019, USA⁷¹Southern Methodist University, Dallas, Texas 75275, USA⁷²Rice University, Houston, Texas 77005, USA⁷³University of Virginia, Charlottesville, Virginia 22904, USA⁷⁴University of Washington, Seattle, Washington 98195, USA

(Received 9 May 2014; published 17 July 2014)

We measure the mass of the top quark in lepton + jets final states using the full sample of $p\bar{p}$ collision data collected by the D0 experiment in Run II of the Fermilab Tevatron Collider at $\sqrt{s} = 1.96$ TeV, corresponding to 9.7 fb^{-1} of integrated luminosity. We use a matrix element technique that calculates the probabilities for each event to result from $t\bar{t}$ production or background. The overall jet energy scale is constrained *in situ* by the mass of the W boson. We measure $m_t = 174.98 \pm 0.76 \text{ GeV}$. This constitutes the most precise single measurement of the top-quark mass.

DOI: 10.1103/PhysRevLett.113.032002

PACS numbers: 14.65.Ha

Since its discovery [1,2], the determination of the properties of the top quark has been one of the main goals of the Fermilab Tevatron Collider, recently joined by the CERN Large Hadron Collider. The measurement of the top quark mass m_t , a fundamental parameter of the standard model (SM), has received particular attention. Indeed, m_t , the mass of the W boson M_W , and the mass of the Higgs boson are related through radiative corrections that provide an internal consistency check of the SM [3]. Furthermore, m_t dominantly affects the stability of the SM Higgs potential, which has related cosmological implications [4–6]. Currently, with $m_t = 173.34 \pm 0.76 \text{ GeV}$, a world-average combined precision of about 0.5% has been achieved [7–9].

In this Letter, we present a measurement of m_t using a matrix element (ME) technique, which determines the probability of observing each event under both the $t\bar{t}$ signal and background hypotheses described by the respective MEs [10]. The overall jet energy scale (JES) is calibrated *in situ* by constraining the reconstructed invariant mass of the hadronically decaying W boson to $M_W = 80.4 \text{ GeV}$ [11]. The measurement is performed using the full set of $p\bar{p}$ collision data at $\sqrt{s} = 1.96 \text{ TeV}$ recorded by the D0 detector in Run II of the Fermilab Tevatron Collider, corresponding to an integrated luminosity of 9.7 fb^{-1} . This is an update of a previous D0 measurement that used 3.6 fb^{-1} of integrated luminosity and measured $m_t = 174.94 \pm 1.14(\text{stat} + \text{JES}) \pm 0.96(\text{syst}) \text{ GeV}$ [12]. In the present measurement, we not only use a larger data sample to improve the statistical precision, but also refine the estimation of systematic uncertainties through an updated

detector calibration, in particular improvements to the b -quark JES corrections [13], and using recent improvements in modeling the $t\bar{t}$ signal. The analysis was performed blinded in m_t .

The D0 detector central-tracking system consists of a silicon microstrip tracker and a central fiber tracker, both located within a 1.9 T superconducting solenoidal magnet [14,15], with designs optimized for tracking and vertexing at pseudorapidities $|\eta| < 3$ and $|\eta| < 2.5$, respectively [16]. A liquid-argon calorimeter with uranium absorber plates has a central section covering pseudorapidities up to $|\eta| \approx 1.1$, and two end calorimeters that extend coverage to $|\eta| \approx 4.2$, with all three housed in separate cryostats [17]. An outer muon system, at $|\eta| < 2$, consists of a layer of tracking detectors and scintillation trigger counters in front of 1.8 T iron toroids, followed by two similar layers after the toroids [18].

The top quark decays into a b quark and a W boson with $\approx 100\%$ probability assuming unitarity of the CKM matrix, resulting in a $W^+W^-b\bar{b}$ final state. This analysis is performed using lepton + jets ($\ell + \text{jets}$) final states, where one of the W bosons decays leptonically, and the other hadronically. Here, ℓ denotes either an electron (e) or a muon (μ), including those from leptonic tau decays. This analysis requires the presence of one isolated electron [19] or muon [20] with transverse momentum $p_T > 20 \text{ GeV}$ and $|\eta| < 1.1$ or $|\eta| < 2$, respectively. In addition, exactly four jets with $p_T > 20 \text{ GeV}$ within $|\eta| < 2.5$, and $p_T > 40 \text{ GeV}$ for the jet of highest p_T , are required. Jets are reconstructed using an iterative cone algorithm [21] with a cone parameter of $R = 0.5$. Jet energies are corrected to the

particle level using calibrations derived from exclusive γ + jet, Z + jet, and dijet events [13]. These calibrations account for differences in detector response to jets originating from a gluon, a b quark, and u, d, s , or c quarks. Furthermore, each event must have an imbalance in transverse momentum of $p_T > 20$ GeV expected from the undetected neutrino. Additional selection requirements to suppress background contributions from multijet (MJ) production are discussed in more detail in Ref. [22]. To further reduce background, at least one jet per event is required to be tagged as originating from a b quark (b tagged) through the use of a multivariate algorithm [23]. The tagging efficiency is on average $\approx 65\%$ for b -quark jets in this analysis, while the mistag rate for gluons and for light (u, d, s) quark jets is $\approx 5\%$. In total, 1468 and 1124 events are selected in the e + jets and μ + jets channels, respectively, which is consistent with expectation from SM predictions.

The extraction of m_t is based on the kinematic information in the event and performed with a likelihood technique using per-event probability densities (PDs) defined by the MEs of the processes contributing to the observed events. Assuming only two noninterfering contributing processes, $t\bar{t}$ and W + jets production, the per-event PD is

$$P_{\text{evt}} = A(\vec{x})[fP_{\text{sig}}(\vec{x}; m_t, k_{\text{JES}}) + (1 - f)P_{\text{bkg}}(\vec{x}; k_{\text{JES}})], \quad (1)$$

where the observed signal fraction f , m_t , and the overall multiplicative factor adjusting the energies of jets after the JES calibration k_{JES} , are parameters to be determined from data. Here, $\vec{x}$ represents the measured jet and lepton four-momenta, and $A(\vec{x})$ accounts for acceptance and efficiencies. The function P_{sig} describes the PD for $t\bar{t}$ production. Similarly, P_{bkg} describes the PD for W + jets production, which contributes 14% of the data in the e + jets and 20% in the μ + jets channels according to the normalization procedure in Ref. [22]. W + jets and MJ backgrounds have similar PD in the studied kinematic region, and thus, MJ production is accounted for in P_{evt} via P_{bkg} . MJ events contribute 12% to the e + jets and 5% to the μ + jets channels. The combined contribution from all other backgrounds amounts to about 5% in both channels.

In general, the set $\vec{x}$ of measured quantities will not be identical to the set of corresponding partonic variables $\vec{y}$ because of finite detector resolution and parton hadronization. Their relationship is described by the transfer function $W(\vec{x}, \vec{y}; k_{\text{JES}})$, where we assume that the jet and lepton angles are known perfectly. The densities P_{sig} and P_{bkg} are calculated through a convolution of the differential partonic cross section, $d\sigma(\vec{y})$, with $W(\vec{x}, \vec{y}; k_{\text{JES}})$ and the PDs for the initial-state partons, $f(q_i)$, where the q_i are the momenta of the colliding partons, by integrating over all possible parton states leading to $\vec{x}$

$$P_{\text{sig}} = \frac{1}{\sigma_{\text{obs}}^{\bar{t}t}(m_t, k_{\text{JES}})} \int \sum d\sigma(\vec{y}, m_t) d\vec{q}_1 d\vec{q}_2 f(\vec{q}_1) f(\vec{q}_2) \times W(\vec{x}, \vec{y}; k_{\text{JES}}). \quad (2)$$

The sum in the integrand extends over all possible flavor combinations of the initial state partons. The longitudinal momentum parton density functions (PDFs), $f(q_{i,z})$, are taken from the CTEQ6L1 set [24], while the dependencies $f(q_{i,x})$, $f(q_{i,y})$ on transverse momenta are taken from PDs obtained from the PYTHIA simulation [25]. The factor $\sigma_{\text{obs}}^{\bar{t}t}(m_t, k_{\text{JES}})$, defined as the expected total $t\bar{t}$ cross section, ensures that $A(\vec{x})P_{\text{sig}}$ is normalized to unity. The differential cross section, $d\sigma(\vec{y}, m_t)$, is calculated using the leading order (LO) ME for the process $q\bar{q} \rightarrow t\bar{t}$. The integration in Eq. (2) is performed over the masses of the t and $\bar{t}$ quarks which are assumed to be equal, the masses of the $W^\pm$ bosons, the energy E (curvature $1/p_T$) of the electron (muon), and $E_q/(E_q + E_{\bar{q}})$ for the quarks from the $W \rightarrow q\bar{q}'$ decay. The $M_W = 80.4$ GeV constraint for the *in situ* JES calibration is imposed by integrating over W boson masses from a Breit-Wigner prior. There are 24 possible jet-parton assignments that are summed with weights based on their consistency with the b -tagging information.

The density P_{sig} is calculated by numerical Monte Carlo (MC) integration and is identical to that in Ref. [12], except as described. The transfer function $W(\vec{x}, \vec{y}; k_{\text{JES}})$ and $\sigma_{\text{obs}}^{\bar{t}t}(m_t, k_{\text{JES}})$ are rederived using improved detector calibrations. Instead of pseudorandom numbers, we utilize the implementation of Bratley and Fox [26] of the Sobol low discrepancy sequence [27] for MC integration, which provides a reduction of about 1 order of magnitude in calculation time. Furthermore, we approximate the exact results of Eq. (2) for a grid of points in (m_t, k_{JES}) space by calculating the ME only once for each m_t and multiplying the results with the transfer function $W(\vec{x}, \vec{y}; k_{\text{JES}})$ to obtain P_{sig} for any k_{JES} . This results in another order of magnitude reduction in computation time. Both improvements are verified to provide a performance of the ME technique consistent with that in Ref. [12]. They proved essential to reduce the statistical uncertainty in evaluating most of the systematic uncertainties discussed below.

The differential partonic cross section for P_{bkg} is calculated using the LO W + 4jets MEs implemented in VECBOS [28]. The initial-state partons are all assumed to have zero transverse momentum p_T . As in the case of P_{sig} , we apply identical procedures to calculate P_{bkg} to those in Ref. [12], but using the updated transfer function $W(\vec{x}, \vec{y}; k_{\text{JES}})$ and background normalization factor.

We calculate P_{sig} and P_{bkg} on a grid in (m_t, k_{JES}) with spacings of (1 GeV, 0.01). A likelihood function, $\mathcal{L}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N; m_t, k_{\text{JES}}, f)$, is constructed at each grid point from the product of the individual P_{evt} values for the measured quantities $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N$ of the selected events, and f is determined by maximizing $\mathcal{L}$ at that grid point. The

likelihood function $\mathcal{L}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N; m_t, k_{\text{JES}})$ is then projected onto the m_t and k_{JES} axes by integrating over k_{JES} and m_t , respectively. Best unbiased estimates of m_t and k_{JES} and their statistical uncertainties are extracted from the mean and standard deviation (SD) of $\mathcal{L}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N; m_t)$ and $\mathcal{L}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N; k_{\text{JES}})$.

Simulations are used to calibrate the ME technique. Signal $t\bar{t}$ events, as well as the dominant background contribution from $W + \text{jets}$ production, are generated with ALPGEN [29] using the CTEQ6L1 set of PDFs, interfaced to PYTHIA for parton showering using the MLM matching scheme [30]. Therefore, it is the value of m_t as defined in the MC generator that is measured, and this value is expected to correspond within ≈ 1 GeV to m_t as defined in the pole mass scheme [31]. The simulation of parton showers with PYTHIA uses modified tune D0 tune A with the CTEQ6L1 PDF set and fixed Λ_{QCD} . The detector response is fully simulated through GEANT3 [32], followed by the same reconstruction algorithms as used on data. See Ref. [22] for more details on MC simulations. Contributions from MJ production are estimated with the “matrix method” [22] and modeled using a data sample, where lepton isolation requirements are inverted.

Seven samples of $t\bar{t}$ events, five at $m_t^{\text{gen}} = 165, 170, 172.5, 175, 180$ GeV for $k_{\text{JES}}^{\text{gen}} = 1$, and two at $k_{\text{JES}}^{\text{gen}} = 0.95, 1.05$ for $m_t^{\text{gen}} = 172.5$ GeV, are generated. Three samples of $W + \text{jets}$ events, at $k_{\text{JES}} = 0.95, 1$, and 1.05 , are produced. Together, the $t\bar{t}$, $W + \text{jets}$ and MJ samples are used to derive a linear calibration for the response of the ME technique to m_t and k_{JES} . For each generated $(m_t^{\text{gen}}, k_{\text{JES}}^{\text{gen}})$ point, 1000 pseudoexperiments (PEs) are constructed, each containing the same number of events as observed in data. This is done by randomly drawing simulated signal and background events according to the signal fraction f from Eq. (1), which is randomly varied according to a binomial distribution around the value measured in data. Each of the PEs contains the number of MJ events determined from the matrix method.

The signal fraction f used to construct PEs for the calibration of the method response in m_t and k_{JES} is extracted from data by maximizing the likelihood after integrating over m_t and k_{JES} . Five sets of PEs are formed, for $f = 0.5, 0.6, 0.7, 0.8$, and 0.9 at $m_t^{\text{gen}} = 172.5$ GeV, $k_{\text{JES}}^{\text{gen}} = 1$ to linearly calibrate the response of the ME technique to f . We find $f = 63\%$ in the $e + \text{jets}$ and $f = 70\%$ in the $\mu + \text{jets}$ channels, with an absolute uncertainty of 1% due to the finite size of the data sample and the calibration in f . These values are in agreement with the expectation for the signal yield assuming $\sigma_{t\bar{t}} = 7.24$ pb [33].

With f determined as above, we proceed to form PEs at the chosen $(m_t^{\text{gen}}, k_{\text{JES}}^{\text{gen}})$ points, and extract linear calibrations of the ME technique response to m_t and k_{JES} . Applying them to data, we measure $m_t = 174.98 \pm 0.58$ GeV and $k_{\text{JES}} = 1.025 \pm 0.005$, where the total




FIG. 1 (color online). (a) Two-dimensional likelihood $\mathcal{L}(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N; m_t, k_{\text{JES}})/\mathcal{L}_{\text{max}}$ for data. Fitted contours of equal probability are overlaid as solid lines. The maximum is marked with a cross. Note that the bin boundaries do not necessarily correspond to the grid points on which $\mathcal{L}$ is calculated. (b) Expected uncertainty distributions for m_t with the measured uncertainty indicated by the arrow.

statistical uncertainty on m_t also includes the statistical contribution from k_{JES} . Both uncertainties are corrected by the observed SD of the pull distributions [34]. The two-dimensional likelihood distribution in (m_t, k_{JES}) is shown in Fig. 1(a). Figure 1(b) compares the measured total statistical uncertainty on m_t with the distribution of this quantity from the PEs at $m_t^{\text{gen}} = 172.5$ GeV and $k_{\text{JES}}^{\text{gen}} = 1$. In contrast to the previous measurement [12], we do not use the JES determined in exclusive $\gamma + \text{jet}$ and dijet events with an uncertainty of $\approx 2\%$ to constrain k_{JES} . We follow this strategy because the statistical uncertainty on the measured k_{JES} value is substantially smaller than the typical uncertainty on the JES, and because k_{JES} relates jet energies at detector level to parton energies, while JES relates jet energies at detector level to jet energies at particle level. Splitting the total statistical uncertainty into two parts from m_t alone and k_{JES} , we obtain $m_t = 174.98 \pm 0.41(\text{stat}) \pm 0.41(\text{JES})$ GeV.

Comparisons of SM predictions to data for $m_t = 175$ GeV and $k_{\text{JES}} = 1.025$ are shown in Fig. 2 for the invariant mass of the jet pair matched to one of the W bosons and the invariant mass of the $t\bar{t}$ system. The kinematic reconstruction is identical to the one used in




FIG. 2 (color online). (a) Invariant mass of the jet pair matched to one of the W bosons. (b) Invariant mass of the $t\bar{t}$ system. In the ratio of data to SM prediction, the total systematic uncertainty is shown as a shaded band.

Ref. [22]. The $t\bar{t}$ signal is normalized to total cross sections of $\sigma_{t\bar{t}} = 7.8$ pb in the $e + \text{jets}$ and $\sigma_{t\bar{t}} = 7.6$ pb in the $\mu + \text{jets}$ channel, corresponding to the measured signal fraction.

Systematic uncertainties are evaluated using PEs constructed from simulated signal and background events, for three categories: modeling of signal and background events, uncertainties in the simulation of the detector response, and uncertainties associated with procedures used and assumptions made in the analysis. Contributions from these sources are listed in Table I.

The first four sources of systematic uncertainty in Table I are evaluated for $m_t^{\text{gen}} = 172.5$ GeV by comparing results for m_t using different signal models. All other systematic uncertainties are evaluated by rederiving the calibration with simulations reflecting an alternative model, and applying the alternative calibration to data. The statistical components of systematic uncertainties are ≈ 0.05 GeV for the former and ≈ 0.01 GeV for the latter sources of systematic uncertainty. The statistical components are never larger than the net difference between the default and alternative models for any of the sources of systematic uncertainty. One-sided sources of systematic uncertainties are taken as symmetric in both directions in the total quadrature sum.

We refine the evaluation procedure for several sources of systematic uncertainty compared to Ref. [12] as described

TABLE I. Summary of uncertainties on the measured top quark mass. The signs indicate the direction of the change in m_t when replacing the default by the alternative model.

Source of uncertainty	Effect on m_t (GeV)
<i>Signal and background modeling</i>	
Higher order corrections	+0.15
Initial and final state radiation	± 0.09
Hadronization and UE	+0.26
Color reconnection	+0.10
Multiple $p\bar{p}$ interactions	-0.06
Heavy flavor scale factor	± 0.06
b -jet modeling	+0.09
PDF uncertainty	± 0.11
<i>Detector modeling</i>	
Residual jet energy scale	± 0.21
Flavor-dependent response to jets	± 0.16
b tagging	± 0.10
Trigger	± 0.01
Lepton momentum scale	± 0.01
Jet energy resolution	± 0.07
Jet identification efficiency	-0.01
<i>Method</i>	
Modeling of multijet events	+0.04
Signal fraction	± 0.08
MC calibration	± 0.07
Total systematic uncertainty	± 0.49
Total statistical uncertainty	± 0.58
Total uncertainty	± 0.76

below. Details on other, typically smaller, sources of systematic uncertainty can be found in Ref. [12]. The uncertainty from *higher order corrections* is evaluated by comparing events simulated with MC@NLO [35] to ALPGEN interfaced to HERWIG [36]. The uncertainty due to the modeling of *initial and final state radiation* is constrained from Drell-Yan events [37]. As indicated by these studies, we change the amount of radiation via the renormalization scale parameter for the matching scale in ALPGEN interfaced to PYTHIA [38] up and down by a factor of 1.5. In addition, we reweight $t\bar{t}$ simulations in p_T of the $t\bar{t}$ system ($p_T^{t\bar{t}}$) to match data, and combine the two effects in quadrature. The uncertainty originating from the choice of a model for *hadronization and underlying event* (UE) is evaluated by comparing events simulated with ALPGEN interfaced to either PYTHIA or HERWIG. The JES calibration is derived using PYTHIA with a modified tune D0 tune A [13], and is expected to be valid for this configuration only. Applying it to events that use HERWIG for evolving parton showers can lead to a sizable effect on m_t . However, this effect would not be present if the JES calibration were based on HERWIG. To avoid such double counting of uncertainty sources, we evaluate the uncertainty from hadronization and UE by considering as $\vec{x}$ the momenta of particle level jets matched in (η, ϕ) space to reconstructed jets. In this evaluation, we reweight our default $t\bar{t}$ simulations in $p_T^{t\bar{t}}$ to match ALPGEN interfaced to HERWIG. A potential effect of *color reconnection* (CR) on m_t is evaluated by comparing ALPGEN events interfaced to PYTHIA with the Perugia 2011NOCR and Perugia 2011 tunes [39], where the latter includes an explicit CR model. The *residual jet energy scale* uncertainty from a potential dependence of the JES on (p_T, η) is estimated by changing the jet momenta as a function of (p_T, η) by the upper limits of JES uncertainty, the lower limits of JES uncertainty, and a linear fit within the limits of JES uncertainty. The maximum excursion in m_t is quoted as systematic uncertainty. Dedicated calibrations to account for the *flavor-dependent response to jets* originating from a gluon, a b quark, and u, d, c , or s quarks are now an integral part of the JES correction [13], and the uncertainty on m_t from these calibrations is evaluated by changing them within their respective uncertainties. This systematic uncertainty accounts for the difference in detector response to b - and light-quark jets. To evaluate the uncertainty from modeling of b tagging, differential corrections in (p_T, η) to ensure MC—data b -tagging efficiency agreement are changed within their uncertainties. The uncertainty due to the *modeling of multijet events* is evaluated by assuming a 100% uncertainty on its contribution to the data sample, i.e., by leaving it out when deriving the alternative calibration. We construct PEs with $\pm 5\%$ variations on the measured *signal fraction*, which approximately corresponds to the systematic uncertainty on the measured $t\bar{t}$ production cross section using D0 data [40], ignoring the uncertainty from integrated luminosity, and construct the PEs according to this 5% change.

In summary, we have performed a measurement of the mass of the top quark using the matrix element technique in $t\bar{t}$ candidate events in lepton + jets final states using 9.7 fb^{-1} of Run II integrated luminosity collected by the D0 detector at the Fermilab Tevatron $p\bar{p}$ Collider. The result,

$$m_t = 174.98 \pm 0.58(\text{stat}+\text{JES}) \pm 0.49(\text{syst}) \text{ GeV}, \quad \text{or}$$

$$m_t = 174.98 \pm 0.76 \text{ GeV},$$

is consistent with the values given by the current Tevatron and world combinations of the top quark mass [8,9] and achieves by itself a similar precision. With an uncertainty of 0.43%, it constitutes the most precise single measurement of the top quark mass, with a total systematic uncertainty notably smaller than any other single measurement.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); MON, NRC KI, and RFBR (Russia); CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); NRF (Korea); FOM (Netherlands); STFC and the Royal Society (United Kingdom); MSM T and GACR (Czech Republic); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); and CAS and CNSF (China).

^awith visitors from Augustana College, Sioux Falls, SD, USA,

^bwith visitors from The University of Liverpool, Liverpool, UK,

^cwith visitors from DESY, Hamburg, Germany,

^dwith visitors from Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Mexico

^ewith visitors from SLAC, Menlo Park, CA, USA,

^fwith visitors from University College London, London, UK,

^gwith visitors from Centro de Investigacion en Computacion —IPN, Mexico City, Mexico,

^hwith visitors from Universidade Estadual Paulista, São Paulo, Brazil,

ⁱwith visitors from Karlsruher Institut für Technologie (KIT) —Steinbuch Centre for Computing (SCC), D-76128 Karlsruhe, Germany,

^jwith visitors from Office of Science, U.S. Department of Energy, Washington, D.C. 20585, USA,

^kwith visitors from American Association for the Advancement of Science, Washington, D.C. 20005, USA,

^lwith visitors from Kiev Institute for Nuclear Research, Kiev, Ukraine and

^mwith visitors from University of Maryland, College Park, MD 20742, USA.

[1] S. Abachi *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **74**, 2632 (1995).

[2] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **74**, 2626 (1995).

[3] The ALEPH, CDF, D0, DELPHI, L3, OPAL, SLC Collaborations, the LEP Electroweak Working Group, the Tevatron Electroweak Working Group, and the SLD Electroweak and Heavy Flavor Groups, [arXiv:1012.2367](https://arxiv.org/abs/1012.2367); LEP Electroweak Working Group, <http://lepewwg.web.cern.ch/LEPEWWG/>.

[4] G. Degrossi, S. Vita, J. Elias-Miró, J. R. Espinosa, G. F. Giudice, G. Isidori, and A. Strumia, *J. High Energy Phys.* **08** (2012) 098.

[5] F. Bezrukov, and M. Shaposhnikov, *Phys. Lett. B* **659**, 703 (2008).

[6] A. De Simone, M. P. Hertzberg, and F. Wilczek, *Phys. Lett. B* **678**, 1 (2009).

[7] T. Aaltonen *et al.* (CDF and D0 Collaborations), *Phys. Rev. D* **86**, 092003 (2012), and references therein.

[8] T. Aaltonen *et al.* (CDF and D0 Collaborations), [arXiv:1305.3929](https://arxiv.org/abs/1305.3929), and references therein.

[9] G. Aad *et al.* (ATLAS, CDF, CMS, and D0 Collaborations), [arXiv:1403.4427](https://arxiv.org/abs/1403.4427), and references therein.

[10] V. M. Abazov *et al.* (D0 Collaboration), *Nature (London)* **429**, 638 (2004).

[11] T. Aaltonen *et al.* (CDF and D0 Collaborations), *Phys. Rev. D* **88**, 052018 (2013).

[12] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **84**, 032004 (2011).

[13] V. M. Abazov *et al.* (D0 Collaboration), [arXiv:1312.6873](https://arxiv.org/abs/1312.6873) [*Nucl. Instrum. Methods in Phys. Res., Sect. A* (to be published)].

[14] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).

[15] R. Angstadt *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **622**, 298 (2010).

[16] The D0 coordinate system is right-handed, with the z axis pointing in the direction of the Tevatron proton beam and the y axis pointing upwards. The angles ϕ and θ are the azimuthal and polar angles relative to the x and z axes, respectively. We also use, as an angular variable, the pseudorapidity defined by $\eta = -\ln[\tan(\theta/2)]$.

[17] S. Abachi *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **338**, 185 (1994).

[18] V. M. Abazov *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **552**, 372 (2005).

[19] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **750**, 78 (2014).

[20] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **737**, 281 (2014).

[21] G. Blazey *et al.*, [arXiv:hep-ex/0005012](https://arxiv.org/abs/hep-ex/0005012).

[22] V. M. Abazov *et al.* (D0 Collaboration), [arXiv:1401.5785](https://arxiv.org/abs/1401.5785) [*Phys. Rev. D* (to be published)].

[23] V. M. Abazov *et al.* (D0 Collaboration), [arXiv:1312.7623](https://arxiv.org/abs/1312.7623) [*Nucl. Instrum. Methods in Phys. Res., Sect. A* (to be published)].

[24] J. Pumplin, D. R. Stump, J. Huston, H.-L. Lai, P. Nadolsky, and W.-K. Tung, *J. High Energy Phys.* **07** (2002) 012.

[25] T. Sjöstrand, P. Edén, C. Friberg, L. Lönnblad, G. Miu, S. Mrenna, and E. Norrbin, *Comput. Phys. Commun.* **135**, 238 (2001); T. Sjöstrand, S. Mrenna, and P. Skands, *J. High Energy Phys.* **05** (2006) 026.

[26] P. Bratley and B. Fox, *ACM Trans. Math. Softw.* **14**, 88 (1988).

- [27] I. M. Sobol, Zh. Vychisl. Mat. Mat. Fiz. **7** 784 (1967) [[USSR Computational Mathematics and Mathematical Physics](#) **7**, 86 (1967)].
- [28] F. A. Berends, H. Kuijf, B. Tausk, and W. T. Giele, [Nucl. Phys.](#) **B357**, 32 (1991).
- [29] M. L. Mangano, F. Piccinini, A. D. Polosa, M. Moretti, and R. Pittau, [J. High Energy Phys.](#) **07** (2003) 001.
- [30] M. L. Mangano, M. Moretti, F. Piccinini, and M. Treccani, [J. High Energy Phys.](#) **01** (2007) 013.
- [31] J. Beringer *et al.* (Particle Data Group), [Phys. Rev. D](#) **86**, 010001 (2012).
- [32] CERN Application Software Group, *CERN Program Library Long Writeup W5013*, CERN, Geneva (1993).
- [33] P. Barnreuther, M. Czakon, and A. Mitov, [Phys. Rev. Lett.](#) **109**, 132001 (2012).
- [34] The pull is defined as $\sum_i (m_t^i - \langle m_t \rangle) / \Delta m_t$, where $i = 1, 2, \dots, 1000$ runs over all PEs. We find pull widths of 1.16 and 1.19 for m_t and k_{JES} .
- [35] S. Frixione and B. R. Webber, [J. High Energy Phys.](#) **06** (2002) 029.
- [36] G. Corcella, I. G. Knowles, G. Marchesini, S. Moretti, K. Odagiri, P. Richardson, M. H. Seymour, and B. R. Webber, [J. High Energy Phys.](#) **01** (2001) 010.
- [37] V. M. Abazov *et al.* (D0 Collaboration), [Phys. Rev. Lett.](#) **106**, 122001 (2011).
- [38] B. Cooper, J. Katzy, M. L. Mangano, A. Messina, L. Mijovic, and P. Skands, [Eur. Phys. J. C](#) **72**, 2078 (2012).
- [39] P. Z. Skands, [Phys. Rev. D](#) **82**, 074018 (2010); [arXiv:1005.3457v4](#).
- [40] V. M. Abazov *et al.* (D0 Collaboration), [Phys. Rev. D](#) **84**, 012008 (2011).