

Species-specific traits mediate avian demographic responses under past climate change

One-Sentence Summary: Interacting traits influence sensitivity of bird population sizes to climate warming and cooling over the past million years.

Ryan R Germain^{1,2*†}, Shaohong Feng^{3,4,5†}, Guangji Chen^{6,7}, Gary R. Graves^{1,8}, Joseph A. Tobias⁹, Carsten Rahbek^{1,9,10,11}, Fumin Lei¹², Jon Fjeldså^{1,13}, Peter A. Hosner^{1,10,13}, M. Thomas P. Gilbert^{14,15}, Guojie Zhang^{2,3,4,5,16*}, David Nogués-Bravo^{1*}

[†]equal authorship contributions

¹Center for Macroecology, Evolution, and Climate, The Globe Institute, University of Copenhagen, Copenhagen, Denmark.

²Villum Centre for Biodiversity Genomics, Section for Ecology and Evolution, Department of Biology, University of Copenhagen, Copenhagen, Denmark.

³Center for Evolutionary & Organismal Biology, Department of General Surgery of Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China

⁴Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, China

⁵Innovation Center of Yangtze River Delta, Zhejiang University, Jiashan, China

⁶BGI-Shenzhen, Shenzhen, China

⁷College of Life Sciences, University of Chinese Academy of Sciences, Beijing, China.

⁸Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA.

⁹Department of Life Sciences, Imperial College London, Silwood Park, Ascot, UK.

¹⁰Center for Global Mountain Biodiversity, The Globe Institute, University of Copenhagen, Copenhagen, Denmark.

¹¹Danish Institute for Advanced Study, University of Southern Denmark, Odense, Denmark

¹²Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing, China.

¹³Natural History Museum of Denmark, University of Copenhagen, Copenhagen, Denmark.

28 ¹⁴Center for Evolutionary Hologenomics, The Globe Institute, University of Copenhagen,
29 Copenhagen, Denmark.

30 ¹⁵Department of Natural History, University Museum, Norwegian University of Science and
31 Technology, Trondheim, Norway.

32 ¹⁶State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology,
33 Chinese Academy of Sciences, Kunming, China.

34 *Corresponding authors. Email: ryan.germain@bio.ku.dk, guojiezhang@zju.edu.cn,
35 dnogues@sund.ku.dk

Abstract: Anticipating species' responses to environmental change is a pressing mission in biodiversity conservation. Despite decades of research investigating how climate change may affect population sizes, historical context is lacking and the traits which mediate demographic sensitivity to changing climate remain elusive. We use whole-genome sequence data to reconstruct the demographic histories of 263 bird species over the past million years and identify networks of interacting morphological and life-history traits associated with changes in effective population size (N_e) in response to climate warming and cooling. Our results identify direct and indirect effects of key traits representing dispersal, reproduction, and survival on long-term demographic responses to climate change, thereby highlighting traits most likely to influence population responses to on-going climate warming.

Human-induced changes to the global environment are affecting biodiversity at an unprecedented rate, with animal populations having declined drastically since 1970¹⁻⁴. Despite efforts to quantify contemporary population trends and disentangle the effects of different drivers of global change, we currently lack historical context as to whether similar declines have occurred before, and if species-specific traits influence long-term demographic sensitivity to environmental challenges such as climate change⁵⁻⁷. In particular, identifying common demographic patterns over evolutionary timescales, and before the Anthropocene epoch, can reveal how life-history strategies influence population dynamics during periods of wide-spread climatic stress⁸⁻¹⁰. Elucidating these strategies will aid conservation efforts through identifying species with characteristics prone to demographic decline under current and future challenges.

Climate change is regarded as a key environmental regulator of demographic change. It is hypothesized to affect demography via its effects on reproduction, survival/growth, and dispersal^{11,12}. However, responses to climate change can vary dramatically among even closely-

related or co-occurring species because of differing life-history traits and strategies^{13–15}. Theoretical predictions and empirical evidence suggest that larger-bodied, slower-reproducing species with limited dispersal capacity are more sensitive to sustained climate change (Supplementary Table 11) because of reduced adaptive potential and/or limited ability to exploit climate refugia. However, evaluation of the role of traits on demographic sensitivity to climate change is typically tested only with contemporary data over shorter ecological timescales. Because additional environmental stressors such as land-use change and over-exploitation may mask or confound demographic responses^{9,16–18}, the ability to identify relationships between life-history traits and demographic sensitivity to climate change remains constrained when limited to contemporary data.

Periods of climate warming and cooling over the Earth's history offer a unique opportunity to quantify effects of species-specific traits on demographic sensitivity to climate change in the absence of confounding anthropogenic stressors^{5,6}. We use whole-genome sequence data¹⁹, and Pairwise Sequential Markovian Coalescent (PSMC) analysis²⁰ to reconstruct the long-term (~ one million years) demographic histories of 263 bird species, representing 39 orders distributed from the poles to the tropics. We then quantify demographic responses to the most recent warming and cooling periods before widespread human activity, and identify network effects of morphological and life-history traits related to survival/growth, reproduction, and dispersal which influence overall demographic sensitivity to climate change.

Results and Discussion:

Effective population sizes (N_e) varied substantially across avian species and over time and space (Supplementary Note 1, Supplementary Table 1). Demographic clustering revealed seven main demographic patterns, inferred from temporal patterns of N_e over the past million years (Fig. 1, Supplementary Table 3). Overall position of each species in the avian phylogeny

did not explain the observed differences among demographic patterns (Supplementary Figure 7). Passerines (order Passeriformes, which represent more than half of all extant bird species globally²¹; here $n = 123$ species) and Non-Passerines ($n = 140$ species) were unequally distributed among the demographic clusters ($X^2 = 23.81$, $df = 6$, $p < 0.001$), where Passerines were most heavily represented in clusters 1, 3, 4, and 7 (i.e., demographic peaks in the more recent upper/middle Pleistocene; Fig. 1; Supplementary Table 3). In contrast, Non-Passerines were most represented in clusters 5, 6, and 7, depicting demographic peaks during the more ancient periods of the middle/lower Pleistocene (Fig. 1, Supplementary Table 3). These results were further reflected in lower mean N_e values for Passerines in the more distant past, despite Passerines exhibiting consistently higher mean N_e than Non-Passerines over the past million years (Fig. 2A). Species currently classified as “threatened” or “near-threatened” (IUCN Red List; here $n = 34$ species) were evenly distributed among the seven main demographic patterns ($X^2 = 5.77$, $df = 6$, $p = 0.45$) and exhibited varying demographic trends over time (Fig. 2B), indicating that current conservation status is unlikely to be the result of long-term demographic variation.

Globally, mean normalized N_e for all 263 species increased from 1mya to ~500-600kya, after which it steadily declined (Fig. 3, ‘Global Average’). Across the Earth’s major zoogeographic realms (Supplementary Note 1), species differed only slightly in when they reached their mean demographic peaks. Most species followed similar patterns of higher mean N_e in more ancient periods and lower N_e values closer to 30kya (Fig. 3), and there was little concordance between geographic realm and allocation to a given demographic cluster group (Supplementary Figure 21). Overall, we detected minimal variation in demographic trends among realms (Supplementary Figure 22), indicating little effect of geographic variation on overall patterns of demographic change over time.

We used linear mixed-effect models (Gaussian distribution) to first identify the set of morphological and life-history traits most likely to be associated with overall N_e dynamics in response to climate change, for use in downstream analyses (i.e., those explicitly considering phylogenetic and trait interactions, below). For this variable selection step, we designated two periods of relatively recent warming (~147–123kya) and cooling (~122–65kya), which represent some of the most dramatic changes in global climate (i.e., $\Delta \sim 8^\circ\text{C}$ in Global Average Surface Temperature [GAST] in <60k years) over the past million years (Supplementary Figure 8), and are within the time window (~30–200kya) in which PSMC-based estimates of N_e are most precise²⁰). We quantified demographic sensitivity to climate change as the relationship between species-specific N_e and GAST during these two periods via Pearson correlation coefficients, and designated these relationships as “*Climate Warming*” and “*Climate Cooling*” responses. Of eight initial traits (body mass, brain-body ratio, tarsus length, bill length, egg mass, clutch size, incubation duration, and hand-wing index [HWI – a metric of dispersal ability]) expected to influence these estimates of demographic sensitivity to climate change (Supplementary Note 2), six were identified as key potential influencers (i.e., retained in a subset of ‘best fitting’ models, Table 1) for *Climate Warming* and/or *Climate Cooling* responses, and retained in subsequent analysis. Of these six traits, longer incubation durations and larger clutch sizes were most closely associated with increasing N_e during *Climate Warming*, whereas shorter incubation duration, lower HWI, and longer bill lengths were most closely associated with increasing N_e during *Climate Cooling* (Table 1). While goodness-of-fit for these models were low ($R^2 = 0.06$, 0.05), given the low expectation of variation in a single trait directly influencing N_e responses to climate over evolutionary time scales²², these results reveal the suite of key traits among our initial candidate traits which are most likely to be associated with long-term demographic variation under climate change.

Using these six key traits, we further characterize the phylogenetically-explicit interacting trait network of influences on demographic sensitivity to climate change. We categorized each species by their combined *Climate Warming* and *Climate Cooling* responses and used Phylogenetic Path Analysis (PPA). PPA is a hypothesis-driven framework for assessing direct and indirect effects of each trait on differentiating defined species categories, independent of their phylogenetic relationships (Supplementary Figure 10). Species exhibiting decreasing N_e under *Climate Cooling* and increasing N_e under *Climate Warming* were categorized as “Warming Positive” (Fig. 4, Supplementary Table 5). Those exhibiting increasing N_e under *Climate Cooling* and decreasing N_e under *Climate Warming* were categorized as “Warming Negative”, a scenario expected for many temperate and cold-adapted species during the 21st century. We quantified the network of trait effects on differentiating Warming Positive species from all other species in our analysis, and repeated this process for Warming Negative species. Further, we evaluated species which exhibited overall sensitivity to climate warming or cooling (i.e., Warming Positive + Warming Negative responses, hereafter “Climate Sensitive”) versus those with consistent N_e increases or decreases, and those which exhibited consistent decreases in N_e for both the *Climate Warming* and *Climate Cooling* responses (categorized as “Consistent N_e Decrease”) versus remaining species (Supplementary Table 5).

PPA revealed that Warming Positive species were best differentiated by direct effects of reproductive, survival/growth, and dispersal traits (Supplementary Table 6; Core Model D in Supplementary Figure 10). Averaging the best-performing models from this comparison indicates that N_e of species with larger body masses, lower HWI, and smaller egg masses were more likely to increase under *Climate Warming* and decrease under *Climate Cooling* (Fig. 4A). Interestingly, body mass also had indirect effects on differentiating Warming Positive species from remaining species via its significant positive influence on egg mass and its indirect correlation with a lower HWI (Fig. 4A). These results highlight potential trade-offs between the

influences of reproductive, survival/growth, and dispersal traits on changes in population size under climate warming, where the positive effects of larger body mass are offset by the associated increase in egg mass or decrease in HWI. Warming Negative species were also best differentiated via the direct effects of reproductive, survival/growth and dispersal traits, but in the opposite directions compared to what we found in Warming Positive species (Supplementary Table 6; Core Model D in Supplementary Figure 10). Model averaging further revealed that smaller-bodied species with larger eggs, longer incubation durations, and greater HWI were more likely to exhibit decreasing N_e under *Climate Warming* and increasing N_e under *Climate Cooling*, and again highlighted potential trade-offs between body mass and HWI (Fig. 4B). Trait network effects on differentiating all Climate Sensitive species were less defined, where no traits were found to have significant, direct effects on differentiating both Warming Positive and Warming Negative species from remaining species (Fig. 4C). However, species with consistently decreasing N_e tendency for both the *Climate Warming* and *Climate Cooling* responses were differentiated by smaller clutch sizes and tended to have shorter incubation durations than remaining species (Fig. 4D), providing additional evidence for the directional role of these key traits in mediating demographic responses under climate change.

Our observations of trait-network influences on long-term demographic sensitivity to climate change agree with theoretical expectations that larger-bodied, slower-reproducing species with limited dispersal capacity are likely to respond strongly to changing climate (Fig. 4). However, they also suggest that such traits may not necessarily lead to population declines under warming climate conditions as predicted from some empirical observations in contemporary populations^{12,13,15,23–25} (Supplementary Table 11). A lower HWI typical of species with limited dispersal ability was the only predictor found to significantly influence both un-networked (Table 1) and networked (Fig. 4A, B) responses to climate change, where such species tended to increase in abundance under *Climate Cooling* and decrease under *Climate*

183 *Warming*. We further found that larger body mass had direct and indirect effects on increasing
184 N_e under *Climate Warming* and decreasing N_e under *Climate Cooling* (Fig. 4A, B), contrary to
185 results from contemporary studies (i.e., over ecological time scales, Supplementary Table 11).
186 Such differences are likely due to the much longer (evolutionary) time scales investigated here
187 via our PSMC analysis (see below).

188 If morphological and life-history traits play a central role in demographic responses to
189 climate change, as determined from our historical N_e analyses, they may also be reflected in the
190 contemporary global distributions of species. Specifically, trait effects leading to an increasing
191 N_e trend under the *Climate Warming* period may be similar to those of species found in tropical
192 locations today, under the assumption that such species are adapted to warmer climates. Using
193 all available mean trait and breeding/resident range information for 2,745 avian species
194 sampled from around the world, we found that species in tropical latitudes tend to have longer
195 incubation durations and longer bills, but smaller clutch sizes, smaller eggs, lower HWIs, and
196 lower body mass (Supplementary Note S3; Supplementary Figures 23, 24). Thus, our findings
197 of historical responses to periods of climate warming do partially explain contemporary
198 biodiversity patterns. Specifically, tropical species exhibit lower HWIs, in concordance with
199 species exhibiting such traits showing increasing N_e trends under *Climate Warming*
200 (Supplementary Figures 11E, 12E). However, the opposing findings of smaller clutch sizes
201 among tropical species and the tendency toward larger clutch sizes among *Climate Warming*
202 responses (averaged model, Supplementary Figure 12E), as well as the remaining traits having
203 no significant direct effect among *Climate Warming* responses, indicate that historical responses
204 to climate change alone are only partially predictive of contemporary distributions.

205 Our study reveals seven main demographic patterns among a broad geographic (Fig. 3)
206 and phylogenetic (Supplementary Figure 7) sample of birds over the past million years, and a
207 key trait-adaptive network associated with population responses to long-term climate change in

the absence of additional human impacts. Unlike short-term responses to climate change (e.g., studies listed in Supplementary Table 11) where single traits dictate adaptive functions under strong natural selection^{12,15}, long-term adaptation (i.e., over evolutionary timescales) is influenced by overall genomic variation within species, where the effects of individual traits become saturated and diminish over time^{22,23,26,27}. This is reflected both in the very low goodness-of-fit associated with individual trait effects (i.e., un-networked) models (Table 1) and in trade-offs among several traits in PPA analysis of responses to periods of climate warming and cooling (Fig. 4, Supplementary Figures 11, 12). Over hundreds of thousands of years, interacting trait networks and trade-offs among survival/growth and reproductive traits may develop which potentially mask direct trait effects identified in studies of contemporary populations. Reconstructing long-term population dynamics from genomic data is thus a crucial component of revealing how past climatic events influenced the genetic makeup of contemporary populations over time, and for providing demographic baselines before the Anthropocene^{5,6}. Such analyses across the tree of life will provide unique insight into the natural variability of long-term demography, and help direct conservation efforts towards species more sensitive to broad-scale global environmental change.

Methods:

SNP calling, and mutation rate estimation

Whole-genome sequencing (WGS) data for 345 bird species were collected by Feng et al.¹⁹ and released as genomic resources of the Bird 10,000 Genomes (*B10K*) Project Phase II (<https://b10k.genomics.cn>). Genomes for each species were generated under a standardized protocol of library building, sequencing, and assembly as part of this previous study, thereby minimizing potential bioinformatic artefacts which could affect our analyses and interpretations. We inferred heterozygous information of each species using BWA+GATK pipeline^{20,28}. Four filtering steps were applied to obtain high-quality SNPs²⁹: a) removing homozygous SNPs (i.e., those where the genotype was encoded as 'Minor/Minor' in the variant call format file) and SNPs with more than two alternative alleles, b) removing SNPs with an interval below 10bp, c) removing SNPs with a read depth below 1/3 or over twice the average read depth across the genome, and d) removing SNPs with a root-mean-square mapping quality lower than 25.

We used branch-specific estimates of the substitution rate per site (R) from a dated phylogeny provided by the latest B10K family-phase phylogenetic study as proxies for mutation rate (μ). This latest topology was inferred based on WGS data of 363 species, assumed to be the most effective in addressing deep evolutionary relationships^{30–32}. Divergence times (t) were estimated by MCMCTree³³ with a large number of fossil records to provide by far the best calibration information^{34,35}. Both of these measures ensure the highest-possible accuracy of species-specific substitution estimates, as well as divergence times. To further convert the unit of mutation rate from per site to per site per generation, we scaled the species-specific mutation rates (μ) as $\mu = \frac{R}{t} \times T$, where T is the generation time for each species.

Demographic reconstruction over the last million years

We used the Pairwise Sequential Markovian Coalescent (PSMC) method²⁰ to reconstruct long-term changes in effective population size (N_e) for each species. Diploid genomes consist of thousands of independent loci; the coalescent approach underlying PSMC estimates the time to the most recent common ancestor (TMRCA) of the two alleles at each locus, creating an overall TMRCA distribution across the genome. The rate of coalescent events is inversely proportion to N_e , therefore the PSMC method can identify periods of changing N_e over time (e.g., if many loci were observed to coalescent at a given time point, it would indicate a lower N_e at that time)²⁰. To ensure reliability of PSMC analyses, we removed 20 species with lower-quality heterozygosity information (i.e., <18x genome-wide coverage or >25% missing data, as recommended by Nadachowska-Brzyska et al³⁶). For the remaining 325 qualified species (Supplementary Table 1), we used the PSMC settings “-N30 -t5 -r5 -p 4+251*1+4+6+10” with a reduced dataset, and then scaled results to real time using estimated mutation rate (above) and generation time³⁷. This set of parameters had an increased number of free atomic intervals (-p) on the basis of previous study²⁹, which could generate more PSMC estimates without changing the shape of the PSMC curves (Supplementary Figure 1). Data representing N_e estimates ($\times 10^4$) scaled to real time via PSMC for all 325 species are available online³⁸. Because recent and drastic population bottlenecks, and associated severe inbreeding, can lower PSMC-based estimates of N_e in more recent time periods and erase information regarding ancient N_e dynamics, we restricted downstream analyses to the 263 species with full demographic coverage over the focal time period (30kya–1mya) for which we had the most confidence in PSMC-based estimates of effective population size. Of the 62 species excluded via this step for which conservation status was available, 19 are currently listed as ‘threatened’ or ‘near-threatened’ (IUCN Redlist; www.iucnredlist.org), indicating a tendency ($X^2 = 10.78$, p-value = 0.001) towards such species to have limited polymorphism information needed to quantify long-term demographic history, once again likely due to recent population bottlenecks.

Since its initial development²⁰, the PSMC method has been widely applied across taxa to detect changes in N_e based on WGS information. Because PSMC does not rely on population data and can straightforwardly construct historical N_e dynamics over time, it is favored among *de novo* genome projects for its accuracy and precision³⁹, particularly regarding demographic changes in the more distant past^{40–43}. Although there are some generally accepted filtering strategies for sequence data to improve the accuracy of demographic reconstructions³⁶, potential biases in PSMC output can be the result of both scaling parameters (e.g., mutation rate) and population structure. For the former type, changes in the scaling parameters cause the PSMC curve to move along the axes while maintaining the overall curve shape²⁹. To account for this, we performed a robustness test of the PSMC curves in terms of sampling mutation rates from the posterior distributions on a dated phylogenetic tree (see below, Supplementary Figure 1). For the later type of bias, theoretical work has shown that the accuracy and interpretation of PSMC results can be affected by population structure, in that changes in connectivity among sub-populations may influence overall patterns of demographic change^{44–46}. While we have no *a priori* information on changes in connectivity over the last million years for our 263 species, we assume that the patterns observed are likely the result of a combination of changes in both population size and a degree of changes in population connectivity, both of which would be mediated by the effects of trait networks under periods of climate change (i.e., species with stronger dispersal ability [Hand-Wing Index] maintaining greater connectivity during periods of extreme environmental change). Thus, while not infallible, our analyses and results present a key step forward in our understanding of how trait networks can mediate demographic responses to periods of acute climate warming and cooling over the past million years.

Cluster analysis of demographic patterns and robustness tests

To investigate patterns in overall demographic fluctuations among the 263 bird species, we applied clustering analysis based on normalized N_e values. We first selected 121 time points

with equal time intervals (after log10 transformation) from 30 to 1,000kya. For each species, we extracted the corresponding N_e values for these 121 time points along the species-specific N_e trajectory and used min-max normalization to rescale values between 0 and 1. These normalized N_e values resulted in a 263*121 matrix, which was then used as the input for a hierarchical clustering analysis based on the Euclidean distance performed in *pheatmap* (with 'cluster_row = T'). Clustering methods aim to define clusters such that the total intra-group variation is minimized. In this instance (e.g., Fig. 1), species with similar N_e trajectories are found closer together on the clustering dendrogram. To summarize the major demographic patterns among species for the past million years, we used the 'cutree' function in the R package *dendextend*⁴⁷ to split the resulting clustering dendrogram into 3, 4, 5, 6 and 7 groups (i.e., $k = 3-7$), respectively (Supplementary Figures 2–5). Compared with a lower value of k (e.g., $k = 3$, Supplementary Figure 2), dividing the dendrogram into $k = 7$ subtrees (Fig. 1) made the species contained in each subtree more compact in terms of N_e pattern consistency, and was sufficient to clearly identify the major fluctuation patterns.

Different methods of data transformation can lead to different interpretations of both the appropriate number and organization of clusters. The min/max normalization approach used here focuses on the mode of demographic fluctuations, whereas other methods (e.g., rescaling the mean) take into account both the mode and degrees of change. Because our analyses center on the positive vs. negative relationships between demographic change and climate change during periods of warming and cooling, our normalization method should likewise focus on the modes of positive/negative demographic responses, rather than the degrees of change. However, to examine the robustness of our clustering analysis to the choice of data transformation, we used the 'TreeDistance' function in the R package *TreeDist*⁴⁸ to calculate the distance (0-1, where higher values indicate more similarity) between the output clustering trees generated after the min/max normalization approach and three other data transformation methods (rescaling the mean, Z-transformation, and coefficient of variation). The distance value

between the min/max normalization and Z-transformation was 0.83, and the distance value between the min/max normalization and the coefficient of variation was 0.82, indicating that the cluster designations from these three separate methods were relatively consistent. While the distance value between min/max normalization and rescaling the mean was relatively lower (0.66), indicating less consistency between clustering results, this was expected given the points made above (rescaling being based on combining the modes and degrees of fluctuations). We further used Rezende's 'phylo.signal.disc' algorithm⁴⁹ to examine the phylogenetic signal underlying cluster designations from these four data transformation methods (see below for full results from min/max normalization). For rescaling the mean, the observed number of transitions was 153, which was not significantly different than expected by chance (randomized mean transitions = 154, $p = 0.46$), indicating that phylogeny alone did not explain the observed differences among demographic patterns. Similarly, observed transitions for the Z-transformation and coefficient of variation methods did not significantly differ from chance ($p = 0.21$ and 0.99 , respectively). Thus, in terms of phylogenetic signal testing, all alternative normalization methods investigated returned consistent results/conclusions to the min/max normalization we used throughout our analyses.

Considering that transforming PSMC outputs into real time is sensitive to the value of mutation rates²⁰, we let the species-specific mutation rates vary within a reasonable range to generate random input matrixes and assess the robustness of the above clustering result. As a time-estimation algorithm based on Bayesian theory, MCMCTree provides the posterior age distributions for each node³³. Therefore, we first randomly sampled 100 estimates of the divergence time from the posterior distribution of nodes corresponding to each species. With the fixed substitution rates and generation times, 100 mutation rates were calculated for each species using these random values based on the formulas provided above and then used to scale the PSMC output from the coalescent unit to the real time unit. This gave 100 N_e trajectories per species based on varying mutation rates. Next, we randomly selected one of

these trajectories from each species and extracted the same 121 time points as before to form a new 263*121 matrix. This step was repeated 100 times and the resulting 100 matrixes were next used to produce the clustering results, as well as the split results under different number of groups (k). When $k = 7$, we first randomly selected 500 pairs of species and obtained the information on whether each pair of species belonged to the same group in the clustering dendrogram shown in Fig. 1. Then, for the same pair of species, we checked how many times the clustering results obtained from 100 random matrixes are consistent with the Fig. 1 clustering dendrogram, in terms of the group information. For example, species A and species B were from the same group in Fig. 1, and they were also considered to be from the same group in 100 random clustering results. The consistency ratio for these two species was then calculated as a percentage. The average consistency ratio for all 500 pairs of species was used to represent the robustness value for $k = 7$, and we repeated this process for all other k values (Supplementary Table 2). Robustness values increased with the number of clustering groups used, and was 83.49% when $k = 7$. A higher k value implies higher intra-group similarity and lower inter-group similarity. Thus, a pair of species from the same group at $k = 7$ is more likely to have the sufficient similarity of N_e trajectories to be clustered together even when there is some estimation bias in the mutation rate, compared to species from the same group at $k = 4$. Given that the major fluctuation patterns are identified and the robustness is over 80% at $k = 7$, we used splitting results from this grouping number in subsequent analysis (Fig. 1).

We next ran a Chi-square test to examine whether the splitting pattern for $k = 7$ could be explained in part by when species within each cluster group reached their maximum effective population size (here, normalized N_e value greater than 0.9 is considered as a maximum value; Supplementary Table 3). Compared to expected background patterns, the N_e fluctuations of species in Group 1, Group 3 and Group 4 tended to reach their maximums during the upper Pleistocene (30–129kya), while Group 6 and Group 7 reached their maximums in the middle Pleistocene (129–774kya), and species in Group 5 reached their maximums during the middle

and lower Pleistocene (774–1,000kya; Supplementary Table 3). Unlike these groups, Group 2 did not exhibit significant differences from expected background patterns during any of the time periods. To further determine the association between clustering and mean effective population size, we averaged N_e values over 30-1000kya for each species and quantified intra-group and inter-group differences (ANOVA) based on the $k = 7$ clustering groups. From these comparisons, we detected significant differences in mean N_e between Group 5 & Group 1 and between Group 5 & Group 3, where individuals clustered in Group 5 had lower mean N_e over time than those in Groups 1 & 3, but where all other pairwise comparisons revealed no significant differences in mean N_e over time (Supplementary Figure 6).

To analyze how lineages may resemble each other in overall demographic fluctuations (i.e., whether phylogenetically related species are classified into the same cluster groups), we used Rezende's 'phylo.signal.disc' algorithm⁴⁹. This algorithm compares the minimum number of character-state transitions at each node that account for the observed character distribution in the phylogeny, assuming maximum parsimony with the median of a randomized distribution (1000 randomizations were used). If the observed evolutionary transitions are significantly less than the randomized median, a phylogenetic signal is inferred. The observed transitions were 171 when $k = 7$, which were not significantly lower than expected by chance (the randomized median transitions were 175, p -value = 0.21, Supplementary Figure 7A). Even when some phylogenetically related lineages were observed to have similar N_e trajectories under $k = 4$ (red background in Supplementary Figure 7B), the similarities were still not significant (observed transitions = 123, randomized median transitions = 128, p -value = 0.13; Supplementary Figure 7B). These results indicates that there are factors other than phylogeny that contribute to the convergent patterns of demographic fluctuations.

Quantifying demographic responses to recent periods of climate warming and cooling

The last one million years of the Earth's history is punctuated by periods of abrupt climate warming and cooling⁵⁰, the most dramatic of which have occurred in relatively recent paleo-ecological time (cooling from ~122–65kya; warming from ~147–123kya; Supplementary Figure 8). Such periods of more recent paleo-ecological time also correspond to when PSMC-based estimates of N_e provide more detailed representations of demographic fluctuations within a defined window of time, and thus allow for the most accurate quantification of the relationship between climate change and changing N_e for each species.

To quantify the effects of climate warming and cooling on species-specific demographic trends over time, we first obtained the real time points and N_e values scaled from PSMC estimates for each individual within these two climate periods, and inferred the climate values for these time points from Snyder⁵⁰. Snyder⁵⁰ presented climate data over the last million years (and beyond; estimated from a spatially-weighted, multi-proxy database of over 20,000 sea surface temperature reconstructions) as the change in global average surface temperature (one value per thousand years), for example: -6.12 in 65kya and -5.99 in 66kya. Thus, we could infer the climate value corresponding to any time point between two adjacent time points based on the slope of the line between them. For example, based on the above values, the climate value of 65.70kya is calculated as -6.029. We then quantified overall N_e responses to warming and cooling (hereafter designated "*Climate Warming*" and "*Climate Cooling*") via Pearson correlation coefficients for two variables: N_e estimate and climate value. Following David⁵¹, the recommended sample size for running Pearson's r is 25, or higher. In our study, the sample size is the number of corresponding N_e estimates and climate values for each individual during *Climate Warming* or *Climate Cooling*. Under the PSMC settings "-N30 -t5 -r5 -p 4+251*1+4+6+10", the average sample size in *Climate Cooling* was 26.91, while *Climate Warming* only contained 8.60 N_e estimates in average due to a shorter period. To avoid the under-powered calculations caused by such a small sample size in *Climate Warming*, we reran the PSMC analysis with the settings "-N30 -t5 -r5 -p 4+800*1+4+6+10". These modified parameter settings allowed us to increase

the average number of samples to 26.03 in *Climate Warming*, without changing the shape of the PSMC curves (Supplementary Figure 1). In addition, we used 0.55 as the threshold for correlation coefficient to indicate statistical significance of a correlation, based on the algorithm described by Guenther⁵² with *Power* = 80%, *Alpha* = 0.05 and *Sample Size* = 25. Guenther's algorithm also allowed us to determine whether the correlation coefficients calculated for individuals with sample sizes less than 25 were significantly different from zero. If the sample size of such species is smaller than the minimum sample size estimated from its correlation coefficient based on Guenther's algorithm, we consider this correlation coefficient to be 0. For example, the minimum sample size is 13 for Pearson's $r = 0.7$, when *Power* = 80% and *Alpha* = 0.05. Therefore, if there were only 10 N_e estimates for a species, we cannot accept the correlation between N_e estimates and climate values even if its Pearson's r was as high as 0.7. After passing the above criteria, a positive correlation (at $p < 0.05$) corresponded to N_e tendency tracking changing temperature (i.e., increasing N_e under increasing temperature or decreasing N_e under decreasing temperature). All significant positive/negative correlations were visually inspected to ensure they represented a predominantly linear relationship. The above analysis criteria are outlined visually in Supplementary Figure 9.

To test the robustness of our estimated demographic responses to climate change, we randomly generated 1,000 mutation rates for each individual using the same strategy as the clustering robustness analysis (above). This gave 1,000 N_e trajectories per species based on changes in mutation rates. By using the same quantitative criteria as our observed response results (Supplementary Figure 9), we further assigned response labels to the estimated 1,000 trajectories per individual during the *Climate Warming* and *Climate Cooling* periods, respectively. Based on these response labels, we used the confidence level to represent the robustness, which is the percentage of response labels obtained from 1,000 random sampling events that are consistent with the observed response results for each individual. For example, for a species exhibiting a positive correlation in our observed results, if its response labels

inferred from the random sampling were 950 positives and 50 negatives, its confidence level was 0.95. In summary, the mean confidence level is 0.90 during *Climate Warming*, and 0.92 during *Climate Cooling*, which implied strong confidence in our observed demographic response estimates. Three types of demographic responses (“Increase”, “Decrease”, “Unrelated”; determined by the strength of positive/negative correlations at different confidence levels) are summarized in Supplementary Table 4.

For later inclusion in Phylogenetic Path Analysis (below), we categorized each species by their combined *Climate Warming* and *Climate Cooling* responses. Species which exhibited decreasing N_e under *Climate Cooling* and increasing N_e under *Climate Warming* were categorized as “Warming Positive” ($n = 33$), while those with the opposite response (N_e increase under *Climate Cooling* and decrease under *Climate Warming*) were categorized as “Warming Negative” ($n = 29$; Supplementary Table 5). There were 48 species where demographic change was not correlated with changing temperature during either *Climate Warming* and/or *Climate Cooling* (categorized as “ N_e independent of climate change”). The remaining 153 species either showed consistent N_e increases under both *Climate Warming* and *Climate Cooling* ($n = 55$), or consistent N_e decreases under both these periods of climate change ($n = 98$; Supplementary Table 5).

Quantifying the relative influence of individual traits on population responses to climate change

We used linear mixed-effect models (Gaussian distribution) and multi-model inference via the *lme4*⁵³ and *MuMIn*⁵⁴ R packages as a variable selection step to identify the set of morphological and life-history traits most likely to be associated with population responses during the periods of *Climate Warming* and *Climate Cooling* (based on Pearson correlation coefficients, above). For each response variable, we constructed a set of models with Passerine/Non-passerine as a random effect to account for Passerines exhibiting both greater mean and greater variance in N_e values across the full study period (Fig. 2A), since we aimed to

identify traits linked to demographic responses to climate change while reducing the potential effects of broad-scale phylogenetic signal on such results (see Phylogenetic Path Analysis below). Each global (i.e., all variables included) model also included eight non-collinear morphological/life-history traits predicted to influence population responses to climate change (Supplementary Note 2) as fixed effects. These eight traits (selected from an initial set of 17 candidate traits, Supplementary Table 11) were: mean unsexed mass or mean of male and female masses (g, 'body mass'), ratio of brain size to body mass ('brain-body ratio'), mean unsexed tarsus length or mean of male and female tarsus length (mm, 'tarsus length'), mean unsexed bill length or mean of male and female bill length (mm of total exposed culmen, 'bill length'), mean mass of fresh eggs (g, 'egg mass'), mean number of eggs per clutch ('clutch size'), duration of clutch incubation (days, 'incubation duration'), and 'Hand-Wing Index', measured as Kipp's Distance (distance [mm] between the tip of the first secondary feather to the tip of the longest primary feather) divided by total wing chord length (length [mm] from bend of the wing to the longest primary of the unflattened wing) and multiplied by 100. All fixed effects were standardized to mean = 0, sd = 1 to reduce potential influence of measurement scale on results and to allow direct comparison of model coefficients⁵⁵. We further confirmed key model structure assumptions of homogeneity of variance (residuals vs. predicted values) and normal distribution of residuals (Q-Q plots) for each global model. Goodness of fit (R^2) for each model was assessed by the conditional coefficient of determination⁵⁶. We then ran $n = 256$ models for all possible combinations of these eight fixed effects for each response variable (correlation coefficient during *Climate Warming* and *Climate Cooling*, respectively) and selected a subset with a difference in Akaike Information Criterion (ΔAIC) ≤ 5 from the best-fitting model. For each response variable, we then averaged parameter estimates for each predictor included in this subset of models to create one representative (full-average) estimate of the relative effects of each component on N_e responses to *Climate Warming* and N_e responses to *Climate Cooling*

(Table 1). Statistical significance for each fixed effect was assessed by whether 95% confidence intervals (CIs) overlapped zero.

Phylogenetic Path Analysis to identify trait network effects on demographic responses to climate change

We implemented the hypothesis-driven framework of Phylogenetic Path Analysis (PPA; *phylopath* R package⁵⁷) to quantify the network of direct and indirect effects of key morphological/life-history traits (identified via multi-model inference; above) on demographic responses to climate change while accounting for phylogenetic non-independence of species. PPA allows for a large number of models with complex configurations and inter-correlation of variables. In our study, six continuous variables representing our key morphological/life history traits (egg mass, clutch size, incubation duration, body mass, bill length, and hand-wing index) and one binary variable representing differing demographic responses to climate change (see below) were included in each model. A total of 14 core models with different configurations of these variables were evaluated for each of our four comparisons using both p-values and the C-statistic Information Criterion (CICc) corrected for small sample sizes (Supplementary Figure 10). Models were designed to test all possible networks of the traits and their effects on the demographic responses. Since the “Reproduction” category contained three traits (egg mass, clutch size, and incubation duration), core models assuming a direct effect of “Reproduction” on demographic responses in fact have seven derived models (i.e., at least one of the three traits having a direct effect on the demographic responses; Supplementary Figure 10). The same is true for the “Survival/Growth” category with two traits (body mass and bill length). Thus, depending on the network assumed by each core model, one core model would further be modified into 1, 3, 7 or 21 submodels.

In total, we implemented 164 models for four comparisons of the overall responses of species to *Climate Warming* and *Climate Cooling* (Supplementary Table 5) at different

confidence levels to again assess the robustness of our main results (i.e., those assessed without confidence level restrictions): 1) “Warming Positive” responses versus all remaining species, 2) “Warming Negative” species versus all remaining species, 3) species which exhibited overall sensitivity to climate warming or cooling (i.e., Warming Positive + Warming Negative species) versus species with consistent N_e increases or decreases for both the *Climate Warming* and *Climate Cooling* responses, and 4) species which exhibited consistently decreasing N_e versus all remaining species. For each comparison, PPA provides the number of independence claims made by the model, the number of parameters, the C-statistic, and the accompanying p-value, where significance (at $p < 0.05$) indicates that the available evidence rejects the model (i.e., the model does not provide a good fit to the data⁵⁸). Thus, after discarding rejected models we calculated the top-ranked model (based on ΔCICc) and the average of the best performing models ($\Delta\text{CICc} \leq 2$) for each comparison. Detailed results of the best performing models for each confidence level are provided in Supplementary Tables 6-10, with their associated directed acyclic graphs and regression coefficients provided in Supplementary Figures 11-20.

546 **Data Availability:**

547 All data and underlying these analyses (including raw N_e estimates over time for each species)
548 are available in the Dryad Digital Repository³⁸

549 **Code Availability:**

550 All code underlying these analyses is available in the Dryad Digital Repository³⁸

551

552 **Acknowledgments:**

553 This work was made possible by the generous efforts of field biologists and museum staff who
554 contributed samples to the B10k project. **Funding:** This work was supported by a National
555 Natural Science Foundation of China grant (no. 32170626 and no. 31901214) to S.F., an
556 Independent Research Fund Denmark grant to D.N-B. and G.Z. (8021-00282B), and grants
557 from the Strategic Priority Research Program of the Chinese Academy of Sciences
558 (XDB31020000), International Partnership Program of Chinese Academy of Sciences (no.
559 152453KYSB20170002), Carlsberg Foundation (CF16-0663), and Villum Foundation (no.
560 25900) to G.Z.

561 **Author contributions:**

562 R.R.G., S.F., G.Z. and D.N-B. designed the study and wrote the paper. R.R.G., S.F., and G.C.
563 carried out analyses. All authors contributed towards the assembly of genomic and
564 morphological/life-history data, discussion of results, and reviewing and editing the manuscript.

565 **Competing interests:**

566 The authors declare no competing interests.

Table 1 - Parameter estimates, standard errors (SE), and upper and lower 95% confidence intervals (UCI, LCI) from averaged two-sided linear mixed models (random effect = Passerine/Non-passerine) evaluating the relative effects of morphological and life-history traits on demographic responses to climate warming and cooling (measured as correlation coefficient between N_e change and climate change). For each response, we ran 256 models involving all possible combinations of eight predictor variables (body mass [g], ratio of brain size to body mass, tarsus length [mm], hand-wing index [HWI], bill length [mm], egg mass [g], incubation duration [days], and clutch size; all z-scaled to remove the effects of measurement scale), selected a 'best models' subset ($\Delta AIC \leq 5$ from the best-fitting model) and averaged parameter estimates within this subset. R^2 represents the goodness-of-fit of the global model (including all explanatory variables) for each response.

		Estimate	SE	z-value	p	LCI	UCI
<i>Climate Warming</i> (R ² = 0.06)	Intercept	0.0003	0.07				
	Incubation duration	0.15	0.08	1.92	0.06	-0.003	0.30
	Clutch size	0.13	0.07	1.91	0.06	-0.004	0.27
	Egg mass	0.12	0.08	1.61	0.11	-0.03	0.27
	HWI	-0.09	0.07	1.34	0.18	-0.22	0.04
	Body mass	0.05	0.07	0.82	0.41	-0.08	0.19
<i>Climate Cooling</i> (R ² = 0.05)	Intercept	-0.0005	0.08				
	Incubation duration	-0.14	0.07	1.93	0.05	-0.28	0.002
	HWI	-0.14	0.07	2.00	0.05	-0.28	-0.003
	Bill length	0.05	0.07	0.70	0.49	-0.09	0.18

579 *Values marked in bold highlight statistically significant predictors (i.e., CIs do not overlap zero).

Fig. 1. Demographic histories of 263 avian species from 30,000 to 1 million years ago (x-axis presented on the $\log_{10}$ scale). Cluster analysis of normalized N_e values reveal seven main demographic patterns over the Upper/Middle/Lower(L) Pleistocene. Clusters designating Groups 1–7 (see Supplementary Table 1 for species included in each group) were based on overall similarity of long-term N_e patterns but can be distinguished by when the majority of species reached their relative peak (see Supplementary Table 3).

Fig. 2. Demographic histories of different avian groups from 30,000 to 1 million years ago (all x-axes presented on the $\log_{10}$ scale). A) Mean log effective population size (N_e) values ($\pm$ SD) of Passerine and Non-Passerine species. Passerines exhibited consistently higher N_e values (mean sample difference = 9.21) across 120 equally-spaced (loglinear) time points from 30–1000kya (paired two-sided t-test, $t_{119} = 27.9$, $p = 2.2 \times 10^{-16}$). B) Examples of differing demographic histories of species currently designated as “threatened” under IUCN Red List status, where species arriving at similar N_e values at ~30kya follow differing demographic patterns over time (illustrations by JF).

Fig 3. Mean change in normalized N_e from 30,000 to 1 million years ago (30–1000kya) for 263 avian species, summarized by zoogeographic realm (top panel, following Holt et al.⁵⁹). For each panel, the number of species representing each realm is given in parenthesis. Colored dots depict mean normalized N_e at 120 equally-spaced (loglinear) time points from 30–1000kya, while shaded area depicts SD at each point. Full demographic curves of each species are provided in the background (grey) to show overall variation within each realm.

Fig. 4. Network effects of key morphological and life-history traits on demographic responses to climate warming and climate cooling, shown as directed acyclic graphs (left) and corresponding standardized regression coefficients ($\pm$ SE; right) for averaged best performing models. A) Comparison of “Warming Positive” responses ($n = 33$) versus all remaining species ($n = 230$) reveals that larger body masses, smaller egg masses, and lower HWI are associated with demographic increases during *Climate Warming* and decreases during *Climate Cooling* (statistical significance of trait effects in all panels assessed by whether SEs overlap zero). B) Comparison of “Warming Negative” responses ($n = 29$) versus all remaining species ($n = 234$) reveals that smaller body masses, larger egg masses, greater HWI, and longer incubation durations are significantly associated with demographic increases during *Climate Cooling* and decreases during *Climate Warming*. C) Comparison of all species which exhibited sensitivity to *Climate Warming* or *Cooling* ($n = 33 + 29$) versus those with consistent N_e increases or decreases ($n = 98 + 55$) reveals no traits significantly influenced overall species sensitivity to changing climate conditions. D) Comparison of species with consistent N_e decreases during *Climate Warming* and *Climate Cooling* ($n = 98$) versus all remaining species ($n = 165$) again revealed no significant influences, although species with smaller clutch sizes, higher HWI, and shorter incubation durations did tend to exhibit demographic decrease during both *Climate Warming* and *Climate Cooling*.

References

1. Butchart, S. H. M. *et al.* Global biodiversity: indicators of recent declines. *Science* **328**, 1164–1168 (2010).
2. Dirzo, R. *et al.* Defaunation in the Anthropocene. *Science* **345**, 401–406 (2014).
3. IPBES. *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. (IPBES secretariat, 2019).
4. Rosenberg, K. V. *et al.* Decline of the North American avifauna. *Science* **366**, 120–124 (2019).
5. Fordham, D. A. *et al.* Using paleo-archives to safeguard biodiversity under climate change. *Science* **369**, eabc5654 (2020).
6. Nogués-Bravo, D. *et al.* Cracking the code of biodiversity responses to past climate change. *Trends Ecol. Evol.* **33**, 765–776 (2018).
7. Foden, W. B. *et al.* Climate change vulnerability assessment of species. *WIREs Climate Change* **10**, e551 (2019).
8. Chattopadhyay, B., Garg, K. M., Ray, R. & Rheindt, F. E. Fluctuating fortunes: genomes and habitat reconstructions reveal global climate-mediated changes in bats' genetic diversity. *Proc. R. Soc. B.* **286**, 20190304 (2019).
9. Chen, L. *et al.* Large-scale ruminant genome sequencing provides insights into their evolution and distinct traits. *Science* **364**, eaav6202 (2019).
10. Peart, C. R. *et al.* Determinants of genetic variation across eco-evolutionary scales in pinnipeds. *Nat. Ecol. Evol.* **4**, 1095–1104 (2020).
11. Jenouvrier, S. Impacts of climate change on avian populations. *Glob. Change Biol.* **19**, 2036–2057 (2013).

- 644 12. Pacifici, M. *et al.* Species' traits influenced their response to recent climate change. *Nat.*
645 *Clim. Chang.* **7**, 205–208 (2017).
- 646 13. Foden, W. B. *et al.* Identifying the world's most climate change vulnerable species: a
647 systematic trait-based assessment of all birds, amphibians and corals. *PLOS ONE* **8**,
648 e65427 (2013).
- 649 14. Telenský, T., Klvaňa, P., Jelínek, M., Cepák, J. & Reif, J. The influence of climate variability
650 on demographic rates of avian Afro-palearctic migrants. *Sci. Rep.* **10**, 17592 (2020).
- 651 15. Jiguet, F., Gadot, A.-S., Julliard, R., Newson, S. E. & Couvet, D. Climate envelope, life
652 history traits and the resilience of birds facing global change. *Glob. Change Biol.* **13**, 1672–
653 1684 (2007).
- 654 16. Lorenzen, E. D. *et al.* Species-specific responses of Late Quaternary megafauna to climate
655 and humans. *Nature* **479**, 359–364 (2011).
- 656 17. Hung, C.-M. *et al.* Drastic population fluctuations explain the rapid extinction of the
657 passenger pigeon. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 10636–10641 (2014).
- 658 18. Feng, S. *et al.* The genomic footprints of the fall and recovery of the crested ibis. *Curr. Biol.*
659 **29**, 340–349.e7 (2019).
- 660 19. Feng, S. *et al.* Dense sampling of bird diversity increases power of comparative genomics.
661 *Nature* **587**, 252–257 (2020).
- 662 20. Li, H. & Durbin, R. Inference of human population history from individual whole-genome
663 sequences. *Nature* **475**, 493–496 (2011).
- 664 21. Fjeldså, J., Christidis, L. & Ericson, P. G. P. *The Largest Avian Radiation. The Evolution of*
665 *Perching Birds, or the Order Passeriformes.* (Lynx ed, 2020).
- 666 22. Bay, R. A. *et al.* Genomic signals of selection predict climate-driven population declines in a
667 migratory bird. *Science* **359**, 83–86 (2018).
- 668 23. Parmesan, C. Ecological and evolutionary responses to recent climate change. *Annu. Rev.*
669 *Ecol. Evol. Syst.* **37**, 637–669 (2006).

- 670 24. Angert, A. L. *et al.* Do species' traits predict recent shifts at expanding range edges? *Ecol.*
671 *Lett.* **14**, 677–689 (2011).
- 672 25. McCain, C. M. & King, S. R. B. Body size and activity times mediate mammalian responses
673 to climate change. *Glob. Change Biol.* **20**, 1760–1769 (2014).
- 674 26. Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol.*
675 *Evol.* **23**, 38–44 (2008).
- 676 27. Hancock, A. M. *et al.* Adaptation to climate across the *Arabidopsis thaliana* genome.
677 *Science* **334**, 83–86 (2011).
- 678 28. McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing
679 next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
- 680 29. Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G. & Ellegren, H. Temporal dynamics
681 of avian populations during pleistocene revealed by whole-genome sequences. *Curr. Biol.*
682 **25**, 1375–1380 (2015).
- 683 30. Jarvis, E. D. *et al.* Whole-genome analyses resolve early branches in the tree of life of
684 modern birds. *Science* **346**, 1320–1331 (2014).
- 685 31. Rokas, A., Williams, B. L., King, N. & Carroll, S. B. Genome-scale approaches to resolving
686 incongruence in molecular phylogenies. *Nature* **425**, 798–804 (2003).
- 687 32. Wolf, Y. I., Rogozin, I. B., Grishin, N. V. & Koonin, E. V. Genome trees and the tree of life.
688 *Trends Genet.* **18**, 472–479 (2002).
- 689 33. Yang, Z. PAML: a program package for phylogenetic analysis by maximum likelihood.
690 *Comput. Appl. Biosci.* **13**, 555–556 (1997).
- 691 34. Forest, F. Calibrating the tree of life: fossils, molecules and evolutionary timescales. *Ann.*
692 *Bot.* **104**, 789–794 (2009).
- 693 35. Magallón, S. A. Dating lineages: molecular and paleontological approaches to the temporal
694 framework of clades. *Int. J. Plant Sci.* **165**, S7–S21 (2004).

36. Nadachowska-Brzyska, K., Burri, R., Smeds, L. & Ellegren, H. PSMC analysis of effective population sizes in molecular ecology and its application to black-and-white *Ficedula* flycatchers. *Mol. Ecol.* **25**, 1058–1072 (2016).
37. Bird, J. P. *et al.* Generation lengths of the world's birds and their implications for extinction risk. *Cons. Biol.* **34**, 1252–1261 (2020).
38. Germain, R. *et al.* Data from: Species-specific traits mediate avian demographic responses under past climate change. Dryad, Dataset, <http://datadryad.org/stash/dataset/doi:10.5061/dryad.fn2z34tz8> (2022).
39. Patton, A. H. *et al.* Contemporary demographic reconstruction methods are robust to genome assembly quality: a case study in Tasmanian devils. *Mol. Biol. Evol.* **36**, 2906–2921 (2019).
40. Liu, X. & Fu, Y.-X. Exploring population size changes using SNP frequency spectra. *Nat. Genet.* **47**, 555–559 (2015).
41. Terhorst, J. & Song, Y. S. Fundamental limits on the accuracy of demographic inference based on the sample frequency spectrum. *Proc. Nat. Acad. Sci. U.S.A.* **112**, 7677–7682 (2015).
42. Terhorst, J., Kamm, J. A. & Song, Y. S. Robust and scalable inference of population history from hundreds of unphased whole genomes. *Nat. Genet.* **49**, 303–309 (2017).
43. Lapierre, M., Lambert, A. & Achaz, G. Accuracy of Demographic Inferences from the Site Frequency Spectrum: The Case of the Yoruba Population. *Genetics* **206**, 439–449 (2017).
44. Mazet, O., Rodríguez, W., Grusea, S., Boitard, S. & Chikhi, L. On the importance of being structured: instantaneous coalescence rates and human evolution—lessons for ancestral population size inference? *Heredity* **116**, 362–371 (2016).
45. Chikhi, L. *et al.* The IICR (inverse instantaneous coalescence rate) as a summary of genomic diversity: insights into demographic inference and model choice. *Heredity* **120**, 13–24 (2018).

46. Teixeira, H. *et al.* Impact of model assumptions on demographic inferences: the case study of two sympatric mouse lemurs in northwestern Madagascar. *BMC Ecol. Evol.* **21**, 197 (2021).
47. Galili, T. dendextend: an R package for visualizing, adjusting and comparing trees of hierarchical clustering. *Bioinformatics* **31**, 3718–3720 (2015).
48. Smith, M. R. TreeDist: Distances Between Phylogenetic Trees. R Package version 2.5.0 (2019) doi:10.5281/ZENODO.3528124.
49. Paleo-López, R. *et al.* A phylogenetic analysis of macroevolutionary patterns in fermentative yeasts. *Ecol. Evol.* **6**, 3851–3861 (2016).
50. Snyder, C. W. Evolution of global temperature over the past two million years. *Nature* **538**, 226–228 (2016).
51. David, F. N. *Tables of the Ordinates and Probability Integral of the Distribution of the Correlation Coefficient in Small Samples.* (Cambridge University Press, 1938).
52. Guenther, W. C. Desk Calculation of Probabilities for the Distribution of the Sample Correlation Coefficient. *Am. Stat.* **31**, 45–48 (1977).
53. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J Stat. Softw.* **67**, 1–48 (2015).
54. Barton, K. MuMIn: Multi-Model Inference. R package version 1.47.1 (2022).
55. White, G. & Burnham, K. Program MARK: survival estimation from populations of marked animals. *Bird Study* **46**, 120–139 (1999).
56. Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods Ecol. Evol.* **4**, 133–142 (2013).
57. Bijl, W. van der. phylopath: Perform Phylogenetic Path Analysis. R package version 1.1.3 (2021).

- 745 58. Gonzalez-Voyer, A. & von Hardenberg, A. An Introduction to Phylogenetic Path Analysis. in
746 *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology*
747 (ed. Garamszegi, L.) (Springer Berlin Heidelberg, 2014).
748 59. Holt, B. G. *et al.* An update of Wallace's zoogeographic regions of the world. *Science* **339**,
749 74–78 (2013).

750

751


752 **Supplemental Materials**

753 Supplementary Notes 1 to 3

754 Supplementary Tables 1 to 12

755 Supplementary Figures 1 to 24

756 References (60–104)



Log Effective Population Size (N_e)

120
100
80
60
40
20
0
-20

3×10^4

10^5

10^6

Time Since Present (Years)

Non-Passerine

Passerine



Effective Population Size ($N_e \times 10^4$)



Southern cassowary

Yellowhead

MacQueen's bustard


White-crested guan


3×10^4









10^5

10^6

Time Since Present (Years)







→ Positive — Negative abs value < 0.1