

OBSERVATIONS

Bariatric Surgery Does Not Exacerbate and May Be Beneficial for the Microvascular Complications of Type 2 Diabetes

The effects of bariatric surgery on microvascular complications remain underexplored despite more than 40 years of surgery on patients with type 2 diabetes. The literature has focused predominantly on glycemic control and very little on diabetes-related complications, which confer high rates of morbidity and mortality. Our aim was to assess whether the gastric bypass, sleeve gastrectomy, and gastric banding procedures are safe for the retinal and renal complications of type 2 diabetes, especially as they lead to rapid improvements in glycemia, which may paradoxically cause or exacerbate microvascular complications.

Prospectively collected retinal photographs (two-field) and urine albumin creatinine ratios (ACRs) were analyzed retrospectively in 84 consecutive patients with type 2 diabetes before and 12–18 months after surgery. The results were confirmed by an independent ophthalmologist who was blinded to clinical information. Improvement or deterioration was defined as a decrease or increase of at least two grading steps, respectively. Each patient was assigned a numerical retinopathy score based on retinal disease severity to allow noncategorical statistical analysis (1). Metformin and/or long-acting insulin analogs were used for glycemic control after gastric bypass or sleeve gastrectomy, and gastric banding patients had preoperative glucose-lowering medication doses adjusted. Antihypertensives and lipid-lowering medications were continued if tolerated.

Data are expressed as mean $\pm$ SEM, median (interquartile range), or percentage. Eighty-four patients (67% female) 50.2 ± 1.1 years of age with BMI = 47.5 ± 0.8 kg/m² were studied. There were 59 gastric bypass patients

(70.2%), 19 sleeve gastrectomy (22.6%), and 6 gastric banding (7.1%). Of the 67 patients with complete retinal data, 5 (7.5%) had an improvement in retinal disease, 1 (1.5%) had deterioration, and 61 (91.0%) had no change. In the subgroup of 28 patients with diabetic retinopathy preoperatively, 5 (17.8%) improved, 1 (3.6%) deteriorated, 22 (78.6%) had no change, and no one normalized after bariatric surgery. In the subgroup with preoperative disease, mean retinopathy scores decreased significantly from 4.7 ± 0.6 to 3.3 ± 0.5 ($P = 0.004$). Thirty-two patients (42.7%) had preoperative albuminuria, and their ACRs improved from 7.6 (4.7–24.5) to 2.2 (1.0–17.3) mg/mmol ($P < 0.001$). The total coefficient of variation for ACR was 30.2%. These changes did not correlate with reductions in weight, glycemia, or blood pressure. The stabilization and improvements in microvascular complications were observed in the context of a reduction in medication usage in all three groups.

Bariatric surgery does not exacerbate and may also be beneficial for the retinal and renal complications of type 2 diabetes. Considering that diabetes-related complications progressively deteriorate over time (2), the effects of surgery to halt and in some cases reverse retinopathy were encouraging. Improvement of retinopathy is rarely encountered, and the rates of deterioration observed after surgery, albeit in a small number of patients, did not exceed the rates reported in the literature (3). Our results should be confirmed in larger, randomized, controlled clinical trials. Further elucidation of the physiological mechanisms at play, such as reductions in systemic or local inflammation (4) and direct effects of gut hormones (5), could eventually lead to the development of more effective and safe pharmacotherapy for both type 2 diabetes and obesity.

ALEXANDER D. MIRAS, MRCP, BSC^{1,2}

LING LING CHUAH, MRCP¹

GERASSIMOS LASCARATOS, MSC, MRCS³

SANA FARUQ, BSC¹

AJAY A. MOHITE, BSC¹

PRIYA R. SHAH, BSC¹

MAHI GILL, BSC¹

SABRINA N. JACKSON, BSC¹

DESMOND G. JOHNSTON, FRCP, MD⁴

TORSTEN OLBERS, MD, PHD¹

CAREL W. LE ROUX, MRCP, PHD¹

From the ¹Imperial Weight Centre, Imperial College London, London, U.K.; the ²Molecular and Metabolic Imaging Group, Institute of Clinical Sciences, Imperial College London, London, U.K.; ³Moorfields Eye Hospital, London, U.K.; and ⁴Metabolic Medicine, Imperial College London, London, U.K.

Corresponding author: Alexander D. Miras, a.miras@nhs.net.

DOI: 10.2337/dc11-2353

© 2012 by the American Diabetes Association.

Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—A.D.M. is funded by the Medical Research Council Research Training Fellowship G0902002. The Division of Diabetes, Endocrinology and Metabolism, Imperial College London is funded by grants from the Medical Research Council, the Biotechnology and Biological Sciences Research Council, and the National Institute for Health Research (NIHR) and an Integrative Mammalian Biology Capacity Building Award, a FP7-HEALTH-2009-241592 EuroCHIP grant, and the NIHR Imperial Biomedical Research Centre Funding Scheme.

No potential conflicts of interest relevant to this article were reported.

A.D.M. wrote the manuscript, researched data, and contributed to the discussion. L.L.C. and G.L. researched data and contributed to the discussion. S.F., A.A.M., P.R.S., M.G., and S.N.J. researched data. D.G.J., T.O., and C.W.I.R. contributed to the discussion and reviewed and edited the manuscript. C.W.I.R. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank all the staff of the Imperial Weight Centre for their tireless efforts in looking after these patients.

References

1. Klein R, Klein BE, Magli YL, et al. An alternative method of grading diabetic retinopathy. *Ophthalmology* 1986;93:1183–1187
2. Fong DS, Aiello L, Gardner TW, et al.; American Diabetes Association. Retinopathy in diabetes. *Diabetes Care* 2004;27(Suppl. 1):S84–S87
3. Ohkubo Y, Kishikawa H, Araki E, et al. Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 1995;28:103–117
4. Bueter M, Dubb SS, Gill A, et al. Renal cytokines improve early after bariatric surgery. *Br J Surg* 2010;97:1838–1844
5. Zhang Y, Zhang J, Wang Q, et al. Intravitreal injection of exendin-4 analogue protects retinal cells in early diabetic rats. *Invest Ophthalmol Vis Sci* 2011;52:278–285