

# COMPARING THE BEST REPLY STRATEGY AND MEAN FIELD GAMES: THE STATIONARY CASE

MATT BARKER

*Department of Mathematics, Imperial College London, London, SW7 2AZ, UK and Grantham Institute,  
Imperial College London, London, SW7 2AZ, UK  
Email: m.barker17@imperial.ac.uk*

PIERRE DEGOND

*Department of Mathematics, Imperial College London, London, SW7 2AZ, UK  
Email: p.degond@imperial.ac.uk*

MARIE-THERESE WOLFRAM

*University of Warwick, Mathematics Institute, Gibbet Hill Road, CV47AL Coventry, UK and Radon  
Institute for Computational and Applied Mathematics, Altenbergerstr. 69, 4040 Linz, Austria  
Email: m.wolfram@warwick.ac.uk*

## ACKNOWLEDGEMENTS

MB acknowledges support by the Natural Environment Research Council (NERC) under training grant no. NE/L002515/1. PD acknowledges support by the Engineering and Physical Sciences Research Council (EPSRC) under grants no. EP/M006883/1 and EP/N014529/1, by the Royal Society and the Wolfson Foundation through a Royal Society Wolfson Research Merit Award no. WM130048 and by the National Science Foundation (NSF) under grant no. RNMS11-07444 (KI-Net). PD is on leave from CNRS, Institut de Mathématiques de Toulouse, France. MTW acknowledges partial support from the Austrian Academy of Sciences under the New Frontier's grant NST-001 and the EPSRC under the First Grant EP/P01240X/1.

## AMS SUBJECT CLASSIFICATION

35Q84, 35Q91, 35J15, 35J57, 91A13, 91A23, 49N70, 34C60, 37M05

## KEY WORDS

Mean field games, best reply strategy, stationary Fokker–Planck equation

DATA STATEMENT: NO NEW DATA WERE COLLECTED IN THE COURSE OF THIS RESEARCH.

## 1. ABSTRACT

Mean field games (MFGs) and the best reply strategy (BRS) are two methods of describing competitive optimisation of systems of interacting agents. The latter can be interpreted as an approximation of the respective MFG system, see [3, 5, 20]. In this paper we present an analysis and comparison of the two approaches in the stationary case. [We provide novel existence and uniqueness results for the stationary boundary value problems related to the MFG and BRS formulations, and we present an analytical and numerical comparison of the two paradigms in some specific modelling situations.](#)

## 2. INTRODUCTION

Mean field games (MFGs) describe the dynamics of large interacting agent systems, in which individuals determine their optimal strategy by minimising a given cost functional. The extensive current literature is based on the original work of Lasry and Lions [36, 37, 38] and Huang, Caines and Malhamé [34, 33, 35]. MFGs have been used successfully in many different disciplines. A good overview is presented by Caines, Huang and Malhamé in [8], a detailed probabilistic approach by Carmona and Delarue is presented in [11, 12].

MFGs can be formulated as parabolic optimal control problems (under certain conditions on the cost). This connection can be used to construct approximations to MFGs. Degond, Liu and Ringhofer proposed a so-called best reply strategy (BRS) in [21]. It can be derived from the corresponding explicit in time discretisation of the respective optimal control problem, as in [3, 20]. More recently it has been derived in [5] through considering a discounted optimal control problem and taking the discount factor to  $\infty$ . Specifically, in [3, 20] the following cost functional

$$J^{\Delta t}(\alpha; m) = \mathbb{E} \left[ \int_t^{t+\Delta t} \left( \frac{\alpha_s^2}{2} + \frac{1}{\Delta t} h(X_s, m(X_s)) \right) ds \right],$$

is analysed as  $\Delta t \rightarrow 0$ , where  $m$  is a probability measure that is absolutely continuous with respect to the Lebesgue measure,  $\alpha = (\alpha_t)_{t \geq 0}$  is a control process, and  $X_t$  is a controlled stochastic process evolving according to

$$dX_t = \alpha_t dt + \sigma dB_t.$$

The expectation in the cost functional is taken with respect to the controlled process. In [5] the limiting behavior of MFG systems with cost functionals

$$J^\rho(\alpha; m) = \mathbb{E} \left[ \int_0^T \left( \frac{\alpha_s^2}{2} + h(X_s, m(X_s)) \right) e^{-\rho s} ds \right],$$

as the temporal discount factor  $\rho$  tends to infinity is considered. In both cases the resulting dynamics depend instantaneously on the cost function  $h$ . Hence, in these limiting systems, agents do not anticipate future dynamics, which contrasts with MFG approaches where the future behaviour of the system is taken into account.

As far as the authors are aware, no systematic analysis and comparison of the two approaches has been done. However, the BRS is computationally less expensive and therefore more attractive in applications. Therefore it is important to understand under which circumstances the use of each model is appropriate and whether there are situations where the two models are comparable. This paper is a first step towards analysing the similarities and differences between the two models in a systematic way. The comparisons in this paper are restricted to some specific examples, which illuminate the characteristic differences that can be expected between the MFG and BRS and we provide some reasoning for why such differences occur, based on the modelling differences between the MFG and BRS paradigms

The existence and uniqueness of solutions to stationary MFGs has been studied extensively in previous literature, c.f. [9, 25, 27, 38]. However, apart from a small number of papers e.g. [18, 26], almost all results focus on problems posed on the torus in order to avoid dealing with boundary conditions. In [4] the Dirichlet problem was motivated as a stopping time problem, and it was analysed in [26]. In this paper we consider Neumann boundary conditions, which relate to a no-flux boundary. The only other paper we are aware of that deals with such a situation is [18]. In this paper the authors prove existence of solutions to the MFG problem with non-local dependence on the distribution using a Schauder fixed point argument. They then perturb the solutions to prove existence in the case of a local dependence on the distribution. Other typical methods of proof use continuation methods [23, 25, 28], Schauder's fixed point theorem [7, 19] or variational approaches through energy minimisation problems [16, 22]. In our proof we exploit the linear-quadratic nature of the control. This was done in the time-dependent case in [32] where the problem was reduced to a forward-backward system of heat equations, but our method has not been considered in the stationary case. Our result complements the only other result for Neumann boundary conditions [18]. On the one hand the Hamiltonian used in [18] is more general than ours, however the regularity assumptions and the form of nonlinearity  $h$  required in [18] is relaxed under our assumptions.

Due to assumption  $(A_2)$ , which states that the running cost  $h$  is an increasing function of density, we are in the setting of monotone stationary MFGs. Existence and uniqueness of such MFGs has been studied extensively by Gomes and collaborators in a number of papers, e.g. [22, 27, 30]. Although the

setting in these papers focusses on domains with periodic boundary conditions, it is worth mentioning the types of techniques used and how they compare to the method in this paper. In [27] a Hopf–Cole transformation is used to prove existence and uniqueness of minimisers of an energy functional related to a specific case of an MFG with periodic boundary conditions and a cost  $h$  that is logarithmic in the density. The concepts used in our existence and uniqueness proof are similar to those used in [27], though we are able to generalise the density dependence and consider Neumann boundary value problems. In [28] the results of [27] are extended using a continuation method. They were further improved in [25] where a combination of a continuation method and Minty’s method is used. In both cases, the methods allow the authors to perturb a problem for which existence and uniqueness is known to prove existence and uniqueness of the problem of interest. The methods used there and in many subsequent works (e.g. [26]) rely on monotonicity properties of the operators. In our work presented here monotonicity similarly plays a central role in proving existence and uniqueness, **both through** the use of the maximum principle to prove existence and uniqueness for strictly increasing functions  $h$ , and through the ability to uniformly perturb an increasing function into a strictly increasing function **with** the addition of a logarithmic congestion term.

Our non-linear stationary BRS model (8) is an example of a stationary non-linear Fokker–Planck equation. Existence and uniqueness of solutions to non-linear Fokker–Planck equations have been studied extensively (see for example [13, 15] and references therein). Many results (e.g. in [13, 17, 42]) focus on non-local, non-linear terms i.e. they consider Fokker–Planck equations of the form

$$(1) \quad -\frac{\sigma^2}{2}\nabla^2 m - \nabla \cdot (m\nabla W * m) = 0,$$

with suitable boundary conditions. Here  $\nabla W * m = \int_{\Omega} \nabla W(x-y)m(y) dy$  is the usual convolution operator. For our model, we consider a local function of density  $h = h(x, m(x))$ , rather than a convolution term. In this case there are a number of results of existence and uniqueness of solutions to the stationary model, as well as convergence of the dynamic model to the stationary version, see e.g. [6, 14, 15, 39]. These papers all consider the term  $h$  to be either independent of  $x$  i.e.  $h = h(m)$ , or of the form  $h = h_1(x) + h_2(m)$ . So our result extends this case to more general local functions of the density. In previous literature the proof for the local case, as in [39], relies on a related energy functional, for which minimisers can be proven to exist and be unique. Then these minimisers are also solutions to the Fokker–Planck equation. Our result takes a different approach, one that is more closely related to the case with a convolution term, as in [13]. For the non-local case (1) it has been frequently shown (c.f. [13, 41]) that solutions of PDE (1) are equivalent to fixed points of a non-linear map

$$T(m) = \frac{1}{Z} e^{-\frac{\sigma^2}{2} W * m}, \quad \text{where } Z = \int_{\Omega} e^{-\frac{\sigma^2}{2} W * m} dx.$$

We approach existence and uniqueness of solutions to our PDE (8) in a similar vein, considering solutions to the implicit equation (11). While the proof in [13] relies on Schauder’s fixed point theorem, we take advantage of  $h$  being a local function of density, so instead we use the implicit function theorem and intermediate value theorem to prove our result. **It should be noted that our technique used to prove existence and uniqueness relies on monotonicity arguments, and on the quadratic nature of the Hamiltonian. As such, the technique may not be readily extendable to more general situations.**

This paper is organised as follows. We start by briefly introducing the time-dependent MFG and BRS models in Section 2.1, following a more detailed derivation presented in [3, 20]. We then describe how the dynamic problems relate to the stationary case. In Section 3 we present a proof of existence and uniqueness for the stationary BRS and MFG. Both proofs use similar arguments for proving existence and uniqueness, relying on the observation that both models involve a stationary Fokker–Planck equation with integral constraints. In Section 4 we describe an explicit solution to the MFG and BRS model. The explicit solution allows us to analyse **how the problem-specific parameters affect the similarity between the MFG and BRS**. In Section 5 we illustrate the different behavior of solutions to both models with numerical simulations, **and provide an explanation for the differences based on each model’s construction**. Finally, in Section 6 we conclude with summarising the implications of the results found, specifically what they tell us about the similarities and differences between the two models. We also briefly comment on future directions for research.

**2.1. The dynamic MFG problem and the corresponding BRS.** First we briefly review the underlying modeling assumptions of MFGs and the respective BRS models. For the ease of presentation we consider a quadratic cost on the control and restrict ourselves to the  $d$ -dimensional torus  $\mathbb{T}^d$  in the introduction. However, we will consider bounded domains with Neumann boundary conditions from Section 3. Note that some of the following arguments have not been proven for such a set-up. However it

is not unreasonable to assume that the following results extend naturally to the bounded domain case. Consider a distribution of agents **that** is absolutely continuous with respect to the Lebesgue measure. We denote the density of the distribution by a function  $m : \mathbb{T}^d \rightarrow [0, \infty)$ . We **assume** a representative agent **from this distribution**, with state  $X_t \in \mathbb{T}^d$ , **moves** according to the following SDE:

$$\begin{aligned} dX_t &= \alpha_t dt + \sigma dB_t \\ \mathcal{L}(X_0) &= m_0, \end{aligned}$$

where  $\mathcal{L}(X_0)$  denotes the law of the random variable  $X_0$ ,  $m_0$  is a given initial distribution of agents,  $\sigma \in (0, \infty)$  denotes the size of idiosyncratic noise in the model and  $B_t$  is a  $d$ -dimensional Wiener process. The function  $\alpha_t : [0, T] \rightarrow \mathbb{T}^d$  is a control chosen by the representative agent as a result of an optimisation problem, from a set of admissible controls  $\alpha \in \mathcal{A}$ . The representative agent takes the distribution of other agents,  $m$ , to be given and attempts to optimise the following functional

$$(2) \quad J(\alpha; m) = \mathbb{E} \left[ \int_0^T \left( \frac{\alpha_s^2}{2} + h(X_s, m(X_s)) \right) ds \right].$$

The functional  $J$  consists of a quadratic cost **on** the control  $\alpha_t$  and a density dependent cost function  $h : \mathbb{T}^d \times (0, \infty) \rightarrow \mathbb{R}$  over a finite time horizon  $T \in (0, \infty)$ . Then the optimal cost trajectory  $u(x, t)$  is

$$u(x, t) = \inf_{\alpha \in \mathcal{A}} \mathbb{E} \left[ \int_t^T \left( \frac{\alpha_s^2}{2} + h(X_s, m(X_s)) \right) ds \middle| X_t = x \right].$$

The optimal control is given in terms of  $u$  by  $\alpha_t^* = -\nabla u(X_t, t)$ . The optimal cost trajectory evolves backwards in time according to

$$\begin{aligned} \partial_t u &= \frac{|\nabla u|^2}{2} - h(x, m) - \frac{\sigma^2}{2} \nabla^2 u \\ u(x, T) &= 0. \end{aligned}$$

We complete the model by assuming all agents act in the same way as the representative agent, and so the backward PDE is coupled to a forward Fokker–Planck PDE describing the evolution of agents. The full MFG model is given by

$$(3a) \quad \partial_t u = \frac{|\nabla u|^2}{2} - h(x, m) - \frac{\sigma^2}{2} \nabla^2 u$$

$$(3b) \quad \partial_t m = \nabla \cdot [m \nabla u] + \frac{\sigma^2}{2} \nabla^2 m$$

$$(3c) \quad m(x, 0) = m_0$$

$$(3d) \quad u(x, T) = 0.$$

Following the approach in [3, 20], the corresponding BRS model arises through considering a rescaled cost functional over a short, rolling time horizon

$$J^{\Delta t}(\alpha; m) = \mathbb{E} \left[ \int_t^{t+\Delta t} \left( \frac{\alpha_s^2}{2} + \frac{1}{\Delta t} h(X_s, m(X_s)) \right) ds \middle| X_t = x \right].$$

Going through a similar procedure to the MFG problem, approximating the result up to  $O(\Delta t)$  and taking the limit  $\Delta t \rightarrow 0$ , the optimal control is given by  $\alpha_t = -[\nabla h(x, m(x))]_{x=X_t}$ . **The model is completed** by assuming all agents act in the same way as the representative agent. Then the distribution of agents evolves according to the Fokker–Planck equation

$$(4a) \quad \partial_t m = \nabla \cdot [(\nabla h(x, m(x))) m] + \frac{\sigma^2}{2} \nabla^2 m$$

$$(4b) \quad m(x, 0) = m_0.$$

In (4) the dynamics of agents is influenced **only** by the current agent density. Hence the anticipation behavior, which is characteristic **of the MFG model**, is ‘lost’. Only the current state drives the dynamics. We shall refer to equation (4) as the BRS strategy in the following.

**2.2. From the dynamic problems to the stationary case.** For the MFG the interpretation of the stationary problem is slightly subtle because in the dynamic case we are considering a problem set on a fixed time horizon so we cannot consider the stationary problem by setting  $\partial_t u, \partial_t m = 0$  and interpreting it as the long-time behaviour of the dynamic case. Instead we follow the work by Cardaliaguet, Lasry,

Lions and Porretta [10]. To highlight the dependence of the MFG on the time horizon  $T$  we use the notation  $\bar{u}^T, \bar{m}^T$  for solutions satisfying (3). Then we define the rescaled functions  $u^T$  and  $m^T$  by

$$u^T(x, t) = \bar{u}^T(x, tT), \quad m^T(x, t) = \bar{m}^T(x, tT).$$

Then Theorem 1.2 in [10] states that under some mild assumptions on the data we have that as  $T \rightarrow \infty$ :

$$\begin{aligned} u^T - \int_{\mathbb{T}^d} u^T dy &\rightarrow u, \quad \text{in } L^2(\mathbb{T}^d \times (0, 1)) \\ \frac{1}{T} u^T &\rightarrow (1-t)\lambda, \quad \text{in } L^2(\mathbb{T}^d \times (0, 1)) \\ m^T &\rightarrow m, \quad \text{in } L^p(\mathbb{T}^d \times (0, 1)), \end{aligned}$$

where  $p$  depends on the space dimension  $d$ . **Furthermore**, the triple  $(m, u, \lambda) \in C^2(\mathbb{T}^d) \times C^2(\mathbb{T}^d) \times \mathbb{R}$  satisfies the following stationary problem

$$(5a) \quad -\frac{\sigma^2}{2} \nabla^2 m - \nabla \cdot (m \nabla u) = 0$$

$$(5b) \quad -\frac{\sigma^2}{2} \nabla^2 u + \frac{|\nabla u|^2}{2} - h(x, m) + \lambda = 0$$

$$(5c) \quad \int_{\mathbb{T}^d} m dx = 1$$

$$(5d) \quad \int_{\mathbb{T}^d} u dx = 0.$$

The corresponding stationary BRS model is obtained by setting  $\partial_t m = 0$  in (4). Hence we have

$$(6a) \quad \nabla \cdot [(\nabla h(x, m(x))) m] + \frac{\sigma^2}{2} \nabla^2 m = 0$$

$$(6b) \quad \int_{\mathbb{T}^d} m dx = 1.$$

Equation (6) can be understood as either the long-time behaviour of the dynamic BRS or, under suitable convexity conditions (c.f. [39]), a competitive equilibrium distribution of the following minimisation problem

$$(7) \quad \min \mathbb{E} [h(X_t, m(X_t))].$$

By competitive equilibrium we mean a stationary distribution  $m$  for which  $\mathbb{E} [h(X, m(X))]$  is minimised when  $\mathcal{L}(X) = m$ .

### 3. EXISTENCE AND UNIQUENESS OF STATIONARY SOLUTIONS

In this section we show that the MFG (5) and the BRS (6) admit unique solutions on a bounded domain  $\Omega$  with no flux boundary conditions. We make the following assumptions on the function  $h(x, m) : \Omega \times (0, \infty) \rightarrow \mathbb{R}$  and the domain  $\Omega$ :

(A<sub>1</sub>)  $\Omega \subset \mathbb{R}^d$  is an open bounded set with a  $C^{2,\alpha}$  boundary, for some  $\alpha \in (0, 1)$  and  $d \geq 1$ .

(A<sub>2</sub>)  $h(x, \cdot)$  is an increasing function for every  $x \in \Omega$ .

(A<sub>3</sub>) There exists a continuous function  $g : (0, \infty) \rightarrow [0, \infty)$  such that  $\sup_{x \in \Omega} |h(x, m)| \leq g(m)$  for every  $m \in (0, \infty)$ .

Since  $\Omega$  is bounded, we can now define  $|\Omega| = \int_{\Omega} dx$ , where this integral is with respect to the standard Lebesgue measure. Furthermore, we denote the unit outer normal vector by  $\nu$ .

For the BRS we **also** assume:

$$(BRS_1) \quad h \in C^2(\Omega \times (0, \infty)) \cap C^1(\bar{\Omega} \times (0, \infty)).$$

$$(BRS_2) \quad \text{There exists a continuous function } f : (0, \infty) \rightarrow [0, \infty) \text{ such that } \sup_{x \in \Omega} |\nabla_x h(x, m)| \leq f(m) \text{ for every } m \in (0, \infty).$$

While for the MFG we will assume:

$$(MFG_1) \quad h \in C(\Omega \times (0, \infty))$$

$$(MFG_2) \quad \lim_{m \rightarrow 0} \sup_{x \in \Omega} h(x, m) < \inf_{x \in \Omega} h\left(x, \frac{1}{|\Omega|}\right).$$

$$(MFG_3) \quad \sup_{x \in \Omega} h\left(x, \frac{1}{|\Omega|}\right) < \lim_{m \rightarrow \infty} \inf_{x \in \Omega} h(x, m).$$

**Discussion of assumptions:** Since we are interested in classical solutions, generally in  $C^2(\Omega) \cap C^1(\bar{\Omega})$ , the above assumptions on the cost and domain ensure sufficient regularity and boundedness of  $h$ . It is worth mentioning why we need  $h$  to be increasing. This assumption of an increasing function can be related to “crowd aversion”. When  $h$  is increasing in  $m$  then areas of high density are more highly penalised than low density areas, in the optimisation problem related to the MFG and BRS (see Section 2.1). This prevents “accumulation points” occurring, where higher density is preferable and a Dirac delta might be introduced into the solution. As well as being a problem for regularity, the position of the Dirac deltas would be sensitive on the data and so uniqueness could not be guaranteed.

Assumptions (MFG<sub>2</sub>) and (MFG<sub>3</sub>) are also less intuitive than the rest of the assumptions. The MFG problem has two integral constraints related to it. We prove that these constraints can be satisfied using the intermediate value theorem. In doing so we show that two functions  $m_1, m_2$  exist such that  $m_1(x) \leq \frac{1}{|\Omega|} \leq m_2(x)$  for every  $x \in \Omega$ . The existence of these functions is guaranteed if assumptions (MFG<sub>2</sub>) and (MFG<sub>3</sub>) hold. As it may not be initially clear what kind of function  $h$  satisfies our requirements, some sufficient conditions if  $h(x, m) = h_1(x) + h_2(m)$  are:

- $h_1 \in C^2(\Omega) \cap C^1(\bar{\Omega})$
- $h_2 \in C^2((0, \infty))$
- $h_2$  is increasing
- $\lim_{m \rightarrow 0} h_2(m) < h_2\left(\frac{1}{|\Omega|}\right) + \inf_{x \in \Omega} h_1(x)$
- $\lim_{m \rightarrow \infty} h_2(m) > h_2\left(\frac{1}{|\Omega|}\right) + \sup_{x \in \Omega} h_1(x)$

**3.1. Best Reply Strategy.** We start by defining the notion of classical solutions we are aiming for.

**Definition 3.1.** Let assumptions (A<sub>1</sub>)–(A<sub>3</sub>) and (BRS<sub>1</sub>)–(BRS<sub>2</sub>) be satisfied. Then the stationary BRS boundary value problem is to find a function  $m : \Omega \rightarrow (0, \infty)$  satisfying

$$\begin{aligned}
 (8a) \quad & m \in C^2(\Omega) \cap C^1(\bar{\Omega}) \\
 (8b) \quad & -\frac{\sigma^2}{2} \nabla^2 m - \nabla \cdot (m \nabla [h(x, m)]) = 0, \quad x \in \Omega \\
 (8c) \quad & -\frac{\sigma^2}{2} \nabla m \cdot \nu - m \nabla [h(x, m)] \cdot \nu = 0, \quad x \in \partial\Omega \\
 (8d) \quad & \int_{\Omega} m \, dx = 1.
 \end{aligned}$$

Throughout this subsection we will assume (A<sub>1</sub>)–(A<sub>3</sub>) and (BRS<sub>1</sub>)–(BRS<sub>2</sub>) hold.

**Lemma 3.2.** For any  $Z \in (0, \infty)$  there exists a unique  $m_Z : \Omega \rightarrow (0, \infty)$  such that

$$(9) \quad m_Z(x) = \frac{1}{Z} e^{-\frac{2}{\sigma^2} h(x, m_Z(x))}.$$

Furthermore,  $m_Z \in C^2(\Omega) \cap C^1(\bar{\Omega})$ .

*Proof.* Fix  $Z \in (0, \infty)$  and  $x \in \Omega$ . Consider  $G_{Z,x} : (0, \infty) \rightarrow \mathbb{R}$  given by

$$G_{Z,x}(m) = \frac{1}{Z} e^{-\frac{2}{\sigma^2} h(x, m)} - m.$$

This is a strictly decreasing function of  $m$ . Furthermore, since  $h$  is increasing and continuous with respect to  $m$  and  $\sup_{x \in \Omega} h(x, m) \leq g(m)$ , we must have  $\lim_{m \rightarrow 0} \sup_{x \in \Omega} h(x, m) \leq \sup_{x \in \Omega} h(x, 1) \leq g(1) < \infty$ . So we get the following limit inequality as  $m \rightarrow 0$ , which holds uniformly in  $x$ :

$$\lim_{m \rightarrow 0} G_{Z,x}(m) > 0.$$

So there exists some  $\epsilon > 0$ , independent of  $x$ , such that  $G_{Z,x}(\epsilon) > 0$ . Furthermore, after defining a constant  $C = \inf_{x \in \Omega} h(x, \epsilon)$ , it is clear that

$$G_{Z,x} \left( \frac{1}{Z} e^{-\frac{2}{\sigma^2} C} + \epsilon \right) \leq \frac{1}{Z} e^{-\frac{2}{\sigma^2} h(x, \epsilon)} - \frac{1}{Z} e^{-\frac{2}{\sigma^2} C} - \epsilon \leq -\epsilon < 0.$$

Therefore, by the intermediate value theorem and strict monotonicity of  $G_{Z,x}$ , there exists a unique  $m = m_Z(x) > 0$  such that  $G_{Z,x}(m_Z(x)) = 0$ . Hence the first result follows. In order to show the regularity requirement that  $m_Z \in C^2(\Omega) \cap C^1(\bar{\Omega})$ , we need to show

- (1)  $m_Z \in C^2(\Omega)$ .
- (2) For any  $x \in \partial\Omega$ ,  $\lim_{y \rightarrow x, y \in \Omega} m_Z(y)$  exists.
- (3) For any  $x \in \partial\Omega$ ,  $\lim_{y \rightarrow x, y \in \Omega} \nabla m_Z(y)$  exists.

The assertion that  $m_Z \in C^2(\Omega)$  follows from the implicit function theorem. For the implicit function theorem to hold, we require that  $G_{Z,x}(m)$  is a  $C^2$  function with respect to  $x$  and  $m$  at  $(x, m_Z(x))$  and that  $G'_{Z,x}(m_Z(x)) \neq 0$ . The first requirement is true from our assumption that  $h \in C^2(\Omega \times (0, \infty))$ , the second requirement is true since

$$G'_{Z,x}(m) = -\frac{2}{\sigma^2 Z} \partial_m h(x, m) e^{-\frac{2}{\sigma^2} h(x, m)} - 1 \leq -1 < 0.$$

To prove that  $\lim_{y \rightarrow x, y \in \Omega} m(y)$  exists for every  $x \in \partial\Omega$ , it is enough to show  $m_Z$  is uniformly Lipschitz in  $\Omega$ . Since  $m_Z \in C^2(\Omega)$ , it is enough to show  $\|\nabla m_Z\|_\infty < \infty$ . Note that  $C, \epsilon$  defined above are independent of  $x$ , so we must have  $\epsilon \leq \|m_Z\|_\infty \leq \frac{1}{2} e^{-\frac{2}{\sigma^2} C} + \epsilon < \infty$ . Then by differentiating the implicit formula for  $m_Z$  we get

$$(10) \quad \nabla m_Z = \frac{-2m_Z \nabla_x h(x, m_Z)}{\sigma^2 + 2m_Z \partial_m h(x, m_Z)}.$$

But  $\partial_m h \geq 0$  since  $h$  is increasing, also  $m_Z$  is uniformly bounded as seen above. Similarly, using  $f$  from assumption (BRS<sub>2</sub>),  $\|\nabla_x h(\cdot, m_Z(\cdot))\|_\infty < \infty$ , so  $\|\nabla m_Z\|_\infty < \infty$ .

To prove the final assertion that  $\lim_{y \rightarrow x, y \in \Omega} \nabla m_Z(y)$  exists for any  $x \in \partial\Omega$ , note **that** the formula for  $\nabla m_Z$  is given by (10). Since  $m_Z \in C^0(\bar{\Omega})$ ,  $\|m_Z\|_\infty < \infty$ ,  $\nabla_x h, \partial_m h \in C^0(\bar{\Omega} \times (0, \infty))$  and  $\sigma^2 + 2m_Z \partial_m h(x, m_Z) \geq \sigma^2$ , then the right hand side of (10) has **a** limit as  $y \rightarrow x$  for every  $x \in \partial\Omega$ . Hence,  $\nabla m_Z$  does as well.  $\square$

**Definition 3.3.** Let  $m_Z$  be given by (9). Then we define the following function  $\Phi : (0, \infty) \rightarrow (0, \infty)$

$$\Phi(Z) = \int_{\Omega} m_Z \, dx.$$

**Lemma 3.4.** *There exists  $\bar{Z}, Z$  such that  $\Phi(\bar{Z}) \geq 1$  and  $\Phi(Z) \leq 1$ .*

*Proof.* Take  $C_1 = \inf_{x \in \Omega} h(x, \frac{1}{|\Omega|})$ . So,  $C_1 \in [-g(\frac{1}{|\Omega|}), g(\frac{1}{|\Omega|})]$ . Then, with  $Z = |\Omega| e^{-\frac{2}{\sigma^2} C_1} \in (0, \infty)$ , we have

$$G_{Z,x}\left(\frac{1}{|\Omega|}\right) \leq 0 \text{ for every } x \in \Omega.$$

Hence,  $m_Z(x) \leq \frac{1}{|\Omega|}$  because  $G_{Z,x}$  is a strictly decreasing function. So

$$\Phi(Z) \leq \|m_Z\|_\infty |\Omega| \leq 1.$$

We can similarly find  $\bar{Z}$  by taking  $C_2 = \sup_{x \in \Omega} h(x, \frac{1}{|\Omega|})$ . So,  $C_2 \in [-g(\frac{1}{|\Omega|}), g(\frac{1}{|\Omega|})]$ . Then, with  $\bar{Z} = |\Omega| e^{-\frac{2}{\sigma^2} C_2} \in (0, \infty)$ , we have

$$G_{\bar{Z},x}\left(\frac{1}{|\Omega|}\right) \geq 0 \text{ for every } x \in \Omega.$$

Hence,  $m_{\bar{Z}}(x) \geq \frac{1}{|\Omega|}$  because  $G_{\bar{Z},x}$  is a strictly decreasing function. So

$$\Phi(\bar{Z}) \geq \|m_{\bar{Z}}\|_\infty |\Omega| \geq 1.$$

$\square$

**Lemma 3.5.** *There exists a unique  $Z^* \in (0, \infty)$  such that  $\Phi(Z^*) = 1$ .*

*Proof.* If  $\Phi$  is continuous and strictly decreasing the intermediate value theorem and the Lemma 3.4 give the result. We start by proving that  $\Phi$  is strictly decreasing. First note that if  $m_Z(x)$  is strictly decreasing in  $Z$  for every  $x$ , then  $\Phi$  must be strictly decreasing, because  $m_Z$  is continuous with respect to  $x$ . Take  $Z_1 < Z_2$ . Then  $m_{Z_1}(x)$  satisfies

$$\frac{1}{Z_1} e^{-\frac{2}{\sigma^2} h(x, m_{Z_1}(x))} - m_{Z_1}(x) = 0.$$

So,

$$\frac{1}{Z_2} e^{-\frac{2}{\sigma^2} h(x, m_{Z_1}(x))} - m_{Z_1}(x) < \frac{1}{Z_1} e^{-\frac{2}{\sigma^2} h(x, m_{Z_1}(x))} - m_{Z_1}(x) = 0.$$

Hence,  $G_{Z_2, x}(m_{Z_1}(x)) < 0$ . Then,  $m_{Z_2}(x) < m_{Z_1}(x)$  for all  $x \in \Omega$  since  $G_{Z, x}$  is a strictly decreasing function. To show that  $\Phi$  is continuous at  $Z \in (0, \infty)$ , take  $\epsilon < Z' < Z$ . Then

$$\begin{aligned} |\Phi(Z) - \Phi(Z')| &= \Phi(Z') - \Phi(Z) = \int_{\Omega} m_{Z'}(x) dx - \Phi(Z) \\ &= \frac{Z}{Z'} \int_{\Omega} \frac{1}{Z} e^{-h(x, m_{Z'}(x))} dx - \Phi(Z) \\ &\leq \frac{Z}{Z'} \int_{\Omega} \frac{1}{Z} e^{-h(x, m_Z(x))} dx - \Phi(Z) \leq \frac{\Phi(Z)}{\epsilon} (Z - Z') \leq \frac{\Phi(\epsilon)}{\epsilon} (Z - Z'). \end{aligned}$$

By exchanging  $Z$  and  $Z'$ , we can similarly show the analogous result for  $\epsilon < Z < Z'$ , therefore  $\Phi$  is locally Lipschitz and hence continuous.  $\square$

**Theorem 3.6.** *There exists a unique solution  $m : \Omega \rightarrow (0, \infty)$  to the stationary BRS (8).*

*Proof.* Take  $m(x) = m_{Z^*}(x)$ , with  $Z^*$  defined as in Lemma 3.5. Then from Lemmas 3.2 and 3.5 we have shown there exists a unique  $m : \Omega \rightarrow (0, \infty)$  satisfying

$$(11a) \quad m \in C^2(\Omega) \cap C^1(\bar{\Omega})$$

$$(11b) \quad \text{There exists } Z \in (0, \infty) \text{ such that } m = \frac{1}{Z} e^{-\frac{2}{\sigma^2} h(x, m)}, \quad x \in \Omega$$

$$(11c) \quad \int_{\Omega} m dx = 1.$$

Now, for any  $m : \Omega \rightarrow (0, \infty)$  we can define  $\phi(m)$  by  $\phi(m) = e^{-\frac{2}{\sigma^2} h(x, m)}$ . Suppose  $m$  is a solution to (8), then  $\frac{m}{\phi(m)} = m e^{\frac{2}{\sigma^2} h(x, m)}$ , and so  $\frac{m}{\phi(m)} \in H^1(\Omega)$  because  $m \in C^1(\bar{\Omega})$ ,  $h \in C^1(\bar{\Omega} \times (0, \infty))$  and  $h$  is increasing in  $m$ . Therefore, a solution to (8) is equivalent to a solution of

$$(12a) \quad m \in C^2(\Omega) \cap C^1(\bar{\Omega})$$

$$(12b) \quad \frac{m}{\phi(m)} \in H^1(\Omega)$$

$$(12c) \quad \nabla \cdot \left( \phi(m) \nabla \left( \frac{m}{\phi(m)} \right) \right) = 0, \quad x \in \Omega$$

$$(12d) \quad \phi(m) \nabla \left( \frac{m}{\phi(m)} \right) \cdot \nu = 0, \quad x \in \partial\Omega$$

$$(12e) \quad \int_{\Omega} m dx = 1.$$

Now, if we multiply (12c) by  $\frac{m}{\phi(m)}$ , and integrate over  $\Omega$ , then using Green's formula and the boundary condition (12d) we get

$$0 = \int_{\Omega} \frac{m}{\phi(m)} \nabla \cdot \left( \phi(m) \nabla \left( \frac{m}{\phi(m)} \right) \right) dx = - \int_{\Omega} \phi(m) \left| \nabla \left( \frac{m}{\phi(m)} \right) \right|^2 dx.$$

But  $\phi(m) > 0$  and  $\left| \nabla \left( \frac{m}{\phi(m)} \right) \right|^2 \geq 0$  for every  $x \in \Omega$ . Hence,  $\nabla \left( \frac{m}{\phi(m)} \right) = 0$  for every  $x \in \Omega$ , i.e. if there exists  $Z \in (0, \infty)$  such that  $m = \frac{1}{Z} \phi(m)$ . Conversely, if  $m \in C^2(\Omega) \cap C^1(\bar{\Omega})$  and there exists  $Z \in (0, \infty)$  such that  $m = \frac{1}{Z} \phi(m)$ , then  $m$  satisfies (12a)–(12d). So, a solution of (12) is equivalent to a solution of (11). Therefore, the systems (8) and (11) are equivalent, and existence and uniqueness of solutions to (11) follows from existence and uniqueness of solutions to (8).  $\square$

**3.2. Mean Field Games.** Next we discuss existence and uniqueness of classical solutions to (5), which is defined as follows.

**Definition 3.7.** The stationary MFG boundary value problem is to find  $m : \Omega \rightarrow (0, \infty)$ ,  $u : \Omega \rightarrow \mathbb{R}$  and  $\lambda \in \mathbb{R}$  satisfying the following PDE system

$$\begin{aligned}
(13a) \quad & m \in C^2(\Omega) \cap C^1(\bar{\Omega}) \\
(13b) \quad & u \in C^2(\Omega) \cap C^1(\bar{\Omega}) \\
(13c) \quad & -\frac{\sigma^2}{2} \nabla^2 m - \nabla \cdot (m \nabla u) = 0, \quad x \in \Omega \\
(13d) \quad & -\frac{\sigma^2}{2} \nabla^2 u + \frac{|\nabla u|^2}{2} - h(x, m) + \lambda = 0, \quad x \in \Omega \\
(13e) \quad & -\frac{\sigma^2}{2} \nabla m \cdot \nu = 0, \quad x \in \partial\Omega \\
(13f) \quad & -\nabla u \cdot \nu = 0, \quad x \in \partial\Omega \\
(13g) \quad & \int_{\Omega} m \, dx = 1, \\
(13h) \quad & \int_{\Omega} u \, dx = 0.
\end{aligned}$$

*Remark 3.8.* Here, following the method of Section 3.1, we note that for any  $u \in C^2(\Omega) \cap C^1(\bar{\Omega})$ , a solution  $m$  of (13a), (13c), (13e), (13g) is equivalent to a solution of

$$\begin{aligned}
(14a) \quad & m \in C^2(\Omega) \cap C^1(\bar{\Omega}) \\
(14b) \quad & m = \frac{1}{Z} e^{-\frac{\sigma^2}{2} u}, \quad x \in \Omega \\
(14c) \quad & Z = \int_{\Omega} e^{-\frac{\sigma^2}{2} u} \, dx.
\end{aligned}$$

Then, by the arguments in Section 3.1, a unique  $m$  satisfying (14) exists and is the unique solution to (13a), (13c), (13e), (13g). So from now we only consider that solution.

**Proposition 3.9.** *There exists a unique solution  $(m, u, \lambda) \in [C^2(\Omega) \cap C^1(\bar{\Omega})] \times [C^2(\Omega) \cap C^1(\bar{\Omega})] \times \mathbb{R}$  to the stationary MFG boundary value problem if and only if there exists a unique solution  $(u, \lambda, Z) \in [C^2(\Omega) \cap C^1(\bar{\Omega})] \times \mathbb{R} \times (0, \infty)$  to*

$$\begin{aligned}
(15a) \quad & u \in C^2(\Omega) \cap C^1(\bar{\Omega}) \\
(15b) \quad & -\frac{\sigma^2}{2} \nabla^2 u + \frac{|\nabla u|^2}{2} - h\left(x, \frac{1}{Z} e^{-\frac{\sigma^2}{2} u}\right) + \lambda = 0, \quad x \in \Omega \\
(15c) \quad & -\nabla u \cdot \nu = 0, \quad x \in \partial\Omega \\
(15d) \quad & \int_{\Omega} \frac{1}{Z} e^{-\frac{\sigma^2}{2} u} \, dx = 1, \\
(15e) \quad & \int_{\Omega} u \, dx = 0.
\end{aligned}$$

*Proof.* First, assume a unique solution  $(m, u, \lambda)$  to (13) exists, then **using** remark 3.8,  $m = \frac{1}{Z} e^{-\frac{\sigma^2}{2} u}$ , for  $Z$  satisfying (14c). So, the triple  $(u, \lambda, Z)$  is clearly a solution to (15). Furthermore, suppose another solution  $(u', \lambda', Z')$  to (15) exists. Then  $(m', u', \lambda')$ , with  $m' = \frac{1}{Z'} e^{-\frac{\sigma^2}{2} u'}$ , is a solution to (13). But, since we assumed such solutions are unique,  $(m', u', \lambda') = (m, u, \lambda)$  and hence  $(u', \lambda', Z') = (u, \lambda, Z)$ , so the solution to (15) is unique.

Next, assume that a unique solution  $(u, \lambda, Z)$  to (15) exists. Then, defining  $m = \frac{1}{Z} e^{-\frac{\sigma^2}{2} u}$ ,  $(m, u, \lambda)$  is a solution to (13). Now, suppose  $(m', u', \lambda')$  is another solution then, again using remark 3.8,  $m' = \frac{1}{Z'} e^{-\frac{\sigma^2}{2} u'}$ , where  $Z'$  satisfies (14c). So  $(u', \lambda', Z')$  satisfies (15). By uniqueness  $(u', \lambda', Z') = (u, \lambda, Z)$  and so  $(m, u, \lambda)$  is also unique.  $\square$

**Theorem 3.10.** *There exists a unique solution  $(m, u, \lambda)$  of the MFG system (13).*

*Proof (outline).* First note, as a result of Proposition 3.9, we only need to prove existence and uniqueness of a solution to (15) and existence and uniqueness for the MFG system (13) will follow. The proof is split into the following steps:

- (1) Show that for any pair of constants  $(\lambda, Z)$  there exists a unique solution, denoted by  $u_{\lambda, Z}$ , to (15a), (15b) (see Proposition 3.15)
- (2) Show that for any constant  $Z$  there exists a unique  $\lambda = \lambda(Z)$  such that  $u_{\lambda(Z), Z}$  satisfies (15d) (see Proposition 3.16)
- (3) Show that there exists a unique  $Z = Z^*$  such that  $u_{\lambda(Z^*), Z^*}$  satisfies (15e).

Then  $(u_{\lambda(Z^*), Z^*}, \lambda(Z^*), Z^*)$  is a solution to (15). Uniqueness follows from uniqueness obtained at each step of the proof outlined. We prove step 1 using a variant of the method of upper and lower solutions in the spirit of [40], so that it applies to our case of Neumann boundary conditions. We prove steps 2 and 3 by iteratively using the intermediate value theorem, in a similar manner to the proof of Lemma 3.5. Note that we will first do this for  $h$  which is strictly increasing in  $m$ . Then by considering  $h_\epsilon(x, m) = h(x, m) + \epsilon \log(|\Omega|m)$ , and taking the limit as  $\epsilon \rightarrow 0$  we will prove it in the more general setting when  $h$  is increasing.  $\square$

**Lemma 3.11.** *There exists  $\Lambda_1, \Lambda_2 \in [-\infty, \infty]$ , with  $\Lambda_1 < \Lambda_2$ , such that for every  $\lambda \in (\Lambda_1, \Lambda_2)$  and  $Z > 0$ , there exist two constants  $\underline{u}_{\lambda, Z} \leq 0 \leq \bar{u}_{\lambda, Z}$  satisfying*

$$(16) \quad -h\left(x, \frac{1}{Z}e^{-\frac{\sigma^2}{2}u_{\lambda, Z}}\right) + \lambda \leq 0 \leq -h\left(x, \frac{1}{Z}e^{-\frac{\sigma^2}{2}\bar{u}_{\lambda, Z}}\right) + \lambda.$$

*Proof.* Take  $\Lambda_1 = \lim_{m \rightarrow 0} \sup_{x \in \Omega} h(x, m)$  and  $\Lambda_2 = \lim_{m \rightarrow \infty} \inf_{x \in \Omega} h(x, m)$ . First  $\Lambda_1 < \Lambda_2$  by combining assumptions (MFG<sub>2</sub>) and (MFG<sub>3</sub>). Moreover, since  $h$  is continuous and increasing in  $m$ , for any  $\lambda \in (\Lambda_1, \Lambda_2)$  there exists  $M_\lambda^1, M_\lambda^2 \in (0, \infty)$  such that  $h(x, m) \leq \lambda$  for all  $(x, m) \in \Omega \times (0, M_\lambda^1]$ , and similarly  $h(x, m) \geq \lambda$  for all  $(x, m) \in \Omega \times [M_\lambda^2, \infty)$ . We define the upper and lower constants for  $\lambda \in (\Lambda_1, \Lambda_2)$  as

$$\begin{aligned} \bar{u}_{\lambda, Z} &= \max\left(-\frac{\sigma^2}{2} \log Z M_\lambda^1, 0\right) \\ \underline{u}_{\lambda, Z} &= \min\left(-\frac{\sigma^2}{2} \log Z M_\lambda^2, 0\right). \end{aligned}$$

Then clearly

$$-h\left(x, \frac{1}{Z}e^{-\frac{\sigma^2}{2}\bar{u}_{\lambda, Z}}\right) + \lambda = -h(x, \min(M_\lambda^1, 1)) + \lambda \geq -h(x, M_\lambda^1) + \lambda \geq 0,$$

while the reverse inequality is true for  $\underline{u}_{\lambda, Z}$ . Hence,  $\bar{u}_{\lambda, Z}, \underline{u}_{\lambda, Z}$  are the required upper and lower constants.  $\square$

**Proposition 3.12.** *Define  $C^{2, \tau}(\bar{\Omega})$  as the set of functions  $u \in C^2(\bar{\Omega})$  whose second partial derivatives are all Hölder continuous, with exponent  $\tau$ , on  $\bar{\Omega}$ . Assume  $h$  is strictly increasing with respect to  $m$ . Then, for every  $\lambda \in (\Lambda_1, \Lambda_2)$  and every  $Z \in (0, \infty)$ , there exists a unique function,  $u_{\lambda, Z} \in C^{2, \tau}(\bar{\Omega}) \subset C^2(\Omega) \cap C^1(\bar{\Omega})$  for some  $\tau \in (0, 1)$ , which satisfies (15a)–(15c). Furthermore,  $\underline{u}_{\lambda, Z} \leq u_{\lambda, Z} \leq \bar{u}_{\lambda, Z}$ .*

*Proof.* Existence is an application of Corollary 2.9 in [40], which states that a solution  $u_{\lambda, Z} \in C^{2, \tau}(\bar{\Omega})$  to (15a)–(15c) exists provided the following properties hold:

- (1) There exist constants  $\underline{u}_{\lambda, Z} \leq 0 \leq \bar{u}_{\lambda, Z}$  satisfying (16) for every  $x \in \bar{\Omega}$ .
- (2) There exists a continuous function  $f : [0, \infty) \rightarrow [0, \infty)$  such that the following inequality holds for every  $(x, u, p) \in \bar{\Omega} \times \mathbb{R} \times \mathbb{R}^d$

$$\left| \frac{|p|^2}{2} - h\left(x, \frac{1}{Z}e^{-\frac{\sigma^2}{2}u}\right) + \lambda \right| \leq f(|u|)(1 + |p|^2).$$

Property 1 is true from Lemma 3.11. Property 2 can be shown to be true by taking

$$f(u) = \max\left(\frac{1}{2}, |\lambda| + \max\left[g\left(\frac{1}{Z}e^{-\frac{\sigma^2}{2}u}\right), g\left(\frac{1}{Z}e^{\frac{\sigma^2}{2}u}\right)\right]\right),$$

where  $g$  is defined in assumption (A<sub>3</sub>).

We prove uniqueness using the strong maximum principle and Hopf's Lemma as stated in [24] (Section 6.4.2. pp. 330–333). Suppose there are two solutions,  $u_1, u_2 \in C^2(\Omega) \cap C^1(\bar{\Omega})$  to (15a), (15b). Define  $a = \nabla(u_1 + u_2)$ . Then  $a \in L^\infty(\bar{\Omega})$ . Now, suppose  $u_1 \neq u_2$  and define  $v = u_1 - u_2$ . Then,  $v$  must attain

its maximum at some point  $\bar{x} \in \bar{\Omega}$ . First, suppose  $\bar{x} \in \Omega$ . Then, there exists an open bounded and connected region  $V$ , such that  $V \subset \Omega$ ,  $\bar{x} \in V$  and  $v > 0$  for all  $x \in V$ . Hence, since  $h(x, \cdot)$  is increasing,

$$-\frac{\sigma^2}{2} \nabla^2 v + \frac{1}{2} a \cdot \nabla v \leq 0, \quad \text{for every } x \in V.$$

So, by the strong maximum principle,  $v$  is constant in  $V$ . Therefore,

$$h\left(x, \frac{1}{Z} e^{-\frac{2}{\sigma^2} u_1(x)}\right) = h\left(x, \frac{1}{Z} e^{-\frac{2}{\sigma^2} u_2(x)}\right) \quad \text{for every } x \in V.$$

So  $u_1 = u_2$  in  $V$  because  $h$  is strictly increasing, which leads to a contradiction. **Therefore, the only other case to consider is**  $\bar{x} \in \partial\Omega$  and  $v(x) < v(\bar{x})$  for every  $x \in \Omega$ . Hence, by Hopf's Lemma (which we can use because  $\partial\Omega$  is  $C^2$ )  $\frac{\partial v}{\partial \nu}|_{\bar{x}} > 0$ , but by the boundary condition (15c),  $\frac{\partial v}{\partial \nu} = \frac{\partial u_1}{\partial \nu} - \frac{\partial u_2}{\partial \nu} = 0$ . This again leads to a contradiction. Therefore,  $u_1 = u_2$  and the solution is unique.  $\square$

*Remark 3.13.* It should be noted that the same method to prove uniqueness can be used to prove that  $u_{\lambda_1, Z} \geq u_{\lambda_2, Z}$  for all  $\lambda_1 \leq \lambda_2$

**Lemma 3.14.** *Assume  $h$  is strictly increasing with respect to  $m$ . Then, for every  $x \in \Omega$ ,  $u_{\lambda, Z}(x)$  is decreasing with respect to  $\lambda$  and  $Z$ .*

*Proof.* In Light of remark 3.13, we need only to prove  $u_{\lambda, Z_1} \geq u_{\lambda, Z_2}$  for all  $Z_1 \leq Z_2$ . However, by substitution,  $u = u_{\lambda, Z_1} - \frac{\sigma^2}{2} \log \frac{Z_2}{Z_1}$  satisfies (15a)–(15c) with  $Z = Z_2$ . So, by uniqueness of solutions to this PDE proved in Proposition 3.12

$$u_{\lambda, Z_2} = u \leq u_{\lambda, Z_1}.$$

$\square$

**Proposition 3.15.** *Define  $\Phi : (\Lambda_1, \Lambda_2) \times (0, \infty) \rightarrow L^\infty(\Omega)$  by*

$$\Phi(\lambda, Z) = u_{\lambda, Z},$$

*where  $u_{\lambda, Z}$  is the unique solution to (15a)–(15c) as found in the Proposition 3.12. Assume  $h$  is strictly increasing with respect to  $m$ . Then  $\Phi$  is continuous (with respect to **the**  $L^\infty$  norm).*

*Proof.* We will prove  $\Phi$  is sequentially continuous. Let  $(\lambda_n, Z_n)$  be a sequence in  $(\Lambda_1, \Lambda_2) \times (0, \infty)$ , **which** converges to  $(\lambda, Z) \in (\Lambda_1, \Lambda_2) \times (0, \infty)$ . Consider two sequences:  $(\lambda_n^{(i)}, Z_n^{(i)})$  for  $i = 1, 2$ , which **will be used** to sandwich our original sequence. **The sequences are defined term-wise by**

- (1)  $\lambda_n^{(1)} = \inf_{j \geq n} \lambda_j$
- (2)  $\lambda_n^{(2)} = \sup_{j \geq n} \lambda_j$
- (3)  $Z_n^{(1)} = \inf_{j \geq n} Z_j$
- (4)  $Z_n^{(2)} = \sup_{j \geq n} Z_j$

**First, we prove that for  $i = 1$** , there exists a subsequence  $(\lambda_{n_k}^{(i)}, Z_{n_k}^{(i)})$  such that  $u_{\lambda_{n_k}^{(i)}, Z_{n_k}^{(i)}} \rightarrow u_{\lambda, Z}$ . **Note that this property also holds true for  $i = 2$ , but we don't prove it as the case is identical to  $i = 1$ .** The sequence  $(\lambda_n^{(1)}, Z_n^{(1)})$  also converges to  $(\lambda, Z)$ . So, there exists a subsequence  $n_k$  such that  $u_{\lambda_{n_k}^{(1)}, Z_{n_k}^{(1)}} \rightarrow u_*$  in  $C^2(\Omega) \cap C^1(\bar{\Omega})$  because  $u_{\lambda_{n_k}^{(1)}, Z_{n_k}^{(1)}} \in C^{2, \tau}(\bar{\Omega})$  by Proposition 3.12, which is compactly embedded in  $C^2(\bar{\Omega}) \subset C^2(\Omega) \cap C^1(\bar{\Omega})$ . Therefore the following pointwise convergence holds

$$\begin{aligned} 0 &= \lim_{k \rightarrow \infty} \left[ -\frac{\sigma^2}{2} \nabla^2 u_{\lambda_{n_k}, Z_{n_k}} + \frac{1}{2} |\nabla u_{\lambda_{n_k}, Z_{n_k}}|^2 - h\left(x, \frac{1}{Z_{n_k}} e^{-\frac{2}{\sigma^2} u_{\lambda_{n_k}, Z_{n_k}}}\right) + \lambda_{n_k} \right] \\ &= -\frac{\sigma^2}{2} \nabla^2 u_* + \frac{|\nabla u_*|^2}{2} - h\left(x, \frac{1}{Z} e^{-\frac{2}{\sigma^2} u_*}\right) + \lambda \\ 0 &= \lim_{k \rightarrow \infty} \nabla u_{\lambda_{n_k}, Z_{n_k}} \cdot \nu|_{x \in \partial\Omega} = \nabla u_* \cdot \nu|_{x \in \partial\Omega}. \end{aligned}$$

So,  $u_* = u_{\lambda, Z}$ , by uniqueness proved in Proposition 3.12. Now, by design,  $\lambda_{n_k}^{(2)} \geq \lambda_n \geq \lambda_{n_k}^{(1)}$  for all  $n \geq n_k$  and similarly for  $Z_n$ , hence  $u_{\lambda_{n_k}^{(1)}, Z_{n_k}^{(1)}} \geq u_{\lambda_n, Z_n} \geq u_{\lambda_{n_k}^{(2)}, Z_{n_k}^{(2)}}$  by Lemma 3.14. So,  $u_{\lambda_n, Z_n} \rightarrow u_{\lambda, Z}$  in  $L^\infty(\Omega)$ .  $\square$

**Proposition 3.16.** For each  $Z \in (0, \infty)$ , define  $I_1(\cdot; Z) : (\Lambda_1, \Lambda_2) \rightarrow \mathbb{R}$  by

$$I_1(\lambda; Z) = \int_{\Omega} u_{\lambda, Z} \, dx = \int_{\Omega} \Phi(\lambda, Z) \, dx.$$

Assume  $h$  is strictly increasing with respect to  $m$ . Then, for every  $Z \in (0, \infty)$ , there exists a unique  $\lambda = \lambda(Z)$  such that  $I_1(\lambda(Z); Z) = 0$ . Furthermore

$$(17) \quad \inf_{x \in \Omega} h\left(x, \frac{1}{Z}\right) \leq \lambda(Z) \leq \sup_{x \in \Omega} h\left(x, \frac{1}{Z}\right).$$

*Proof.* We use the intermediate value theorem to prove this proposition. There are three parts to prove

- (1) For every  $Z \in (0, \infty)$ , there exists  $\lambda_1 \leq \lambda_2 \in (\Lambda_1, \Lambda_2)$  such that  $I_1(\lambda_1; Z) \leq 0$  and  $I_1(\lambda_2; Z) \geq 0$ .
- (2)  $I_1(\lambda; Z)$  is continuous with respect to  $\lambda$  in  $[\lambda_1, \lambda_2]$ .
- (3)  $I_1(\lambda; Z)$  is strictly decreasing with respect to  $\lambda$ .

Using part (1), part (2) and the intermediate value theorem, for every  $Z \in (0, \infty)$  there exists some  $\lambda$  such that  $I_1(\lambda; Z) = 0$ . Using part (3), this  $\lambda$  is unique, so the function  $Z \mapsto \lambda(Z)$  is well defined.

Part (1): Take  $\lambda_1 = \sup_{x \in \Omega} h\left(x, \frac{1}{Z}\right) > \Lambda_1$ . Recall that  $\bar{u}_{\lambda_1, Z} = \max\left(-\frac{\sigma^2}{2} \log(ZM_{\lambda_1}^1), 0\right)$ , where  $M_{\lambda_1}^1$  satisfies  $h\left(x, M_{\lambda_1}^1\right) \leq \lambda_1$ , but we can take  $M_{\lambda_1}^1 = \frac{1}{Z}$  by our choice of  $\lambda_1$ . So,  $u_{\lambda_1, Z} \leq \bar{u}_{\lambda_1, Z} = 0$ , and therefore  $I_1(\lambda_1; Z) \leq 0$ . The choice for  $\lambda_2$  is  $\lambda_2 = \inf_{x \in \Omega} h\left(x, \frac{1}{Z}\right)$  and the proof is similar to the above.

Part (2): Take  $\lambda_1, \lambda_2$  as above. By Propositions 3.12 and 3.15 and Lemma 3.14,  $u_{\lambda, Z}$  is continuous with respect to  $\lambda$  in  $L^\infty(\Omega)$ , and  $u_{\lambda_2, Z} \leq u_{\lambda, Z} \leq \bar{u}_{\lambda_1, Z}$  for any  $\lambda \in [\lambda_1, \lambda_2]$ . So, by the dominated convergence theorem,  $I_1$  is continuous in  $\lambda$ .

Part (3): Take  $\lambda_1 < \lambda_2$ , so  $u_{\lambda_1, Z} \geq u_{\lambda_2, Z}$ , from Lemma 3.14. Since solutions to the PDE (15a)–(15c) are unique, there exists  $a \in \Omega$  such that  $u_{\lambda_1, Z}(a) \neq u_{\lambda_2, Z}(a)$ . Hence,  $u_{\lambda_1, Z}(a) > u_{\lambda_2, Z}(a)$  and so by continuity  $I_1(\lambda_1, Z) > I_1(\lambda_2, Z)$ .  $\square$

*Remark 3.17.* This proposition ensures that for any  $Z \in (0, \infty)$  there exists  $\lambda = \lambda(Z)$  and  $u = u_{\lambda(Z), Z}$  satisfying (15a)–(15c), (15e), so we are left to find  $Z^*$  such that (15d) holds.

**Lemma 3.18.** Assume  $h$  is strictly increasing with respect to  $m$ . Then, the function  $\lambda(Z)$  is strictly decreasing.

*Proof.* From Lemma 3.14,  $u_{\lambda, Z}$  is strictly decreasing with respect to  $Z$ . Now, suppose  $Z_1 < Z_2$  then

$$0 = I_1(\lambda(Z_2), Z_2) = I_1(\lambda(Z_1), Z_1) > I_1(\lambda(Z_1), Z_2).$$

Therefore, since  $I_1$  is strictly decreasing in  $\lambda$ ,  $\lambda(Z_2) < \lambda(Z_1)$ . So,  $\lambda(Z)$  is strictly decreasing with respect to  $Z$ .  $\square$

**Lemma 3.19.** Assume  $h$  is strictly increasing with respect to  $m$ . Then, the function  $\lambda(Z)$  is continuous.

*Proof.* We will prove  $\lambda(Z)$  is sequentially continuous. Let  $Z_n$  be a sequence in  $(0, \infty)$  that converges to  $Z \in (0, \infty)$ . Consider two sequences:  $Z_n^{(i)}$  for  $i = 1, 2$ , which will be used to sandwich our original sequence. The sequences are defined term-wise by

- (1)  $Z_n^{(1)} = \inf_{j \geq n} Z_j$
- (2)  $Z_n^{(2)} = \sup_{j \geq n} Z_j$

Note that  $Z_n^{(1)}, Z_n^{(2)} \rightarrow Z$  and are increasing and decreasing sequences respectively. Furthermore, there exists  $\bar{Z}, \bar{Z}$  such that  $Z_n^{(1)}, Z_n^{(2)} \in [\bar{Z}, \bar{Z}]$  for every  $n \in \mathbb{N}$ . So, since  $\lambda(Z)$  is decreasing,  $\lambda(Z_n^{(1)}), \lambda(Z_n^{(2)}) \in [\lambda(\bar{Z}), \lambda(\bar{Z})]$  and are decreasing and increasing respectively. Therefore,  $\lambda(Z_n^{(1)}) \rightarrow \lambda^{(1)}$  and  $\lambda(Z_n^{(2)}) \rightarrow \lambda^{(2)}$ . Using continuity of  $I_1$ , the following equality holds for  $i = 1, 2$ :

$$0 = \lim_{n \rightarrow \infty} I_1\left(\lambda(Z_n^{(i)}), Z_n^{(i)}\right) = I_1\left(\lambda^{(i)}, Z\right).$$

Hence, by definition,  $\lambda^{(1)} = \lambda^{(2)} = \lambda(Z)$ . Since  $Z_n^{(i)}$  bound  $Z_n$ , then  $\lambda(Z_n^{(i)})$  bound  $\lambda(Z_n)$  and  $\lambda(Z_n) \rightarrow \lambda(Z)$ .  $\square$

**Proposition 3.20.** Define  $I_2 : (0, \infty) \rightarrow (0, \infty)$  by

$$(18) \quad I_2(Z) = \int_{\Omega} \frac{1}{Z} e^{-\frac{2}{\sigma^2} u_{\lambda(Z), Z}} dx.$$

Assume  $h$  is strictly increasing with respect to  $m$ . Then, there exists a unique  $Z^* \in (0, \infty)$  such that  $I_2(Z^*) = 1$ .

*Proof.* Similar to the proof of Proposition 3.16, we prove this proposition using the intermediate value theorem. Again, there are three parts to prove

- (1) There exists  $Z_1 \leq Z_2 \in (0, \infty)$  such that  $I_2(Z_1) \geq 1$  and  $I_2(Z_2) \leq 1$ .
- (2)  $I_2(Z)$  is continuous with respect to  $Z$  for all  $Z \in [Z_1, Z_2]$ .
- (3)  $I_2(Z)$  is strictly decreasing with respect to  $Z$ .

Steps (1) and (2) prove existence via the intermediate value theorem, step (3) proves uniqueness.

Step (1): From assumption (MFG<sub>2</sub>), **there exists**  $Z_2 \geq |\Omega|$  such that

$$(19) \quad \sup_{x \in \Omega} h\left(x, \frac{1}{Z_2}\right) \leq \inf_{x \in \Omega} h\left(x, \frac{1}{|\Omega|}\right).$$

Since  $\lambda(Z_2) \leq \sup_{x \in \Omega} h\left(x, \frac{1}{Z_2}\right)$ , from (17), it follows that

$$u_{\lambda(Z_2), Z_2} \geq u_{\sup_{x \in \Omega} h\left(x, \frac{1}{Z_2}\right), Z_2} \geq u_{\sup_{x \in \Omega} h\left(x, \frac{1}{|\Omega|}\right), Z_2} = \min\left(-\frac{\sigma^2}{2} \log Z_2 M, 0\right),$$

where  $M$  satisfies  $h(x, M) \geq \sup_{x \in \Omega} h\left(x, \frac{1}{Z_2}\right)$  for all  $x$ , from the proof of Lemma 3.11. But, from (19), this is satisfied by  $M = \frac{1}{|\Omega|}$ . In this case,  $\min\left(-\frac{\sigma^2}{2} \log Z_2 M, 0\right) = -\frac{\sigma^2}{2} \log \frac{Z_2}{|\Omega|}$ . Thus

$$I_2(Z_2) \leq \int_{\Omega} \frac{1}{Z_2} e^{-\frac{2}{\sigma^2} \left(-\frac{\sigma^2}{2} \log \frac{Z_2}{|\Omega|}\right)} dx = \int_{\Omega} \frac{1}{|\Omega|} dx = 1.$$

A similar procedure works to find  $Z_1$ , in which case  $Z_1$  satisfies  $Z_1 \leq |\Omega|$  and  $\inf_{x \in \Omega} h\left(x, \frac{1}{Z_1}\right) \geq \sup_{x \in \Omega} h\left(x, \frac{1}{|\Omega|}\right)$ .

Step (2): Take  $Z_1 \leq Z_2$  as in Step (1). Then, by (17), for every  $Z \in [Z_1, Z_2]$  there exists  $C_1, C_2 \in \mathbb{R}$  such that

$$C_2 = \inf_{x \in \Omega} h\left(x, \frac{1}{Z_2}\right) \leq \lambda(Z_2) \leq \lambda(Z) \leq \lambda(Z_1) \leq \sup_{x \in \Omega} h\left(x, \frac{1}{Z_1}\right) = C_1.$$

So  $u_{C_1, Z_2} \leq u_{\lambda(Z), Z} \leq u_{C_2, Z_1}$  for every  $Z \in [Z_1, Z_2]$ . So, **using** the dominated convergence theorem, continuity of  $u_{\lambda, Z}$  with respect to  $(\lambda, Z)$  and continuity of  $\lambda(Z)$  with respect to  $Z$ ,  $I_2(Z)$  is continuous.

Step (3): Take  $Z < \bar{Z}$ . Then, there exists  $a \in \Omega$  such that

$$u_{\lambda(Z), \bar{Z}}(a) < u_{\lambda(\bar{Z}), \bar{Z}}(a).$$

Therefore, at  $a \in \Omega$ :

$$\frac{1}{Z} e^{-\frac{2}{\sigma^2} u_{\lambda(Z), Z}} = \frac{1}{Z} e^{-\frac{2}{\sigma^2} u_{\lambda(Z), Z}} > \frac{1}{\bar{Z}} e^{-\frac{2}{\sigma^2} u_{\lambda(\bar{Z}), \bar{Z}}}.$$

So,  $I_2(Z) > I_2(\bar{Z})$ , because of the continuity of  $u_{\lambda, Z}$ . This proves  $I_2$  is strictly decreasing.  $\square$

*End of proof of Theorem 3.10.* First, assume  $h$  is a strictly increasing function in  $m$ . Then, **there exists** a unique  $Z^* \in (0, \infty)$  such that  $I_2(Z^*) = 1$ . **Moreover**, the triple  $(u_{\lambda(Z^*), Z^*}, \lambda(Z^*), Z^*)$  is a solution to the system (15). Furthermore, suppose  $(u', \lambda', Z')$  is also a solution of (15). This implies  $u'$  satisfies (15a)–(15c), so  $u' = u_{\lambda', Z'}$ , from uniqueness proven in Proposition 3.12. Then,  $u_{\lambda', Z'}$  also solves (15e). So, by uniqueness proven in Proposition 3.16,  $\lambda' = \lambda(Z')$ . Finally,  $u' = u_{\lambda(Z'), Z'}$  meets the integral constraint (15d). So, from uniqueness proven in Proposition 3.20,  $Z' = Z^*$ . Therefore,  $(u', \lambda', Z') = (u_{\lambda(Z^*), Z^*}, \lambda(Z^*), Z^*)$  and the unique solution to the MFG problem is given by  $(m_{Z^*}, u_{\lambda(Z^*), Z^*}, \lambda(Z^*))$ , where  $m_{Z^*}$  is defined by  $m_{Z^*} = \frac{1}{Z^*} e^{-\frac{2}{\sigma^2} u_{Z^*}}$ .

Now assume  $h$  is an increasing function in  $m$ , and define  $h_{\epsilon}(x, m)$  by

$$h_{\epsilon}(x, m) = h(x, m) + \epsilon \log(|\Omega|m).$$

Then, for every  $\epsilon \in (0, 1]$ ,  $h_\epsilon$  is a strictly increasing function of  $m$ . Furthermore,  $h_\epsilon$  still satisfies assumptions (MFG<sub>1</sub>)–(MFG<sub>3</sub>). Therefore, there exists a unique solution  $(u_\epsilon, \lambda_\epsilon, Z_\epsilon)$  to the MFG system (15). From Proposition 3.20,  $Z_\epsilon \in [Z_\epsilon^1, Z_\epsilon^2]$  for some  $Z_\epsilon^1, Z_\epsilon^2 \in (0, \infty)$  such that

$$\begin{aligned} 0 < Z_\epsilon^1 \leq |\Omega| \leq Z_\epsilon^2 < \infty \\ \sup_{x \in \Omega} h_\epsilon \left( x, \frac{1}{Z_\epsilon^2} \right) &\leq \inf_{x \in \Omega} h_\epsilon \left( x, \frac{1}{|\Omega|} \right) \\ \inf_{x \in \Omega} h_\epsilon \left( x, \frac{1}{Z_\epsilon^1} \right) &\geq \sup_{x \in \Omega} h_\epsilon \left( x, \frac{1}{|\Omega|} \right). \end{aligned}$$

By definition,  $h_\epsilon \left( x, \frac{1}{Z_\epsilon^2} \right) \leq h \left( x, \frac{1}{Z_\epsilon^2} \right)$  and  $h_\epsilon \left( x, \frac{1}{|\Omega|} \right) = h \left( x, \frac{1}{|\Omega|} \right)$ , and a similar inequality holds for  $Z_\epsilon^1$ . So, **there exist**  $Z^1 \in (0, |\Omega|]$  and  $Z^2 \in [|\Omega|, \infty)$ , **which are** independent of  $\epsilon$ , such that  $Z_\epsilon \in [Z^1, Z^2]$  for every  $\epsilon \in (0, 1]$ . Now, from Lemma 3.18

$$\lambda_\epsilon = \lambda(Z_\epsilon) \in [\lambda(Z^2), \lambda(Z^1)].$$

So, take a sequence  $\epsilon_n$  such that  $\lim_{n \rightarrow \infty} \epsilon_n = 0$ . Then, since  $u_\epsilon \in C^{2,\tau}(\bar{\Omega})$ , which is compactly embedded in  $C^2(\Omega) \cap C^1(\bar{\Omega})$ , there exists a subsequence, also denoted by  $n$ , such that  $u_{\epsilon_n} \rightarrow u_0$  with convergence in  $C^2(\Omega) \cap C^1(\bar{\Omega})$ ,  $Z_{\epsilon_n} \rightarrow Z_0 \in [Z^1, Z^2]$ , and  $\lambda_{\epsilon_n} \rightarrow \lambda_0 \in [\lambda(Z^2), \lambda(Z^1)]$ . So, by taking limits

$$-\frac{\sigma^2}{2} \nabla^2 u_0 + \frac{|\nabla u_0|^2}{2} - h \left( x, \frac{1}{Z_0} e^{-\frac{\sigma^2}{2} u_0} \right) + \lambda_0 = 0.$$

Similarly,  $(u_0, \lambda_0, Z_0)$  satisfies (15). So, **solutions to (15) exist if  $h$  is increasing in  $m$ .**

For uniqueness, assume  $(u_1, \lambda_1, Z_1)$  and  $(u_2, \lambda_2, Z_2)$  are both solutions of (15) and  $\lambda_1 \leq \lambda_2$ . Define  $u = u_2 - \frac{\sigma^2}{2} \log \left( \frac{Z_1}{Z_2} \right)$ , then  $u$  satisfies

$$-\frac{\sigma^2}{2} \nabla^2 u + \frac{|\nabla u|^2}{2} - h \left( x, \frac{1}{Z_1} e^{-\frac{\sigma^2}{2} u} \right) + \lambda_1 = 0.$$

Define  $v = u - u_2$  and suppose there exists  $x \in \bar{\Omega}$  such that  $v(x) > 0$ . Then,  $v$  has a maximum at  $x^*$  and  $v(x^*) > 0$ . By Hopf's lemma  $x^* \in \Omega$  and by the maximum principle  $v$  is constant in the set  $\Omega_+ = \{x \in \Omega : v(x) \geq 0\}$  (see the proof of Proposition 3.12 for the details of such an argument). By assumption,  $x^* \in \Omega_+$  and  $v(x^*) > 0$ , so  $\Omega_+ = \Omega$  by continuity of  $v$ . Hence,  $v$  is constant in  $\Omega$  and  $v > 0$ . However, from the integral constraint (15d)

$$(20) \quad 0 = \int_{\Omega} \frac{1}{Z_1} e^{-\frac{\sigma^2}{2} u_1} \left( 1 - e^{-\frac{\sigma^2}{2} v} \right) dx = |\Omega| \left( 1 - e^{-\frac{\sigma^2}{2} v} \right),$$

since  $1 - e^{-\frac{\sigma^2}{2} v}$  is constant. So  $v = 0$ , contradicting the assumption  $v(x^*) > 0$ . Therefore,  $v(x) \leq 0$  for all  $x \in \Omega$ . This implies that  $1 - e^{-\frac{\sigma^2}{2} v} \leq 0$ , and subsequently that

$$0 = \int_{\Omega} \frac{1}{Z_1} e^{-\frac{\sigma^2}{2} u_1} \left( 1 - e^{-\frac{\sigma^2}{2} v} \right) dx \leq 0,$$

with equality if and only if  $v = 0$ . Therefore  $u_2 = u$ , which implies  $Z_1 = Z_2$ , using the integral constraint (15e), and subsequently that  $u_1 = u_2$ . Finally, by subtracting the PDE (15b) satisfied by  $u_1$  from the one satisfied by  $u_2$ ,  $\lambda_2 = \lambda_1$ . Therefore solutions are unique.  $\square$

#### 4. QUADRATIC POTENTIAL

In this section we consider a specific example, **in  $\mathbb{R}$ , of a** quadratic potential and a logarithmic congestion term,  $h(x, m) = \beta x^2 + \log m$  for some constant  $\beta \geq 0$ . This problem has been studied extensively in [29] and [31] and admits explicit solutions, **which mean the MFG and BRS solutions can be compared analytically**. Note that **no** boundary conditions or integral constraints on  $u$  **are imposed, since the domain is  $\mathbb{R}$** , rather than a bounded domain. Therefore, it doesn't fit directly into the framework for existence and uniqueness proven in the previous section. It is, however, one of the few illustrative examples where explicit solutions are known. This allows us to make an analytical comparison of the two models and use the solution to validate the proposed numerical methods.

**4.1. The MFG.** The stationary MFG model studied in [29] and [31], with the integral constraints used in this paper, is given by:

$$(21a) \quad \frac{\sigma^2}{2} \partial_{xx}^2 m + \partial_x (m \partial_x u) = 0, \quad x \in \mathbb{R},$$

$$(21b) \quad -\frac{|\partial_x u|^2}{2} + \log m + \beta x^2 + \frac{\sigma^2}{2} \partial_{xx}^2 u + \lambda = 0, \quad x \in \mathbb{R},$$

$$(21c) \quad \int_{\mathbb{R}} m \, dx = 1,$$

where  $\lambda \in (-\infty, \infty)$  is a constant to be found as part of the solution, and  $\sigma, \beta \geq 0$  are given parameters.

**Proposition 4.1.** *A solution to the stationary MFG system (21) exists and has an explicit form*

$$(22a) \quad m(x) = \left(\frac{a}{\pi}\right)^{1/2} e^{-ax^2}$$

$$(22b) \quad u(x) = bx^2$$

$$(22c) \quad \lambda = \log\left(\frac{\pi}{a}\right) - \sigma^2 b,$$

where the constants  $a, b \geq 0$  are given by

$$a = \beta, \quad b = 0,$$

if  $\sigma = 0$ , or

$$a = \frac{-1 + (1 + 2\sigma^4\beta)^{1/2}}{\sigma^4}, \quad b = \frac{-1 + (1 + 2\sigma^4\beta)^{1/2}}{2\sigma^2},$$

if  $\sigma > 0$ .

The proof is straight-forward using substitution.

**4.2. The BRS.** Next, we consider the respective stationary BRS model. It is given by

$$(23a) \quad \partial_x (m \partial_x (\log m + \beta x^2)) + \frac{\sigma^2}{2} \partial_{xx}^2 m = 0, \quad x \in \mathbb{R}$$

$$(23b) \quad \int_{\mathbb{R}} m \, dx = 1.$$

**Proposition 4.2.** *The solution to the stationary BRS equation (23) is*

$$m(x) = \left(\frac{2\beta}{(2 + \sigma^2)\pi}\right)^{1/2} e^{-\frac{2\beta}{(2 + \sigma^2)}x^2}.$$

Again, the claim follows from substitution.

**4.3. Comparison.**

**Proposition 4.3.** *For  $\sigma > 0$ , the stationary distributions of the MFG system (21) and the BRS (23) are given by normal distributions, with mean 0 and variances  $a_1$  and  $a_2$  respectively, where*

$$a_1 = \frac{\sigma^4}{-2 + 2(1 + 2\sigma^4\beta)^{1/2}}$$

$$a_2 = \frac{2 + \sigma^2}{4\beta}.$$

Then, for fixed  $\beta \geq 0$

$$(24a) \quad \lim_{\sigma^2 \rightarrow 0} \frac{a_2}{a_1} = 1$$

$$(24b) \quad \lim_{\sigma^2 \rightarrow \infty} \frac{a_2}{a_1} = \frac{1}{(2\beta)^{1/2}}.$$

While, for fixed  $\sigma > 0$

$$(25a) \quad \lim_{\beta \rightarrow 0} \frac{a_2}{a_1} = 1 + \frac{\sigma^2}{2}$$

$$(25b) \quad \lim_{\beta \rightarrow \infty} (2\beta)^{1/2} \frac{a_2}{a_1} = \frac{2 + \sigma^2}{\sigma^2}.$$







(A)  $\beta = 0.1, \frac{\sigma^2}{2} = 10$ (B)  $\beta = 0.1, \frac{\sigma^2}{2} = 0.2$ (C)  $\beta = 1, \frac{\sigma^2}{2} = 10$ (D)  $\beta = 1, \frac{\sigma^2}{2} = 0.2$ (E)  $\beta = 10, \frac{\sigma^2}{2} = 10$ (F)  $\beta = 10, \frac{\sigma^2}{2} = 0.2$ 

FIGURE 1. Simulations of BRS and MFG with quadratic potential and logarithmic congestion

*Proof.* The first part of this proof is trivial from the previous propositions. Now

$$\frac{a_2}{a_1} = \frac{(2 + \sigma^2) ((1 + 2\sigma^4\beta)^{1/2} - 1)}{2\sigma^4\beta}.$$

Using a Taylor expansion of  $(1 + x)^{1/2}$  around  $x = 0$  gives behaviour for small  $\sigma^2$  i.e.

$$\frac{a_2}{a_1} = \frac{(2 + \sigma^2)(\sigma^4\beta + o(\sigma^4))}{2\sigma^4\beta}.$$

Hence,  $\lim_{\sigma^2 \rightarrow 0} \frac{a_2}{a_1} = \lim_{\sigma^2 \rightarrow 0} \frac{2+\sigma^2}{2} = 1$ . The other limit can be calculated as

$$\lim_{\sigma^2 \rightarrow \infty} \frac{a_2}{a_1} = \lim_{\sigma^2 \rightarrow \infty} \frac{(2 + \sigma^2)(1 + 2\sigma^4\beta)^{1/2}}{2\sigma^4\beta} = \lim_{\sigma^2 \rightarrow \infty} \frac{(2 + \sigma^2)(2\beta)^{1/2}\sigma^2}{2\sigma^4\beta} = \frac{1}{(2\beta)^{1/2}}.$$

The limits as  $\beta \rightarrow 0, \infty$ , for fixed  $\sigma$ , follows from straight forward calculations.  $\square$

This result is an important **initial insight into** how the behaviour of the BRS and MFG may vary, as well as the importance certain parameters play in the difference **between the models**. The limit in (24a) shows that the existence of noise is **required for the two models to behave differently**. However, as soon as there is noise, its effect on the relative difference plays a less important role than the strength of the quadratic potential, this can be seen in (24b) and (25b). Specifically, the limit (25b) shows that the relative difference between the variances of the two distributions grows like  $\beta^{\frac{1}{2}}$ , which means the BRS distribution reacts more rapidly with changes to the potential strength than the MFG. This suggests that the MFG is more affected by congestion or is a more congestion-averse model than the BRS one.

At a conceptual level this agrees with the formulation of the MFG and BRS systems. The agents in the BRS are acting myopically, only reacting to the situation as it currently exists, which isn't the case in the MFG. As a result, the BRS agents don't **anticipate any** future congestion **arising** from their behaviour, and hence they move towards the minimum of  $\beta x^2$  more rapidly than the MFG agents, **which** do **anticipate** the future cost of the congestion that results from their behaviour. Therefore, thinking of the stationary solutions as the long time, time-averaged, behaviour of the models, the stationary BRS will result in a distribution that appears to take into account the congestion less than the MFG, and hence one with a smaller variance. This is confirmed by the result (25b) and it quantifies the extent to which the BRS ignores the congestion compared with the MFG.

For this model we have run a variety of simulations — both to confirm our numerical methods (see Section 5 for methods) and to visualise how the parameters affect the distributions. Figure 1 shows the results of these simulations on a bounded domain for a variety of parameter choices. Although the formulation on a bounded domain is slightly different than the one in this section, the same behaviour can be seen. For small  $\sigma$ , the difference between the models doesn't change much as  $\beta$  increases, while, for large values of  $\sigma$ , the BRS model is affected by changes to  $\beta$  **to a greater extent than the MFG model**. In both cases, the BRS and MFG are more closely aligned when  $\beta$  is small.

## 5. SIMULATIONS

Due to the complexity of the MFG, and the integral constraints that have to be satisfied for both the MFG and BRS models, we are unable to provide a systematic analysis of the difference between the distribution of agents in the two models. However, a good intuition can be gained through numerical simulations. Therefore, we conclude with presenting various computational experiments, which illustrate the difference between solutions to the BRS and MFG for different choices running costs and potentials.

**5.1. Solving the stationary BRS and MFG.** Solutions to the stationary BRS (8) can be computed by finding the zeros of the function  $G_{Z,x}(m)$ , at every discrete grid point  $x$  on a grid given  $Z$ . To compute the roots of  $G_{Z,x}$  at each grid point  $x$ , we use a Newton–Raphson method. Then, having found  $m_Z(x)$  for a particular value of  $Z$ , we can differentiate the implicit formula  $m_Z = \frac{1}{z} e^{-\frac{2}{\sigma^2} h(x, m_Z)}$  with respect to  $Z$ , and use a Newton–Raphson method to find  $Z$  such that  $\Phi(Z) = \int_{\Omega} m_Z dx = 1$ . In practice this means iterating between the two Newton–Raphson methods: first finding  $m_{Z_n}$ , then computing  $Z_{n+1}$ , then recomputing  $m_{Z_{n+1}}$  and repeating until convergence.

The forward-backward structure of MFG presents challenges not only for the analysis but also for the numerical simulations. Most solvers are based on a consistent finite difference discretization of the Fokker-Planck and its adjoint Hamilton-Jacobi Bellman equation, using a fixed point iteration. The numerical analysis of these schemes for the finite and infinite time horizon problem were analysed in a series of papers, see [2]. Certain types of MFG can be reformulated as parabolic optimal control problems. Then computational approaches, such as alternating directions methods, can be used. We refer to the excellent overview by Achdou, Camilli and Capuzzo-Dolcetta [1] for more detailed information.

The solution to the stationary MFG (13) is found using an iterative procedure. Given an admissible initial iterate  $m^l$ ,  $l = 0$ , we solve the HJB equation (13d) to obtain  $u^l$  and  $\lambda^l$ . Note that we include the constraint (13h) via a Lagrange multiplier. In the final step of the iteration, we update the distribution of agents by solving the FPE (13c) using  $u^{l+1}$  to obtain  $m^{l+1}$ . This procedure is repeated until convergence. Note that we performed a damped update

$$u^l = \omega u^{l-1} + (1 - \omega) v^l \text{ and } m^l = m^{l-1} + (1 - \omega) q^l$$

where  $v^l, q^l$  are the undamped solutions of each iteration process. Solutions to the HJB and the FPE are obtained by using an  $H^1$  conforming finite element discretisation. While conducting the computational experiments, we found that for cost functions with larger potentials or with more sensitive dependence on the density, greater damping was required to ensure the iterative procedure converged. In the simulations, using a trial and error approach, a damping ratio of  $\omega = 0.9$  was decided upon because it ensured convergence for all of the potentials used, without detrimentally impacting the speed of the simulations.

When considering the similarity of the two models, there are two possible comparisons to make. The first is to compare the distribution of agents. The second is to compare the function  $u$ , which we compute in the MFG, to a corresponding function that we can compute for the BRS model. In Section 2.1 it can be seen that  $u(x, 0)$  is the expected cost, over the entire time horizon, that an agent, starting in position  $x$ , incurs in the MFG model. For the dynamic BRS model, one would have to explicitly compute the expected cost.

When it comes to the stationary model, as in the setup of the paper, the expected cost over the whole time horizon blows up as the time horizon goes to  $\infty$ , because there is no discount factor in the cost functional (2). Instead, for the MFG in the stationary case, the constant  $\lambda$  is the limit of the time averaged expected cost of an agent starting at any position, as the time horizon goes to  $\infty$ . Furthermore, the function  $u$ , which we compute as part of the solution to the stationary MFG, is the difference between the expected long-term cost of an agent that started in position  $x$ , and the space-averaged expected long-term cost, which is given approximately by  $\lambda T$  for large time horizons  $T$ .

On the other hand, for the stationary BRS, the information about the time-averaged long-term cost, and the fluctuations in the long-term cost, are not readily available from the computations that are done to find the stationary BRS distribution. So, although it would be beneficial to compare the long-term costs related to the two models, this information is unfortunately unavailable to us in the case of the BRS. Therefore, our numerical simulations and comparisons are focused on comparing the distribution of agents.

**5.2. Single well potential.** In the first examples, we investigate the behavior of solutions to both models for cost functionals of the form  $h(x, m) = F(m) + \beta x^2$ , using a variety of functions  $F$  and parameters  $\beta$ . We also analyse how the noise level  $\sigma$  affects the similarity of the models. Note that the case  $F(m) = \log(m)$  was already discussed in Section 4. We are particularly interested in how penalising congestion affects the solutions. We have done this by considering functions  $F(m)$  of the form  $F(m) = m^\alpha$  for some  $\alpha > 0$  or  $F(m) = \frac{1}{m_{\max} - m}$  for some  $m_{\max} > \frac{1}{\Omega}$  affect solutions. The last choice introduces a ‘barrier’ above which the density can not exceed. Using such a congestion term is more realistic from a modelling perspective than either the logarithmic or power-law term as it forces densities to stay below a certain physical reasonable limit. The simulations showed a similar dependence on the parameters  $\beta$  and




FIGURE 2. Simulations of BRS and MFG with  $h(x, m) = m^{10} + 10x^2$




FIGURE 3. Simulations of BRS and MFG with  $h(x, m) = \frac{1}{m_{\max} - m} + x^2$

$\sigma$  compared with the logarithmic congestion term, in fact the same can be said for all of our simulations. So, for all values of  $\sigma$ , the MFG model responded less to changes in the strength of the potential  $\beta$ , compared with the BRS model. However, the difference **becomes more** pronounced as  $\sigma$  increases and again it may be expected that as  $\sigma \rightarrow 0$  that the two models align very closely.

The most notable difference between the use of a logarithmic congestion term and a power-law congestion term is a difference in the shape of the distribution, particularly the flatness of the peak of the distribution, as shown in figure 2. Importantly, this characteristic is shared by both the MFG and the BRS, suggesting the congestion terms  $F(m)$  affect both models in similar ways. When looking at congestion terms of the form  $F(m) = \frac{1}{m_{\max} - m}$ , we can find certain regimes where the behaviour is similar to the logarithm, **and other regimes where the behaviour is closer to the power-law congestion**. When  $m_{\max}$  is large, as in figure 3a, the resulting distribution for both models looks like a normal distribution, similar to the case with logarithmic congestion. In fact when  $m_{\max}$  is very large, a formal asymptotic analysis, using a Taylor expansion around  $m_{\max}$  can be made, which shows

$$\frac{1}{m_{\max} - m} \approx \frac{1}{m_{\max}}.$$

Therefore the BRS satisfies the equation

$$m \approx \frac{1}{Z} e^{-\frac{2}{\sigma^2} \frac{\beta m_{\max} x^2 - 1}{m_{\max}}} \approx \frac{1}{Z} e^{-\frac{2\beta}{\sigma^2} x^2},$$

with a normalisation constant  $Z = \int_{\Omega} e^{-\frac{2\beta}{\sigma^2} x^2}$ , which corresponds, on  $\mathbb{R}$ , to a normal distribution with zero mean and variance  $\frac{\sigma^2}{4\beta}$ . The variance of the BRS solution found here differs from the logarithmic congestion case by  $\frac{2}{4\beta}$ . A similar analysis shows that when  $m_{\max} \rightarrow \infty$  the MFG approximately resembles a normal distribution with zero mean and variance  $\frac{\sigma^2}{2\beta}$ . In summary, solutions to the MFG and the BRS are both normal distributions with zero mean and with variances whose relative difference is  $\frac{1}{2}$ .

However, when  $m_{\max}$  is reduced, as in figure 3b, the peak flattens out in a similar but exaggerated way compared to the power-law congestion. It is interesting to note that the BRS seems to respond more to the change in  $m_{\max}$  than the MFG, this is in contrast to the role that  $\sigma$  plays in the two models, where the MFG responds more to changes in  $\sigma$ , compared with the BRS.

**5.3. Double well potential.** The previous subsection **provided** insight into how the form of the congestion term affects both models. In this section, we explore how the potential term affects each model. For this section, we will consider costs of the form  $h(x, m) = F_1(x) + \log(m)$ , where  $F_1(x)$  will be a double well potential (see figure 4). Since the key insight of this section is to understand how varying  $F_1$  affects the similarity of **the two models' solutions**, we have decided not to include results with congestion terms other than the logarithm. From simulations, **we found** that the effect of changing the congestion term, from  $\log(m)$  to another term, is very similar, regardless of whether we are considering a single well or a double well.

Our simulations focus on five different double wells, which can be seen in figure 4. We vary the potentials as follows




FIGURE 4. Double well potentials for simulations

- (1) same depth, same width,
- (2) different depth, same width,
- (3) same depth, different width,
- (4) approximately similar perimeter,
- (5) approximately similar volume.

The simulations with the first two potentials, see figures 5 and 6, where the widths of the two wells are always the same, show that the two models display similar qualitative behaviour. As with the single well potential, as  $\sigma$  increases the discrepancy between the two models also increases, with the MFG model being more affected by the level of noise than the BRS. As expected, when the two wells are of equal depth, as in figure 5, then both the MFG and BRS attribute equal weight between the wells, while when one well is deeper than the other, as in figure 6, both models **distribute** more mass in the location of the deeper well. This is true for all values of  $\sigma$ , although the effect reduces as  $\sigma$  increases, as can be seen by comparing figures 5a and 6a to figures 5b and 6b respectively.

**So far**, we have seen that the qualitative behaviour of the two models **has tended to agree**. However, when we look at double well potentials where the **two wells have different widths**, we **observe some differences between the behaviour of the MFG and the BRS**. In figure 7, where the wells have the same depth but different width, the BRS still distributes density equally to the two wells. However, the MFG model results in a higher density in the wider well, than in the narrower well. The reason the width has no effect on the BRS can be seen from studying the implicit equation. The BRS solves  $m = \frac{1}{Z} e^{-\frac{2}{\sigma^2} h(x, m)}$ , where  $Z$  is a fixed constant. In the current case,  $h(x, m) = G(x) + F(m)$ . Since the potential  $G(x)$  **takes the same value** in each well, then the height of the BRS distribution will be the same in each well. The




FIGURE 5. Simulation of BRS and MFG with logarithmic congestion and potential given in figure 4a




FIGURE 6. Simulation of BRS and MFG with logarithmic congestion and potential given in figure 4b

MFG, however, is affected by the width of the well, because the MFG function  $u$  solves an elliptic equation. Hence, at each point  $x$ ,  $u(x)$  will depend on  $u(y)$  at points  $y \neq x$ , and not only on  $G(x)$ . In other words, the BRS depends only on the local value of the cost  $G$ , whereas the MFG depends also on the value of  $G$  across the whole domain. To understand why the MFG assigns greater density to the wider well, than the narrower well, we need to look at the underlying optimisation problem related to the MFG.

When considering the optimisation problems related to the dynamic MFG and its long-term behaviour, which results in its stationary state, we can see that the MFG model anticipates the future system. Therefore, agents in the wider well are willing to remain in it for longer, compared with agents in the narrower well. This is because the cost functional being optimised by the MFG agents includes a quadratic cost on their control, so the cost to move out of the wider well is higher than the cost to move out of the narrower well. Hence, in the stationary case the wider well has a higher density associated to it than the narrower well.

We have seen that increasing well width affects only the MFG, while increasing well depth affects both the MFG and BRS. Now, we can balance these effects to create situations where the two models qualitatively disagree. Figure 8 involves a double well where the width and depth of the wells differ but the area of each well is the same, while in figure 9 the perimeter of each well is the same. In the case of a small noise term, then both the MFG and BRS favour the deeper wells. However, with a larger noise




FIGURE 7. Simulation of BRS and MFG with logarithmic congestion and potential given in figure 4c




FIGURE 8. Simulation of BRS and MFG with logarithmic congestion and potential given in figure 4d

term, figure 8b shows that there are cases where the wider, shallower well is favoured by the MFG while the narrower, deeper well is favoured by the BRS.

## 6. CONCLUSION AND OUTLOOK

In this paper we have compared two models of interacting multi-agent systems in the stationary case. Through a proof of existence and uniqueness for each model, we have seen that the BRS model can be reformulated as an implicit equation. This shows that the BRS model only depends on local data of the cost function, while the MFG model, the solution of which is given by an elliptic equation, may have non-local dependencies on the data. The existence and uniqueness proofs were based on the important assumption that the congestion term is increasing. However, the regularity requirements on the MFG data are less strict than those on the BRS data. Finally the proof gave an insight into the dependence of each model on the diffusion coefficient. We want to remark that the strategy of the proof is interesting on its own and that the only similar results, presented in [18], are based on different assumptions. The analytic results are supported by numerical simulations and we investigated the similarities and differences of the MFG and BRS models, in various computational experiments. We plan to extend the analysis and simulations to the dynamic case in the future, and consider cost functions other than linear-quadratic ones.




FIGURE 9. Simulation of BRS and MFG with logarithmic congestion and potential given in figure 4e

#### REFERENCES

- [1] Y. Achdou, F. Camilli, and I. Capuzzo-Dolcetta. Mean field games: Convergence of a finite difference method. *SIAM Journal on Numerical Analysis*, 51(5):2585–2612, 2013.
- [2] Y. Achdou and I. Capuzzo-Dolcetta. Mean Field Games: Numerical Methods. *SIAM Journal on Numerical Analysis*, 48(3):1136–1162, jan 2010.
- [3] M. Barker. From mean field games to the best reply strategy in a stochastic framework. *Journal of Dynamics & Games*, 6(4):291–314, 2019.
- [4] J. D. Benamou, G. Carlier, and F. Santambrogio. Variational mean field games. *Modeling and Simulation in Science, Engineering and Technology*, 1:141–171, 2017.
- [5] C. Bertucci, J.-M. Lasry, and P.-L. Lions. Some remarks on mean field games. *Communications in Partial Differential Equations*, 44:205–227, 2019.
- [6] P. Biane and R. Speicher. Free diffusions, free entropy and free Fisher information. *Annales de l’institut Henri Poincaré (B) Probability and Statistics*, 37:581–606, 2001.
- [7] L. Boccardo, L. Orsina, and A. Porretta. Strongly coupled elliptic equations related to mean-field games systems. *Journal of Differential Equations*, 261:1796–1834, 2016.
- [8] P. E. Caines, M. Huang, and R. P. Malhamé. Mean Field Games. In T. Basar and G. Zaccour, editors, *Handbook of Dynamic Game Theory*, pages 1–28. Springer Berlin Heidelberg, 2017.
- [9] P. Cardaliaguet and P. Jameson Graber. Mean field games systems of first order. *ESAIM - Control, Optimisation and Calculus of Variations*, 21:690–722, 2015.
- [10] P. Cardaliaguet, J.-M. Lasry, P.-L. Lions, and A. Porretta. Long time average of mean field games. *Networks and Heterogeneous Media*, 7:279–301, 2012.
- [11] R. Carmona and F. Delarue. *Probabilistic Theory of Mean Field Games with Applications I*, volume 83 of *Probability Theory and Stochastic Modelling*. Springer International Publishing, Cham, 2018.
- [12] R. Carmona and F. Delarue. *Probabilistic Theory of Mean Field Games with Applications II*, volume 84 of *Probability Theory and Stochastic Modelling*. Springer International Publishing, Cham, 2018.
- [13] J. A. Carrillo, R. S. Gvalani, G. A. Pavliotis, and A. Schlichting. Long-Time Behaviour and Phase Transitions for the McKean-Vlasov Equation on the Torus. *Archive for Rational Mechanics and Analysis*, 2019.
- [14] J. A. Carrillo, R. J. McCann, and C. Villani. Kinetic equilibration rates for granular media and related equations: Entropy dissipation and mass transportation estimates. *Revista Matemática Iberoamericana*, 19:971–1018, 2003.
- [15] J. A. Carrillo, R. J. McCann, and C. Villani. Contractions in the 2-Wasserstein length space and thermalization of granular media. *Archive for Rational Mechanics and Analysis*, 179:217–263, 2006.
- [16] A. Cesaroni and M. Cirant. Concentration of ground states in stationary mean-field games systems. *Analysis and PDE*, 12:737–787, 2018.
- [17] L. Chayes and V. Panferov. The McKean-Vlasov equation in finite volume. *Journal of Statistical Physics*, 138:351–380, 2010.
- [18] M. Cirant. Multi-population Mean Field Games systems with Neumann boundary conditions. *Journal de Mathématiques Pures et Appliquées*, 103:1294–1315, 2015.
- [19] M. Cirant. Stationary focusing mean-field games. *Communications in Partial Differential Equations*, 41:1324–1346, 2016.
- [20] P. Degond, M. Herty, and J. G. Liu. Meanfield games and model predictive control. *Communications in Mathematical Sciences*, 15:1403–1422, 2017.
- [21] P. Degond, J. G. Liu, and C. Ringhofer. Large-scale dynamics of mean-field games driven by local nash equilibria. *Journal of Nonlinear Science*, 24:93–115, 2014.
- [22] D. Evangelista, R. Ferreira, D. A. Gomes, L. Nurbekyan, and V. Voskanyan. First-order, stationary mean-field games with congestion. *Nonlinear Analysis, Theory, Methods and Applications*, 173:37–74, 2018.
- [23] D. Evangelista and D. A. Gomes. On the Existence of Solutions for Stationary Mean-Field Games with Congestion. *Journal of Dynamics and Differential Equations*, 30:1365–1388, 2018.

- [24] L. C. Evans. *Partial Differential Equations*. American Mathematical Society, USA, 1998.
- [25] R. Ferreira and D. Gomes. Existence of Weak Solutions to Stationary Mean-Field Games through Variational Inequalities. *SIAM Journal on Mathematical Analysis*, 50:5969–6006, 2018.
- [26] R. Ferreira, D. Gomes, and T. Tada. Existence of weak solutions to first-order stationary mean-field games with Dirichlet conditions. *Proceedings of the American Mathematical Society*, 147:4713–4731, 2019.
- [27] D. Gomes and H. Sánchez Morgado. A stochastic Evans-Aronsson problem. *Transactions of the American Mathematical Society*, 366:903–929, 2013.
- [28] D. A. Gomes, S. Patrizi, and V. Voskanyan. On the existence of classical solutions for stationary extended mean field games. *Nonlinear Analysis, Theory, Methods and Applications*, 99:49–79, 2014.
- [29] D. A. Gomes, E. A. Pimentel, and V. Voskanyan. *Regularity Theory for Mean-Field Game Systems*. SpringerBriefs in Mathematics. Springer International Publishing, Cham, 2016.
- [30] D. A. Gomes and V. K. Voskanyan. Extended deterministic mean-field games. *SIAM Journal on Control and Optimization*, 54:1030–1055, 2016.
- [31] O. Guéant. A reference case for mean field games models. *Journal de Mathématiques Pures et Appliquées*, 92:276–294, 2009.
- [32] O. Guéant. *Mean field games equations with quadratic Hamiltonian: A specific approach*, volume 22. 2012.
- [33] M. Huang, P. E. Caines, and R. P. Malhamé. Distributed multi-agent decision-making with partial observations: asymptotic Nash equilibria. *Proc. the 17th Internat. Symposium on Math. Theory on Networks and Systems (MTNS2006), Kyoto, Japan*, pages 2725–2730, 2006.
- [34] M. Huang, P. E. Caines, and R. P. Malhamé. An invariance principle in large population stochastic dynamic games. *Journal of Systems Science and Complexity*, 20:162–172, 2007.
- [35] M. Huang, R. Malhamé, and P. Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle. *Communications in Information & Systems*, 6:221–251, 2006.
- [36] J.-M. Lasry and P.-L. Lions. Jeux à champ moyen. I – Le cas stationnaire. *Comptes Rendus Mathématique*, 343:619–625, 2006.
- [37] J.-M. Lasry and P.-L. Lions. Jeux à champ moyen. II – Horizon fini et contrôle optimal. *Comptes Rendus Mathématique*, 343:679–684, 2006.
- [38] J.-M. Lasry and P.-L. Lions. Mean field games. *Japanese Journal of Mathematics*, 2(1):229–260, mar 2007.
- [39] R. J. McCann. A convexity principle for interacting gases. *Advances in Mathematics*, 128:153–179, 1997.
- [40] K. Schmitt. Boundary value problems for quasilinear second order elliptic equations. *Nonlinear Analysis, Theory, Methods & Applications*, 2:263–309, 1978.
- [41] Y. Tamura. On asymptotic behaviours of the solution of a non-linear diffusion process. *J. Fac. Sci. Univ. Tokyo Sect. 1A, Math*, 31:195–221, 1984.
- [42] J. Tugaut. Phase transitions of McKean-Vlasov processes in double-wells landscape, 2014.