

Quantifying the Environmental and Food Biodiversity Impacts of Ultra-Processed Foods - Evidence from the EPIC Study

Jeroen Berden^{1,2}, Giles T. Hanley-Cook^{2,3}, Bernadette Chimera¹, Dagfinn Aune^{4,5,6}, Maria Gabriela M. Pinho⁷, Geneviève Nicolas¹, Bernard Srour⁸, Christopher J. Millett^{9,10}, Emine Koc Cakmak^{11,12}, Emmanuelle Kesse-Guyot⁸, Esther M. González-Gil¹, Eszter P. Vamos⁹, Jessica Blanco Lopez¹, Julia Baudry⁸, Justine Berlivet⁸, Kiara Chang⁹, Mathilde Touvier⁸, Charlotte Le Cornet¹³, Chloé Marques¹⁴, Christina C. Dahm¹⁵, Daniel B. Ibsen^{15,16}, Franziska Jannasch¹⁷, Guri Skeie¹⁸, Maria-José Sanchez^{19,20,21}, Matthias B. Schulze^{17,22}, Sara Grioni²³, Yvonne T. van der Schouw²⁴, Ana M. Jimenez Zabala^{25,26}, Anna Winkvist^{27,28}, Anne Tjønneland^{29,30}, Carlotta Sacerdote^{31,32}, Cecilie Kyrø²⁹, Elisabeth Weiderpass³³, Marcela Guevara^{19,34,35}, Pauline Frenoy¹⁴, Rosario Tumino³⁶, Salvatore Panico³⁷, Verena Katzke¹³, Xuan Ren¹⁴, Paolo Vineis³⁸, Pietro Ferrari¹, Carl Lachat^{2*}, Inge Huybrechts^{1*}

*Joint last authors

¹Nutrition and Metabolism Branch, International Agency for Research on Cancer, Lyon, France

²Department of Food Technology, Safety and Health, Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium

³Food and Nutrition Division, Food and Agriculture Organization of the United Nations, Rome, Italy



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980025101067

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

⁴Department of Research, Cancer Registry of Norway, Norwegian Institute of Public Health, Oslo, Norway

⁵Department of Nutrition, Oslo New University College, Oslo, Norway

⁶Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK

⁷Copernicus Institute of Sustainable Development, Utrecht University

⁸Université Sorbonne Paris Nord and Université Paris Cité, INSERM, INRAE, CNAM, Centre of Research in Epidemiology and Statistics (CRESS), Nutritional Epidemiology Research Team (EREN), F-93017 Bobigny, France

⁹Public Health Policy Evaluation Unit, School of Public Health, Imperial College London, London, United Kingdom

¹⁰NOVA National School of Public Health, Public Health Research Centre, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Portugal

¹¹Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, UK

¹²Medical Research Council Centre for Environment and Health, Imperial College London, London, United Kingdom

¹³Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

¹⁴Paris-Saclay University, UVSQ, Inserm, Gustave Roussy, CESP, Villejuif, France

¹⁵Department of Public Health, Aarhus University, Aarhus, Denmark

¹⁶Steno Diabetes Center Aarhus, Aarhus University Hospital, Aarhus, Denmark

¹⁷Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany

¹⁸Department of Community Medicine, UiT The Arctic University of Norway, Tromsø, Norway

¹⁹CIBER in Epidemiology and Public Health (CIBERESP), Madrid, Spain

²⁰Escuela Andaluza de Salud Pública (EASP), 18011 Granada, Spain

²¹Instituto de Investigación Biosanitaria ibs. GRANADA, 18012 Granada, Spain

²²Institute of Nutritional Science, University of Potsdam, Nuthetal, Germany

²³Fondazione IRCCS Istituto Nazionale dei Tumori di Milano Via Venezian, Milan, Italy

²⁴Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands

²⁵Ministry of Health of the Basque Government, Sub-Directorate for Public Health and Addictions of Gipuzkoa, San Sebastián, Gipuzkoa, Spain

²⁶Biogipuzkoa Health Research Institute, Group of Epidemiology of Chronic and Communicable Diseases, San Sebastián, Gipuzkoa, Spain

²⁷Department of Diagnostics and Intervention, Oncology, Umeå University, SE-90187 Umeå, Sweden

²⁸Department of Internal Medicine and Clinical Nutrition, University of Göteborg, SE-405 30 Göteborg, Sweden

²⁹Danish Cancer Institute, Copenhagen, Denmark

³⁰Department of Public Health, University of Copenhagen, Denmark

³¹Department of Health Sciences. University of Eastern Piedmont. Novara, Italy

³²Unit of Epidemiology, Local Health Unit of Novara, Novara, Italy.

³³International Agency for Research on Cancer, Lyon, France

³⁴Instituto de Salud Pública y Laboral de Navarra, 31003 Pamplona, Spain

³⁵Navarra Institute for Health Research (IdiSNA), 31008 Pamplona, Spain

³⁶Hyblean Association for Epidemiology Research, AIRE ONLUS Ragusa, Italy

³⁷School of Medicine, Federico II University, Naples, Italy

³⁸Cancer Epidemiology and Prevention Research Unit, School of Public Health, Imperial College London, UK

Corresponding author: Jeroen Berden, jeroen.berden@ugent.be, +32496237676, International Agency for Research on Cancer, Section of Nutrition and Metabolism, 25 Avenue Tony Garnier, CS 90627, 69366 LYON CEDEX 07, France.

Short title: Food processing, biodiversity and environment

Keywords: food processing, environmental impact, food biodiversity, ultra-processed foods

Financial Support: The coordination of EPIC is financially supported by the International Agency for Research on Cancer (IARC) and the Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London which has additional infrastructure support provided by the NIHR Imperial Biomedical Research Centre (BRC).

The national cohorts are supported by: Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Éducation Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM) (France); German Cancer Aid, German Cancer Research Center (DKFZ), German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Federal Ministry of Education and Research (BMBF) (Germany); Associazione Italiana per la Ricerca sul Cancro-AIRC-Italy, Compagnia di SanPaolo and National Research Council (Italy); Dutch Ministry of Public Health, Welfare and Sports (VWS), Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF) (The Netherlands); Health Research Fund (FIS) - Instituto de Salud Carlos III (ISCIII), Regional Governments of Andalucía, Asturias, Basque Country, Murcia and Navarra, and the Catalan Institute of Oncology - ICO (Spain); Swedish Cancer Society, Swedish Research Council and County Councils of Skåne and Västerbotten (Sweden); Cancer Research UK (14136 to EPIC-Norfolk; C8221/A29017 to EPIC-Oxford), Medical Research Council (1000143 to EPIC-Norfolk; MR/M012190/1 to EPIC-Oxford). (United Kingdom), UiT The Arctic University of Norway. EKC was supported by the Ministry of National Education, Türkiye, through the YLSY International Graduate Education Scholarship Programme.

Funding for grant number IIG_FULLL_2020_034 was obtained from Wereld Kanker Onderzoek Fonds (WKOF), as part of the World Cancer Research Fund International grant programme. Funding for grant IIG_FULLL_2020_033 was obtained from World Cancer Research Fund (WCRF UK), as part of the World Cancer Research Fund International grant programme. Researchers were independent from the funders. Funders had no role in the collection, analysis, and interpretation of data, the writing of the manuscript, and the decision to submit the article for publication. PV and CJM are supported by the NIHR GHRC on NCDs and Environmental Change (NIHR203247) using UK aid from the UK Government to support global health research. Researchers were independent from the funders. Funders had no role in the collection, analysis, and interpretation of data, the writing of the manuscript, and the decision to submit the article for publication.

Conflict of Interest: The authors have declared that no competing interests exist.

IARC disclaimer: Where authors are identified as personnel of IARC/WHO, the authors alone are responsible for the views expressed in this article, and they do not necessarily represent the decisions, policy, or views of IARC/WHO.

FAO disclaimer: The views expressed in this publication are those of the authors and do not necessarily reflect the views or policies of the Food and Agriculture Organization of the United Nations.

Author Contribution:

Authorship: Conceptualization: Jeroen Berden, Carl Lachat, Inge Huybrechts

Data curation: Inge Huybrechts, Geneviève Nicolas, Pietro Ferrari

Formal analysis: Jeroen Berden, Inge Huybrechts, Geneviève Nicolas

Funding acquisition: Giles T. Hanley-Cook, Inge Huybrechts, Carl Lachat, Mathilde Touvier

Investigation: Jeroen Berden, Giles T. Hanley-Cook, Bernadette Chimera, Dagfinn Aune, Maria Gabriela M. Pinho, Geneviève Nicolas, Bernard Srour, Christopher J. Millett, Emine Koc Cakmak, Emmanuelle Kesse-Guyot, Esther M. González-Gil, Eszter P. Vamos, Jessica Blanco Lopez, Julia Baudry, Justine Berlivet, Kiara Chang, Mathilde Touvier, Charlotte Le Cornet, Chloé Marques, Christina C. Dahm, Daniel B. Ibsen, Franziska Jannasch, Guri Skeie, Maria-José Sanchez, Matthias B. Schulze, Sara Grioni, Yvonne T. van der Schouw, Ana M. Jimenez Zabala, Anna Winkvist, Anne Tjønneland, Carlotta Sacerdote, Cecilie Kyrø, Elisabeth Weiderpas, Marcela Guevara, Pauline Frenoy, Rosario Tumino, Salvatore Panico, Verena Katzke, Xuan Ren, Paolo Vineis, Pietro Ferrari, Carl Lachat, Inge Huybrechts.

Methodology: Jeroen Berden, Carl Lachat, Inge Huybrechts

Project administration: Jeroen Berden, Carl Lachat, Inge Huybrechts

Software: Jeroen Berden

Resources: Jeroen Berden, Giles T. Hanley-Cook, Bernadette Chimera, Dagfinn Aune, Maria Gabriela M. Pinho, Geneviève Nicolas, Bernard Srour, Christopher J. Millett, Emine Koc Cakmak, Emmanuelle Kesse-Guyot, Esther M. González-Gil, Eszter P. Vamos, Jessica Blanco Lopez, Julia Baudry, Justine Berlivet, Kiara Chang, Mathilde Touvier, Charlotte Le Cornet, Chloé Marques, Christina C. Dahm, Daniel B. Ibsen, Franziska Jannasch, Guri Skeie, Maria-José Sanchez, Matthias B. Schulze, Sara Grioni, Yvonne T. van der Schouw, Ana M. Jimenez Zabala, Anna Winkvist, Anne Tjønneland, Carlotta Sacerdote, Cecilie Kyrø, Elisabeth Weiderpas, Marcela Guevara, Pauline Frenoy, Rosario Tumino, Salvatore Panico, Verena Katzke, Xuan Ren, Paolo Vineis, Pietro Ferrari, Carl Lachat, Inge Huybrechts.

Supervision: Carl Lachat, Inge Huybrechts

Visualization: Jeroen Berden

Validation: Geneviève Nicolas, Carl Lachat, Inge Huybrechts

Writing original draft: Jeroen Berden

Writing paper draft: Jeroen Berden, Giles T. Hanley-Cook, Bernadette Chimera, Dagfinn Aune, Maria Gabriela M. Pinho, Geneviève Nicolas, Bernard Srouf, Christopher J. Millett, Emine Koc Cakmak, Emmanuelle Kesse-Guyot, Esther M. González-Gil, Eszter P. Vamos, Jessica Blanco Lopez, Julia Baudry, Justine Berlivet, Kiara Chang, Mathilde Touvier, Charlotte Le Cornet, Chloé Marques, Christina C. Dahm, Daniel B. Ibsen, Franziska Jannasch, Guri Skeie, Maria-José Sanchez, Matthias B. Schulze, Sara Grioni, Yvonne T. van der Schouw, Ana M. Jimenez Zabala, Anna Winkvist, Anne Tjønneland, Carlotta Sacerdote, Cecilie Kyrø, Elisabette Weiderpas, Marcela Guevara, Pauline Frenoy, Rosario Tumino, Salvatore Panico, Verena Katzke, Xuan Ren, Paolo Vineis, Pietro Ferrari, Carl Lachat, Inge Huybrechts.

Ethical Standards Disclosure: The present study is relevant to guide local agrifood systems transformation and nutrition research. Local research teams were invited to contribute to the manuscript and were included as co-authors in accordance with ICJME criteria.

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by local ethics committees and by the Internal Review Board of the International Agency for Research on Cancer. Written informed consent was obtained from all subjects. For participants who were illiterate or otherwise unable to provide written consent, informed consent was obtained from a legal guardian or an appropriate representative on their behalf.

Data availability: EPIC data and biospecimens are available to investigators in the context of research projects that are consistent with the legal and ethical standard practices of IARC/WHO and the EPIC Centres. The use of a random sample of anonymized data from the EPIC study can be requested by contacting epic@iarc.fr. For information on the EPIC data access policy and on how to submit an application for gaining access to EPIC data and/or biospecimens, please follow the instructions at iarc.who.int

Abstract

Objective: While associations of ultra-processed food (UPF) consumption with adverse health outcomes are accruing, its environmental and food biodiversity impacts remain underexplored. This study examines associations between UPF consumption and dietary greenhouse gas emissions (GHGe), land use, and food biodiversity.

Design: Prospective cohort study. Linear mixed models estimated associations between UPF intake (grams/day and kcal/day) and GHGe (kg CO₂-equivalents/day), land use (m²/day), and dietary species richness (DSR). Substitution analyses assessed the impact of replacing UPFs with unprocessed or minimally processed foods.

Participants: 368,733 participants in the European Prospective Investigation into Cancer and Nutrition (EPIC) study.

Setting: Europe

Results: Stronger associations were found for UPF consumption in relation with GHGe and land use compared to unprocessed or minimally processed food consumption. Substituting UPFs with unprocessed or minimally processed foods was associated with lower GHGe (8.9%; 95%CI: -9.0; -8.9) and land use (9.3%; -9.5; -9.2) when considering consumption by gram per day and higher GHGe (2.6%; 95% CI: 2.5; 2.6) and land use (1.2%; 1.0; 1.3) when considering consumption in kilocalories per day. Substituting UPF by unprocessed or minimally processed foods led to negligible differences in DSR, both for consumption in grams (-0.1%; -0.2; -0.1) and kilocalories (1.0%; 1.0; 1.1).

Conclusion: UPF consumption was strongly associated with GHGe and land use as compared to unprocessed or minimally processed food consumption, while associations with food biodiversity were marginal. Substituting UPFs with unprocessed or minimally processed foods resulted in differing directions of associations with environmental impacts, depending on whether substitutions were weight- or calorie-based.

1. Introduction

The food system's environment impact has become a pressing concern due to its contributions to greenhouse gas emissions (GHGe), land use, and biodiversity loss¹. Intensive agricultural practices, especially monocultures like maize, wheat, and soy, degrade ecosystems and narrow crop diversity. Ultra-processed foods (UPFs), composed largely of ingredients produced from these high yield crops and livestock, have been indicated to have a negative impact on the environment due to their contribution to limited crop diversity and increased vulnerability to environmental pressures². In addition, many UPFs are characterized by hyperpalatability, low satiety potential, and heavy marketing that can encourage overconsumption, leading to excessive food production and associated environmental pressures, while also contributing to significant public health challenges^{3,4}.

UPFs have been linked to negative health outcomes such as obesity, cardiovascular diseases, depressive symptoms, and certain cancers⁵. Consequently, countries like Mexico have incorporated recommendations to limit UPF consumption in dietary guidelines⁶. However, the environmental impacts of UPFs have received less attention, and the potential implications of substituting UPFs with unprocessed or minimally processed foods remain underexplored. With diets shifting toward greater UPF consumption globally, understanding their impact on the environment is critical, particularly in terms of GHGe, land use, and preservation of food biodiversity⁷⁻⁹. This convergence suggests that UPF-driven overconsumption represents a shared pathophysiological mechanism underlying both human and environmental health. The same hyperpalatable formulations, low satiety signals, and marketing strategies that promote excessive energy intake, could simultaneously drive increased food demand and production, amplifying environmental pressures. This dual pathway through overconsumption represents a novel framework for understanding how food processing impacts both human and planetary health through a common mechanism.

This study examined the relationship between dietary intake across food processing levels and environmental outcomes—specifically GHGe, land use, and food biodiversity—and evaluates the potential impact of substituting minimally processed foods for UPFs among adults in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohort.

2. Methods

2.1. The EPIC cohort

The EPIC cohort is a large multicentre cohort examining the links between metabolic, lifestyle, and environmental factors of cancer and chronic diseases. Between 1991 and 2000, over 500,000 participants aged 25–70 were recruited across 23 centres in 10 European countries: Denmark, France, Germany, Greece, Italy, the Netherlands, Norway, Spain, Sweden, and the United Kingdom. Dietary intake at enrolment was assessed using validated, country-specific questionnaires capturing habitual consumption over the past 12 months¹⁰. In order to study associations in a disease-free population participants with missing dietary data, extreme energy intake-to-requirement ratios, lack of follow-up, or prevalent diseases at baseline were excluded. Due to administrative constraints, cohorts from Greece, Norway, and Sweden were excluded, resulting in 368,733 participants (Supplemental Figure 1).

2.2. Dietary Assessment

In the 1990s, participants' usual food intake over the previous 12 months was assessed at baseline with country-specific dietary questionnaires. Depending on the study centre, quantitative dietary questionnaires, semi-quantitative food-frequency questionnaires, or a combination of semi-quantitative food-frequency questionnaires and 7-day food records were used. Data on frequencies, portion sizes, or intakes in grams per day were stored in a central IARC database¹⁰. Post-harmonization of dietary data was conducted, following standardized procedures (e.g., disaggregating recipes into ingredients), to obtain a standardized food list for which the level of detail is comparable between countries. The EPIC food composition database comprises more than 11,000 food and beverage items reflecting the specificities of each country.

2.3. Exposure - Nova classification

Standardized EPIC food items were categorized by processing level using the Nova classification: Nova 1 (unprocessed or minimally processed foods, e.g., fruits, vegetables), Nova 2 (processed culinary ingredients, e.g., oils, sugar), Nova 3 (processed foods, e.g., cheese, bread), and Nova 4 (UPFs, e.g., soft drinks, flavoured yoghurts). Since the Nova classification system was developed after the EPIC dietary data collection, there was some uncertainty in classifying certain food items according to their level of food processing. Therefore, three classification

scenarios were developed to address this uncertainty, a lower, middle, and upper-bound scenario. This study used the most probable, middle-bound scenario¹¹. Dietary contribution from each Nova class was expressed in both grams and kcal per day, as grams reflect absolute consumption, while kcal accounts for energy density, providing complementary insights into environmental impacts.

2.4. Outcomes - Environmental impacts and food biodiversity

Environmental outcomes were assessed using the SHARP indicators database, which estimates GHGe and land use from life cycle assessment data encompassing production, packaging, transport, and preparation¹². Food items were matched between the EPIC and SHARP databases using EFSA's FoodEx2 base-term codes¹³. Diet-related GHGe and land use were computed for each individual by summing the amounts for all foods consumed; GHGe was expressed as kg CO₂-equivalents per day and land use as m² per day^{12,13}. Food biodiversity was quantified using dietary species richness (DSR), defined as the count of unique biological species consumed across foods, beverages, and mixed dishes¹⁴. Composite foods were decomposed into ingredients using standard recipes and foods consumed "never or less than once per month" were not considered in the DSR computation.¹⁴

2.5. Study covariates

Sociodemographic and anthropometric covariates included in the models were: age at recruitment, body mass index height, sex, educational level, smoking status at baseline, physical activity using the Cambridge index, and alcohol intake.

2.6. Statistical analysis

Consumption of the Nova classes (gram/day or kcal/day) was modelled as continuous variables. Multivariable mixed linear models with random intercepts for study centres and adjustment for sociodemographic and anthropometric variables were fitted to assess associations between Nova class consumption, GHGe, land use, and DSR. Additive models assessed associations of the additional consumption of a Nova class. For this, weight- and energy-based all-component models were constructed, mutually adjusting for each Nova class, to account for the total weight or energy intake¹⁵. Additionally, substitution analyses were performed, using the 'leave-one-out' method estimated associations of replacing a specific amount of Nova 4 with Nova 1, by keeping

total intake constant¹⁶. For instance, the substitution of Nova 4 by Nova 1 in GHGe can be parameterised as:

$$\text{GHGe} = \alpha + \gamma_1 \text{Nova 1} + \gamma_2 \text{Nova 2} + \gamma_3 \text{Nova 3} + \gamma_4 \text{total intake} + \text{covariates} \\ + (1 \mid \text{Study Centre}) + \varepsilon$$

Here, γ_1 represents the relative estimate for replacing a quantity of Nova 4 with an equivalent amount of Nova 1, keeping the total intake constant.

Estimates were expressed as: 1) a 1-standard deviation (SD) increment in consumption of a Nova class, or 2) a 10% increase from the mean absolute total dietary intake. To interpret the results as percentage differences, these estimates were divided by the mean value of the respective outcome measure.

Sensitivity analyses included baseline models only mutually adjusted for each Nova class and main models further adjusted for the Mediterranean diet score (0–18 points)¹⁷. Statistical analyses were performed in RStudio (v4.0.4.1) with two-sided testing, and P-values <0.05 were considered statistically significant.

3. Results

3.1. Sample characteristics

This study included 368,733 participants from the EPIC cohort, of whom 259,268 (70.3%) were females. The mean (SD) age at recruitment was 51.3 (9.9) years, and the average BMI was 25.4 (4.3) kg/m² at baseline. On average, participants consumed 364 gram (278) or 672.9 kcal (412.0) of UPFs daily, representing 12.9% (8.5) of total intake by weight and 30.5% (15.3) by energy. Mean dietary GHGe and land use were 5.3 (1.82) kg CO₂-equivalents per day and 6.9 (2.6) m² per day, respectively. Average DSR was 68.2 (15.2) species per year (**Table 1**).

3.2. Associations between Nova class consumptions and GHGe, land use and food biodiversity




Figure 1 illustrates the percentage difference relative to the mean of GHGe, land use, and DSR associated with higher consumption of each Nova class. A 1-SD increment in consumption of each Nova class, either in gram or kcal per day, was associated with significantly higher GHGe, land use, and DSR, with Nova 4 consumption being more strongly associated with GHGe and land use compared to Nova 1. To illustrate, a 1-SD increment of Nova 4 consumption in kcal per day was related to 15.8% (95%CI: 15.8; 16.0) higher GHGe, 16.9% (16.9; 17.1) higher land use, and 1.0% (0.9;1.1) higher DSR, while for Nova 1 this was 13.8% (13.8; 14.0) for GHGe, 12.8% (12.7; 14.0) for land use. Similar findings were reported for consumption of the different Nova classes in grams/day. Strengths of associations differed within Nova 4 subgroups, with animal-based products showing the strongest positive associations with GHGe and land use, while plant-based alternatives and savoury snacks showed the weakest associations (**Supplemental table 1**).

3.3. Substitution of ultra-processed with unprocessed or minimally processed foods

10% of the mean total intake in grams per day substitution of Nova 4 substitution with Nova 1 was associated with 8.9% (95%CI: -9.0; -8.9) lower GHGe and 9.3% (-9.5; -9.2) lower land use (**Figure 1**). However, such a substitution was related to marginally lower DSR (-0.1%; -0.2; -0.1). Conversely, a Nova 4 substitution with Nova 1, 10% of the mean total intake in kcal per

day, was associated with higher GHGe (2.6%; 2.5: 2.6), land use (1.2%; 1.0; 1.3), and DSR (1.0%; 1.0; 1.1) (**Figure 1**).

Sensitivity analysis confirmed our main findings (data not shown).

4. Discussion

This study found that higher UPF consumption was more strongly associated with increased dietary GHGe and land use compared to unprocessed or minimally processed foods. For DSR, associations were shown to be marginal. Energy-based substitution of UPFs with unprocessed or minimally processed foods were associated with higher environmental impacts, whereas weight-based substitutions were associated with lower environmental impacts.

These discrepancies likely stem from the higher energy density of UPFs. Energy-based substitutions require larger quantities of unprocessed or minimally processed foods to achieve isocaloric substitutions, potentially increasing environmental impacts¹⁸. Research suggests that individuals consuming diets high in unprocessed or minimally processed foods tend to have lower energy intake compared to those with UPF-rich diets, meaning isocaloric substitution may not fully capture these differences¹⁹. In contrast, weight-based substitutions, which emphasize food weight rather than caloric equivalence, show environmental benefits that align with UPFs' well-documented tendency to promote overconsumption through their hyper palatability, low satiety, softer textures requiring less chewing, widespread availability, and lower cost per calorie, which could lead to excessive energy intake, contributing to rising obesity rates^{3,4,19}. Such overconsumption drives higher demand for foods, amplifying environmental impacts further. Additionally, while low-impact plant-based UPFs have lower environmental footprints, animal-based UPFs remain highly impactful, underlining the importance of considering UPF subgroups^{20,21}. These findings support the hypothesis that overconsumption serves as a critical link between UPF consumption and environmental harm, paralleling established mechanisms for UPF-associated health risks.

Additionally, negligible DSR differences were observed when substituting UPFs with unprocessed or minimal foods, diverging from findings in Brazilian diets where UPFs involved fewer species⁸. This discrepancy may reflect methodological differences: the Brazilian study examined species diversity within UPF products at the food system level, while our analysis assessed how individual dietary patterns relate to overall species consumption.

Our findings suggest that food biodiversity operates independently from processing level in individual diets. Substituting UPFs with unprocessed foods may not increase species diversity if individuals simply as individuals might simply consume larger quantities of the same limited set of species they already consume. Therefore, reducing UPF consumption alone may be insufficient to improve dietary biodiversity without concurrent efforts to promote species diversification. Alternatively, UPF-driven overconsumption may increase total food intake, maintaining dietary species richness through higher consumption volumes rather than dietary diversification.

Limited observational evidence on UPFs' environmental impacts exists, with most insights coming from life cycle assessments²². In a French cohort study, it was found that UPFs accounted for 19% of energy intake in the diet and contributed to 24% of greenhouse gas emissions (GHGe), 23% of land use, and 26% of energy demand. These highlight the significant environmental burden associated with diets rich in UPFs, with higher contributions from post-farm stages, in particular processing regarding energy demand²³. A longitudinal study showed reducing UPF consumption lowered GHGe and energy demand, but increased water use²⁴. Our study is unique in its large, diverse European cohort, allowing a comprehensive assessment of food processing levels and substitution effects.

Several limitations must be acknowledged. The EPIC cohort may differ substantially from current European populations. UPF intake has risen dramatically—from approximately one-third of energy intake in our cohort to over half in recent studies, due to changes in food environments and consumption patterns²⁵. Although educational attainment has increased across EU member states, this has not corresponded with expected reductions in UPF consumption, suggesting altered socioeconomic determinants of dietary choices²⁶. The shift toward sedentary lifestyles correlates with increased convenience food reliance, while younger populations exhibit greater price sensitivity toward UPF products^{27,28}. These transitions suggest our cohort likely underestimates the environmental impacts of contemporary European diets. Misclassification within the Nova system and reliance on SHARP database estimates, which lack country specificity and farming method variations, may introduce error. Additionally, many UPF-specific ingredients (e.g., additives) lack environmental impact assessments, and UPFs typically rely on more intensively produced commodity ingredients than non-UPFs, differences our analysis cannot fully capture. These errors might attenuate the true associations due to non-differential

measurement error. Variations in dietary assessment methods and the number of items included between centres could also affect DSR, and taxonomic limitations hinder further analysis of food biodiversity. The questionnaires did not distinguish between homemade and industrially processed foods, which could overlook ingredient differences leading to varying environmental impacts. Lastly, this study compared individuals rather than actual substitutions, and context-specific factors such as preparation time, cost, and food safety may influence dietary shifts and willingness to make substitutions^{29,30}. For instance, while unprocessed or minimally processed foods are often more nutrient-dense, UPFs offer greater accessibility and food safety³¹.

In conclusion, UPF consumption was more strongly associated with GHGe and land use as compared to unprocessed or minimally processed food consumption, while associations with food biodiversity were marginal. Substituting UPFs with unprocessed or minimally processed foods resulted in differing directions of associations with environmental impacts, depending on whether substitutions were weight- or calorie-based.

Table 1. Baseline characteristics of 368,733 middle-aged adults enrolled in the European Prospective Investigation into Cancer and Nutrition (EPIC) study.

	Mean (SD) / N (%)
Nova 4 (% gram per day)	12.93 (8.52)
Nova 3 (% gram per day)	14.28 (10.56)
Nova 2 (% gram per day)	1.23 (1.06)
Nova 1 (% gram per day)	71.56 (12.50)
Nova 4 (% kcal per day)	30.55 (15.3)
Nova 3 (% kcal per day)	25.81 (12.04)
Nova 2 (% kcal per day)	7.95 (6.23)
Nova 1 (% kcal per day)	35.70 (10.62)
Dietary greenhouse gas emissions (kg CO₂ equivalents per day)	5.30 (1.82)
Dietary land use (m² per day)	6.86 (2.62)
DSR (count of unique species consumed per year)	68.22 (15.22)
Age at recruitment (years)	51.29 (9.90)
Body mass index (kg per m²)	25.35 (4.25)
Height (cm)	165.67 (8.92)
Sex	
Male	109,465 (29.7%)
Female	259,268 (70.3%)
Country	
Denmark	55,014 (14.9%)
France	67,920 (18.4%)
Germany	49,352 (13.4%)
Italy	44,547 (12.1%)
Spain	39,990 (10.8%)
The Netherlands	36,538 (9.9%)
United Kingdom	75,372 (20.4%)
Education level	

None or primary school completed	102,198 (27.7%)
Technical/professional school	80,266 (21.8%)
Secondary school	75,288 (20.4%)
Longer education (including university degree)	94,312 (25.6%)
Unknown	16,669 (4.5%)
Smoking status	
Never	184,435 (50.0%)
Current	99,923 (27.1%)
Former	78,175 (21.2%)
Unknown	6,200 (1.7%)
Cambridge physical activity index	
Inactive	76,776 (20.8%)
Moderately inactive	125,817 (34.1%)
Moderately active	88,476 (24.0%)
Active	70,923 (19.2%)
Unknown	6,741 (1.8%)
Alcohol intake	
Non-drinker	44,761 (12.1%)
> 0 to 6 gram per day	96,866 (26.3%)
> 6 to 12 gram per day	96,048 (26.0%)
> 12 to 24 gram per day	64,086 (17.4%)
> 24 gram per day	66,972 (18.2%)

DSR, dietary species richness. Nova 1, unprocessed or minimally processed foods. Nova 2, processed culinary ingredients. Nova 3, processed foods. Nova 4, ultra-processed foods

References

1. Springmann M, Clark M, Mason-D'Croz D, et al. Options for keeping the food system within environmental limits. *Nature*. 2018;562:519-525.
2. Leite FHM, Khandpur N, Andrade GM, et al. Ultra-processed foods should be central to global food systems dialogue and action on biodiversity. *BMJ Glob Health*. 2022;7:e008269.
3. Anastasiou K, Baker P, Hadjikakou M, et al. Conceptualising the drivers of ultra-processed food production and consumption and their environmental impacts: A group model-building exercise. *Glob Food Sec*. 2023;37:100688.
4. Seferidi P, Scrinis G, Huybrechts I, et al. The neglected environmental impacts of ultra-processed foods. *Lancet Planet Health*. 2020;4:e437-e438.
5. Lane MM, Gamage E, Du S, et al. Ultra-processed food exposure and adverse health outcomes: umbrella review of epidemiological meta-analyses. *BMJ*. 2024:e077310.
6. Colansa. Brazil's Food Guide turns 10 [Internet]. 2024 [cited 2024]. Available from: <https://colansa.org/en/library/brazils-food-guide-turns-10/>
7. Popkin BM, Ng SW. The nutrition transition to a stage of high obesity and noncommunicable disease prevalence dominated by ultra-processed foods is not inevitable. *Obes Rev*. 2022;23.
8. Leite F, Khandpur N, Andrade G, et al. A multi-step methodology to identify the food biodiversity that underlies brazilian diets. *Popul Med*. 2023;5.
9. FAO. The State of the World's Biodiversity for Food and Agriculture. Rome: FAO; 2019.
10. Riboli E, Hunt KJ, Slimani N, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr*. 2002;5:1113-1124.
11. Huybrechts I, Rauber F, Couto E, et al. Characterization of the degree of food processing in the European Prospective Investigation into Cancer and Nutrition: application of the Nova classification and validation using selected biomarkers of food processing. *Front Nutr*. 2022;9.
12. Mertens E, Kuijsten A, Dofková M, et al. SHARP-Indicators Database towards a public database for environmental sustainability. *Data Brief*. 2019;27:104617.
13. EFSA. The food classification and description system FoodEx 2 (revision 2). *EFSA Support Publ*. 2015;12.

14. Hanley-Cook GT, Huybrechts I, Biessy C, et al. Food biodiversity and total and cause-specific mortality in 9 European countries: An analysis of a prospective cohort study. *PLoS Med.* 2021;18:e1003834.
15. Tomova GD, Arnold KF, Gilthorpe MS, Tennant PW. Adjustment for energy intake in nutritional research: a causal inference perspective. *Am J Clin Nutr.* 2022;115:189-198.
16. Tomova GD, Gilthorpe MS, Tennant PW. Theory and performance of substitution models for estimating relative causal effects in nutritional epidemiology. *Am J Clin Nutr.* 2022;116:1379-1388.
17. Couto E, Boffetta P, Laggiou P, et al. Mediterranean dietary pattern and cancer risk in the EPIC cohort. *Br J Cancer.* 2011;104:1493-1499.
18. Gupta S, Hawk T, Aggarwal A, Drewnowski A. Characterizing Ultra-Processed Foods by Energy Density, Nutrient Density, and Cost. *Front Nutr.* 2019;6.
19. Hall KD, Ayuketah A, Brychta R, et al. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab.* 2019;30:67-77.e3.
20. Clark M, Domingo NGG, Colgan K, et al. Estimating the environmental impacts of 57,000 food products. *Proc Natl Acad Sci USA.* 2022;119.
21. Cordova R, Viallon V, Fontvieille E, et al. Consumption of ultra-processed foods and risk of multimorbidity of cancer and cardiometabolic diseases: a multinational cohort study. *Lancet Reg Health Eur.* 2023;35:100771.
22. Jones AD, Hoey L, Blesh J, et al. A Systematic Review of the Measurement of Sustainable Diets. *Adv Nutr.* 2016;7:641-664.
23. Kesse-Guyot E, Allès B, Brunin J, et al. Environmental impacts associated with UPF consumption: which food chain stages matter the most? Findings from a representative sample of French adults. *medRxiv.* 2022. doi:10.1101/2022.05.28.22275717.
24. García S, Bouzas C, Mateos D, et al. Ultra-processed foods consumption as a promoting factor of greenhouse gas emissions, water, energy, and land use: A longitudinal assessment. *Sci Total Environ.* 2023;891:164417.
25. Monteiro CA, Moubarac JC, Levy RB, et al. Household availability of ultra-processed foods and obesity in nineteen European countries. *Public Health Nutr.* 2018;21:18-26.

26. Dicken SJ, Qamar S, Batterham RL. Who consumes ultra-processed food? A systematic review of sociodemographic determinants of ultra-processed food consumption from nationally representative samples. *Nutr Res Rev.* 2024;37:416-456.
27. Costa CS, Del-Ponte B, Assunção MCF, Santos IS. Consumption of ultra-processed foods and body fat during childhood and adolescence: a systematic review. *Public Health Nutr.* 2018;21:148-159.
28. Machado PP, Steele EM, Levy RB, et al. Ultra-processed food consumption and obesity in the Australian adult population. *Nutr Diabetes.* 2020;10:39.
29. Gupta S, Hawk T, Aggarwal A, Drewnowski A. Characterizing Ultra-Processed Foods by Energy Density, Nutrient Density, and Cost. *Front Nutr.* 2019;6.
30. Wolfson JA, Martinez-Steele E, Tucker AC, Leung CW. Greater Frequency of Cooking Dinner at Home and More Time Spent Cooking Are Inversely Associated With Ultra-Processed Food Consumption Among US Adults. *J Acad Nutr Diet.* 2024. doi:10.1016/j.jand.2024.03.005.
31. Amorim A, Silva VL, Sobral PJ do A. Food Processing: An overview on links between safety, security, supply chains, and NOVA classification. *Clean Circ Bioeconomy.* 2023;5:100047.