

Local nearby bifurcations lead to synergies in critical slowing down: The case of mushroom bifurcations

Mariona Fucho-Rius ^{*}

Departament de Matemàtiques, Universitat Politècnica de Catalunya (UPC), Pau Gargallo 14, 08028 Barcelona, Spain

Smitha Maretvadakethope ^{*}

Department of Life Sciences, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom

Àlex Haro 

*Departament de Matemàtiques i Informàtica, Universitat de Barcelona (UB), Gran Via de les Corts Catalanes 585, 08007 Barcelona, Spain
and Centre de Recerca Matemàtica, Edifici C, 08193 Cerdanyola del Vallès, Barcelona, Spain*

Tomás Alarcón 

*ICREA, Passeig Lluís Companys, 23 08010 Barcelona, Spain;
Centre de Recerca Matemàtica, Edifici C, 08193 Cerdanyola del Vallès, Barcelona, Spain;
and Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Cerdanyola del Vallès, Barcelona, Spain*

Josep Sardanyés [†]

Centre de Recerca Matemàtica, Edifici C, 08193 Cerdanyola del Vallès, Barcelona, Spain

Rubén Pérez-Carrasco [‡]

Department of Life Sciences, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom



(Received 9 August 2024; accepted 30 January 2025; published 28 February 2025)

The behavior of nonlinear systems near critical transitions has significant implications for stability, transients, and resilience in complex systems. Transient times, τ , become extremely long near phase transitions (or bifurcations) in a phenomenon known as critical slowing down, and are observed in electronic circuits, circuit quantum electrodynamics, ecosystems, and gene regulatory networks. Critical slowing down typically follows universal laws of the form $\tau \sim |\mu - \mu_c|^\beta$, with μ being the control parameter and μ_c its critical value. For instance, $\beta = -1/2$ close to saddle-node bifurcations. Despite intensive research on slowing down phenomena for single bifurcations, both local and global, the behavior of transients when several bifurcations are close to each other remains unknown. Here, we investigate transients near two saddle-node bifurcations merging into a transcritical one. Using a nonlinear gene-regulatory model and a normal form exhibiting a mushroom bifurcation diagram we show, both analytically and numerically, a synergistic, i.e., nonadditive, lengthening of transients due to coupled ghost effects and transcritical slowing down. We also show that intrinsic and extrinsic noise play opposite roles in the slowing down of the transition, allowing us to control the timing of the transition without compromising the precision of timing. This establishes molecular strategies to generate genetic timers with transients much larger than the typical timescales of the reactions involved.

DOI: [10.1103/PhysRevE.111.024213](https://doi.org/10.1103/PhysRevE.111.024213)

I. INTRODUCTION

Complex systems close to a phase transition or a bifurcation undergo critical slowing down [1,2]. This phenomenon

lengthens the relaxation times of a system following a disturbance, indicating reduced resilience and increased susceptibility to perturbations. Critical slowing down has been described in a multitude of nonlinear physical systems in optics [3], electronic circuits [4], circuit quantum electrodynamics [5,6], ferromagnetic materials [7], magnetic quantum phase transitions [8], and models of charge density waves [9]. The investigation of critical slowing down has also been very intensive in the field of ecology, where many works have provided evidence of these phenomena in experimental and field data [1,10–12].

The transient time, τ , for the case of orbits in a small neighborhood of an equilibrium close to a bifurcation typi-

^{*}These authors contributed equally to this work.

[†]Contact author: jsardanyes@crm.cat

[‡]Contact author: r.perez-carrasco@imperial.ac.uk

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

cally follows laws of the form $\tau \sim |\mu - \mu_c|^\beta$ for deterministic systems [9,13–16]. Here, μ is the control (bifurcation) parameter, and μ_c is the critical value at which the bifurcation occurs. The exponents β are universal and they are known for local bifurcations. For example, $\beta = -1$ for pitchfork and transcritical bifurcations [15–17], and $\beta = -1/2$ for saddle-node (SN) bifurcations [13,14], with examples of applications in deterministic models of cooperation [14,18–21]. Similar power laws for transients with exponents -1 have been recently identified for global bifurcations such as the trans-heteroclinic bifurcation introduced in [22,23], and in bifurcations involving the destruction of quasineutral curves [24]. Recent research has also explored the effects of intrinsic noise on such transients, finding more complicated scaling laws for the SN bifurcation [25,26]. Such delays for SN bifurcations are caused by so-called “ghosts”. This phenomenon appears right after the SN bifurcation where the two equilibria have collapsed and moved onto the complex plane. However, in the vicinity of the region where the SN occurs, the vector field is close to zero and therefore the system undergoes slowing down. Here, we will study a scenario where the effect of transients close to a transcritical bifurcation occurs concomitantly with the ghost phenomena appearing in two close SN bifurcations. This coupling produces a nonlinear (synergistic) increase in transient times, causing a nonadditive effect of the duration of transients.

Phenomena involving the slowing down near SN bifurcations are common in systems and synthetic biology, where cell-type switching dynamics often involve the emergence of bistability and bifurcations [27,28]. Cellular decisions in biological processes usually require several SNs coexisting within the same system, making it an ideal case study for our research [29,30]. For instance, during embryonic development, cellular subtypes arise from the cross-repression of genes expressed in each cell type, which together form gene regulatory networks (GRNs). Under spatial signaling, these networks encode the robust spatiotemporal orchestration of cell-type specification in the embryo [31–33]. Nevertheless, there is a lack of systematic understanding of the role that different close bifurcations may have in the resulting timing of cell transitions. Therefore, gaining insights into the timing properties of critical phenomena near multiple bifurcations is essential for fully comprehending the temporal dynamics that govern cell decision-making processes.

Slowing down in GRNs has been achieved experimentally in synthetic bistable switches in bacteria [34]. This has opened the door to the proposal of synthetic circuits that perform specific dynamical functions with applications in biomedicine, such as biosensors, customized drug delivery, or the design of responsive materials [35,36]. Some of these gene regulatory circuits are capable of undergoing several bifurcations while keeping a minimum set of molecular components. This is the case of the AC-DC genetic circuit that exhibits bistability between a stable steady state and oscillations [37]. This multifunctional behavior is paramount in synthetic biology where dynamical instructions need to be encoded with a minimal set of components to avoid the metabolic overload of the cell.

A multifunctional behavior of special interest for synthetic biology applications arises in so-called mushroom bifurcation

diagrams, named after its mushroom shape. These diagrams were initially observed in models for neural stem cell differentiation [38,39], and later in heat-shock protein dynamics in anxiety disorders [40]. The mushroom bifurcation diagram is formed by combining two toggle switches, resulting in four SN bifurcations and three disconnected stable branches [see Fig. 1(b)]. The presence of a mushroom-shaped locus of equilibria provides the system with unique hysteresis properties resulting from two different regions of bistability. Furthermore, it supplies the system with emergent behaviors such as transition-to-isola states [41], resettable memory, or detection of transient signals [42]. Some of these properties can be related to the potential for isola bifurcation diagrams [43], which have not yet been confirmed experimentally.

Due to this interesting phenomenology, mushroom bifurcations have received special attention lately, mostly in studies trying to dissect the mechanistic rules under which the mushroom bifurcation can emerge in genetic circuits [42,44–47]. While these studies have focused on the specific shape of the bifurcation diagrams, they only address the steady-state properties of the circuit, without exploring its dynamics. In particular, the possibility of controlling the proximity of two SN bifurcations and exploiting the properties of emerging transients.

To acquire a suitable understanding of the dynamics close to bifurcations in GRNs it is important to consider the effect of noise. Noise may occur intrinsically due to the random nature of biochemical reactions of gene expression, or extrinsically through environmental factors such as noise in cell signaling or cell-cell variability [48,49]. Intrinsic noise sources occur at different stages of the process of protein expression such as stochasticity in promoter binding, mRNA translation, or protein degradation, leading to significant molecular heterogeneity [50–52]. Despite its prevalence, molecular noise is usually studied in the context of steady-state dynamics and rarely in the context of transient dynamics. Therefore, further study of the noise on the transient dynamics close to bifurcations is required to fully elucidate the impact of stochastic gene expression on cellular decision-making processes.

Here, we address these questions focusing on the mushroom bifurcation diagram found in a minimal gene regulation circuit. In Sec. II, we introduce the mathematical model for the mushroom bifurcation diagram, which models a self-activating gene that is regulated nonmonotonically by an external signal. In Sec. III we explore the resulting deterministic dynamics of the mushroom and its associated normal form. We present analytical and numerical results for the scaling laws, where the synergy of two SN bifurcations close to a transcritical bifurcation is introduced. Finally, in Sec. IV, the effect of intrinsic and extrinsic noise in transient times is investigated for coupled bifurcations and compared to single SN bifurcations. Section V is devoted to conclusions.

II. DETERMINISTIC ONE-DIMENSIONAL MUSHROOM MODEL

To investigate the properties of the mushroom bifurcation diagram, we propose a minimal nonlinear one-dimensional system that exhibits this behavior, i.e. a model of a self-activating gene, u , which is controlled by an external signal,




FIG. 1. (a) Schematic diagram of the minimal gene regulatory system studied. A gene (u) with both constitutive and self-activated expression, where the strength of the self-activation of protein u is regulated by an external signal (s). The effect of s is nonmonotonic, resulting in maximum autoactivation at intermediate values of s . (b) The resulting bifurcation diagram shows the steady-state u^* as a function of the external signal s , corresponding to Eq. (1). The choice of parameters controls the shape of the bifurcation diagram. For high enough values of q the system shows the mushroom bifurcation diagram (left panel with $r = 0.14$ and $q = 3.0$). The neck of the mushroom becomes narrower as q is reduced to $q \rightarrow q_c$, resulting in the collapse of SN1 and SN4 towards a transcritical (T) bifurcation (right panel $r = 0.14$ and $q = q_c \simeq 2.78$). The left panel shows the four saddle-node (SN) bifurcations characteristic of the mushroom bifurcation containing two zones of bistability (shaded blue areas with flow direction indicated with black arrows). The slowing down occurs close to the bifurcations (blue arrows).

s , [Fig. 1(a)]. This external signal has an activating non-monotonic effect, inducing maximum autoactivation of u at intermediate signals, s . In the deterministic limit, this system is described by the ordinary differential equation (ODE)

$$\frac{du}{dt} = r + a(s)H(u) - u, \quad (1)$$

where u has a constant basal/leaky expression, r , and a linear decay (with a rate set to 1 without loss of generality). The self-activation term, $H(u)$, given by the positive Hill function

$$H(u) = \frac{u^2}{1 + u^2},$$

is regulated by the external function $a(s) = sq - s^2$. To ensure bistability r must satisfy

$$r \in \left(0, \frac{1}{3\sqrt{3}}\right).$$

For $r > 1/(3\sqrt{3})$, the system is monostable, rendering it unsuitable for our study. The parameter q controls the shape of the signal function $a(s)$, which remains positive [$a(s) > 0$] in the interval $[0, q]$. The quadratic shape of $a(s)$ captures the incoherent nature of the signal, which induces a maximum activation of u at an intermediate signal $s = q/2$, and fully represses u self-activation at the extreme values $a(0) = a(q) = 0$ [see Fig. 1(a)]. The shape of the resulting bifurcation diagram can be controlled with parameter q . At high values of q the system exhibits a mushroom bifurcation diagram with two zones of bistability [Fig. 1(b), left]. As q is reduced to a critical value q_c , two of the SN bifurcations collide, undergoing a transcritical bifurcation. After this collision, the mushroom

shape transforms into an isola, which becomes disconnected from the lower stable branch [Figs. 2(a)–2(d)].

III. DETERMINISTIC SLOWING DOWN IN MUSHROOM BIFURCATION DIAGRAMS

In this section, we investigate the deterministic dynamics arising in the mushroom bifurcation diagram, focusing on the case where the lower SN bifurcations, SN1 and SN4, approach each other [Fig. 1(b) and Figs. 2(a)–2(c)]. To do so, we first introduce a normal form for two SN bifurcations able to collapse into a transcritical one. This model will be useful to obtain analytical results that will allow us to analyze the behavior of the full genetic model described by Eq. (1). The numerical results for this section have been obtained using a seventh- to eighth- order Runge-Kutta-Fehlberg-Simó method with automatic time step size control and local error tolerance of 10^{-15} .

A. Symmetric collision of two saddle nodes into a transcritical bifurcation model: a normal form

To describe a normal form for a dynamical system exhibiting two symmetric SN bifurcations converging into a transcritical bifurcation (see Fig. 3), we analyze the degenerate fixed point at the origin for $r = 0$. Introducing the unfolding parameter r , the system is described as an unfolding of the transcritical bifurcation (see Appendix C for details on the derivation):

$$\frac{dx}{dt} = x^2 - s^2 + r. \quad (2)$$

For $r < 0$, no bifurcation occurs. Instead, as r approaches 0, we observe a family of branches of fixed points that approach and collide with the degenerate fixed point at the origin




FIG. 2. Evolution of the mushroom bifurcation diagram for Eq. (1) at decreasing q with $r = 0.14$. Panels (a) $q = 3$ and (b) $q = 2.786$ show the diagram with four SN bifurcations. (c) Transcritical bifurcation at $q = q_c = 2.784953 \dots$. Two examples of isolas are shown with $q = 2.7$ (d) and $q = 2.605$ (e). Panel (f) shows how the isola collapses at $q = 2.600914$. Bistability regions are shown in blue.

[see Fig. 3(b)]. In the context of the mushroom bifurcation, this would correspond to the isola scenario, illustrated in Figs. 2(d) and 2(e). When r crosses zero ($r = 0$), the transcritical bifurcation emerges [Fig. 2(c)]. As r increases ($r > 0$), the unfolding of the transcritical bifurcation captures the behavior of interest, as shown in Fig. 3(a), where for two symmetrical SN bifurcations form and diverge from the transcritical bifurcation.

Differentiating Eq. (2) reveals that the equilibria of the system are stable when they are located at $x < 0$, and unstable for $x > 0$. In particular, for the case $r = 0$, Eq. (2) simplifies to $\dot{x} = x^2 - s^2$, containing the steady-state branches $x^* = s$ and $x^* = -s$. Thus, both branches exchange stability at the origin, indicating that the system undergoes a transcritical bifurcation.

To study the role that both SN bifurcations have on the transient times, we will consider the transition from $x = -\epsilon$ to $x = \epsilon$ for a fixed value of $r > 0$, ϵ being an arbitrarily small real value. The scaling law of the SN bifurcation of the normal form model [13,14] is examined by fixing a value of $s = A\sqrt{r}$, leading to an analysis of the extended transients close to each SN point, i.e., $s_{sn}^- = -\sqrt{r}$ (left SN point) and $s_{sn}^+ = +\sqrt{r}$ (right SN point). Here, A ranges from -1 to 1 resulting in values of s that span the full range of values between both SN bifurcations. The duration τ of the transients as the parameter $s \rightarrow s_{sn}^\pm$ (see Fig. 3), is computed using $s^2 = A^2r$, from the integral

$$\tau = \int_{-\epsilon}^{\epsilon} \frac{1}{r - s^2 + x^2} dx \stackrel{s=A\sqrt{r}}{=} \int_{-\epsilon}^{\epsilon} \frac{1}{r - A^2r + x^2} dx.$$

Obtaining the asymptotic behavior of $\tau(r, A)$ in the vicinity of the SN bifurcations as $(1 - A^2) \rightarrow 0$, where $r \rightarrow 0^+$ indicates the proximity of both SNs.

$$\tau(r, A) \stackrel{(1-A^2) \rightarrow 0}{\sim} \frac{\pi}{\sqrt{1 - A^2} \sqrt{r}}. \quad (3)$$

Notably, in contrast to the normal form for a single SN bifurcation [53], notice that the times in Eq. (3) have a prefactor dependent on r , specifically $\frac{\pi}{\sqrt{r}}$, which indicates that as the two SN bifurcations converge, i.e., $r \rightarrow 0^+$, the slowing down is significantly amplified, thereby lengthening the transient times τ while keeping the inverse square root law.

To assess the slowing down close to the transcritical bifurcation, we study a trajectory along the symmetry axis $s = 0$ or equivalently, the neck, between the two SN bifurcations, while progressively approaching the SNs to their convergence, equivalently, as $r \rightarrow 0^+$, as illustrated in Fig. 3(a). The integral representing the timescale of transient times $\tau(r, s)$ is expressed as follows:

$$\tau(r, s) = \int_{-\epsilon}^{\epsilon} \frac{1}{r - s^2 + x^2} dx \stackrel{s^2 \leq r}{=} \frac{2}{\sqrt{r - s^2}} \arctan \left(\frac{\epsilon}{\sqrt{r - s^2}} \right). \quad (4)$$

Finally, we derive the asymptotic behavior of $\tau(r, s)$ near the transcritical bifurcation as $r \rightarrow 0^+$. In this limit, the transcritical bifurcation represents an asymptotic state that is never fully reached, with the “neck” being infinitesimally opened. This allows us to examine the timescales in the vicinity of




FIG. 3. (a) Bifurcation diagram obtained from the normal form, Eq. (2). The solid and dashed lines correspond to stable and unstable steady-state branches, respectively. (a) As r decreases (for $r > 0$) two saddle nodes (SN, circles) at a distance $\Delta s = 2\sqrt{r}$ move towards each other colliding into a transcritical bifurcation at $r = 0$ (diagonal lines). For all values of $r > 0$ the system can evolve from $x < 0$ to $x > 0$ (purple arrow) with a transient time that will depend on the location of the SN bifurcations. (b) For $r < 0$ no bifurcations are found.

$x = \epsilon$ and $x = -\epsilon$, obtaining

$$\begin{aligned}
 \tau(r, s) &= \lim_{r \rightarrow 0^+} \frac{2}{\sqrt{r - s^2}} \arctan\left(\frac{\epsilon}{\sqrt{r - s^2}}\right) \\
 &= \lim_{r \rightarrow 0^+} \frac{\pi}{\sqrt{(\sqrt{r} - s)(\sqrt{r} + s)}} \\
 &= \lim_{(s_{sn}^- - s_{sn}^+) \rightarrow 0^+} \frac{\pi}{(s_{sn}^- - s_{sn}^+)} \\
 &\sim \frac{\pi}{(s_{sn}^- - s_{sn}^+)}.
 \end{aligned} \tag{5}$$

Note that we recover the scaling law for the transcritical bifurcation: $\tau \sim (s_{sn}^- - s_{sn}^+)^{-1}$. It is also noteworthy that one could follow a similar strategy to study the transients at the

formation of the isolas [see Appendix A, Figs. 2(d) and 2(e)] [42,44]. To analyze this case, one should use the normal form:

$$\dot{x} = x^2 + s^2 + r, \tag{6}$$

where the value of r controls the appearance of the isolas (for more details on a criterion to identify isolas and transcritical bifurcations, see Appendix A).

B. Gene regulatory system

We now study the dynamics of the minimal genetic system introduced in Eq. (1) [see Fig. 1(a)], regarding transients and their scaling properties.

Without loss of generality, we will focus on transient times close to SN1. To study how delays are affected by nearby bifurcations, we will study the change in transient times as SN1 and SN4 approach each other. To do so, we introduce the quantities $\Delta s \equiv s - s_{SN1}$ and $\Delta_{4,1} \equiv s_{SN4} - s_{SN1}$, where s_{SN1} and s_{SN4} denote the values of the signal s at which the bifurcations SN1 and SN4 occur, respectively [see Fig. 4(a)]. The magnitude Δs allows us to analyze the effect of a given signal relative to the value at which SN1 occurs. This is the basis for examining the scaling law in a single SN bifurcation and understanding how other bifurcations affect the transient times. On the other hand, $\Delta_{4,1}$ controls the proximity of SN1–SN4, allowing us to study synergistic effects between both SNs and the transcritical bifurcation occurring when $\Delta_{4,1}$ approaches 0. As previously mentioned, by synergistic we mean that the transient times arising from the coupling of these bifurcations is longer than their independent addition. As $\Delta_{4,1} \rightarrow 0^+$ we can derive the power law with exponent -1 , associated with the transcritical bifurcation. This behavior is affected by SN1 and SN4, by introducing prefactors that extend their respective timescales (see Appendix B). The neck width $\Delta_{4,1}$ can be controlled by keeping r constant and decreasing parameter q , thereby changing the signaling effect on the autoactivation.

As expected, when both SN bifurcations are far from each other ($\Delta_{4,1} > 1$), numerical analysis of the transient time shows the well-known inverse square-root power-law scaling for transient times ($\tau \sim \Delta s^{-1/2}$) [see Fig. 4(b)]. That is, just after the bifurcation, the orbits experience delays and long plateaus associated with the ghost [Fig. 4(c)] [9,13,14,31]. For an analytical study of the SN bifurcation scaling law in the self-activating gene model, see Appendix C.

We posit that the ghosts corresponding to SN1 and SN4, which are influenced by the slowing down associated with the transcritical bifurcation, will have a nonlinear influence on each other, so that transient times will increase synergistically. To validate this hypothesis, we examined the slowing down at signal values close to SN1 ($\Delta s \ll 1$) for scenarios in which the neck of the mushroom is very narrow [$\Delta_{4,1} \ll 1$; see Fig. 5(a)]. The resulting slowing down is much longer than what we would expect from a simple additive effect of two SNs. This synergistic amplification can be seen in Figs. 5(b) and 5(c) where transients are displayed for different mushroom necks ($\Delta_{4,1}$). By comparing the resulting transients with the expected values from simply adding the transient times of two independent SNs [blue line in Figs. 5(b) and 5(c), and 6)], we observed that the resulting transients can be orders of magnitude longer. In Appendix B we compare the delays for




FIG. 4. (a) Mushroom bifurcation diagram showing the steady-state u^* as a function of the signal s for the self-activating gene model given by Eq. (1) when the lower saddle nodes SN_1 and SN_4 are far from each other ($q = 5$, $r = 0.14$) corresponding to a mushroom neck distance $\Delta_{4,1} \simeq 4$. (b) Transient times τ as a function of the distance signal Δs across the slowing-down region close to SN_1 [corresponding to the shaded zone in panel (a)]. Transient times are calculated computationally as the time τ required to reach $u(\tau) = 0.4$ from an initial state $u(0) = 0.2$ (SN_1 is located at $u^* = 0.31$). (c) Characteristic bottleneck of ghost transients for $\log_{10}(\Delta s) = -5.6$ [blue circle in panel (b)].




FIG. 5. (a) Mushroom bifurcation diagram computed from Eq. (1) with a very narrow neck using $\log_{10}(\Delta_{4,1}) \approx -6.62$. (b) Transient times as a function of Δs for different neck sizes (from bottom to top) $\log_{10}(\Delta_{4,1}) \approx \{0.62, -0.54, -1.63, -2.63, -3.63, -4.63, -5.63, -6.34, -6.62\}$; other parameters and conditions are the same as in Fig. 4. The results show a $-1/2$ scaling law with transient times longer than what would be expected by the additive effect of two independent SNs (blue line, obtained by doubling the time expected by a single SN in the case of a wide-neck mushroom). Inset: the same scaling law is obtained for the transient times as a function of neck width for a fixed value of $\log_{10}(\Delta s) = -7.5$ (vertical green line and dots). (c) Time series for the same neck sizes and conditions as in panel (b) for a fixed signal value $\log_{10}(\Delta s) = -6.7$, corresponding to the vertical blue line in panel (b). (d) Transient times keeping $\Delta s = \Delta_{4,1}/2$ shown in log-linear scales. Three examples are displayed: (i) $q = 5$, (ii) $q = 2.786$, and (iii) $q = q_c + 10^{-14}$, corresponding to $\log_{10}(\Delta_{4,1}) = 0.62$, $\log_{10}(\Delta_{4,1}) = -1.12$, and $\log_{10}(\Delta_{4,1}) = -6.62$, respectively. The color maps indicate the duration of transients in the gap between SN_1 and SN_4 , which is computed by dividing this gap into 10^2 equal intervals.




FIG. 6. Transient times τ close to the bifurcation value (in log-log scale) for six cases given by the normal form (NF) of the SN bifurcation (blue triangles); twice the times for the NF of the SN (violet triangle); transcritical NF (open blue circles); sum of the times for the transcritical NF plus twice the times of the SN NF (solid green circles); NF of the coupled SN bifurcations collapsing into the transcritical one [Eq. (2)] (orange circles); and the mushroom model given by Eq. (1) (black circles). The inset compares these times (in log-linear scales) for the mushroom NF (orange) and the times assuming additivity of two independent SNs with the transcritical slowing down using their NFs (green).

single local bifurcations and consider additivity in transient times with the multiple bifurcations described above.

Furthermore, the smaller the gap (smaller values of $\Delta_{4,1}$), the slower the transition, deviating further from additivity. This reveals a synergistic amplification of transients. On the other hand, the scaling law $\tau \sim \Delta s^{-1/2}$ is preserved, suggesting that the slower transient results in an increase in the prefactor of the scaling law that cannot be explained merely by the appearance of the transcritical bifurcation but requires the proximity of both SNs.

In an attempt to clarify the behavior of the prefactor and better understand the synergistic effects, we analyzed the prefactor in the scaling law. We did so by studying the increase of the transients as the width of the neck varies for a fixed value of Δs [green dots and inset in Fig. 5(b)]. This analysis showed that the prefactor also follows an inverse square-root power-law behavior. Such behavior is consistent with the results of the normal form [Eq. (2)] found in the previous section. Furthermore, it can be derived analytically for the mushroom model (see Appendix C).

To further study the effect of the coupling between both SN bifurcations as they approach each other, we studied the slowing down for signals exactly at the middle point between both saddle nodes $\Delta s = \Delta_{4,1}/2$ for different values

of $\Delta_{4,1}$ [Fig. 5(d)]. Similar results were found, where for a narrower mushroom neck [$\log_{10}(\Delta_{4,1}) \approx -6.6$], transient times are seven orders of magnitude slower compared to a wider neck. This observation reiterates the significant influence of the interaction between the two collapsing SN bifurcations. Figure 5(d) shows (in log-linear scale) how the delays increase as the width of the mushroom neck closes. This curve follows the scaling behavior with a power exponent -1 associated with the transcritical bifurcation (see the case with black data in Fig. 6 for the same results in the log-log scale). This indicates that the slowing down effect on the system is influenced not only by the two ghosts but also by the transcritical bifurcation. An alternative approach to derive this result is to express τ as a coupling of the SN slowing down as a function of a in Eq. (1), along with the decreasing slope of $a(s)$. Specifically, we can write $\tau \sim [a(s) - a_c(s)]^{-1/2} \sim (s - s_c)^{-1}$, using the fact that near the transcritical bifurcation, $a'(s) = 0$ and $a''(s) < 0$.

IV. STOCHASTIC DYNAMICS IN MUSHROOM BIFURCATIONS

In order to investigate the prevalence of these transients in realistic scenarios, it is crucial to examine the effects of noise on the observed dynamics. To achieve this, we derive molecular reactions compatible with the mushroom gene regulatory network introduced in Sec. II [Fig. 1(a) and Eq. (1)]. We begin by analyzing the effects of intrinsic (demographic) and extrinsic (environmental) noise separately, before exploring the system with both sources of noise combined.

A. Intrinsic noise

Intrinsic noise is studied by explicitly implementing the reactions of protein production and degradation controlling the dynamics of the absolute number of proteins, $U = \Omega u$, where the system volume, Ω , captures the relevance of the fluctuations. The rate functions in Table I describe the stochastic model. These rates result in a birth-death process of protein production and degradation with propensities $W_1(U)$ and $W_2(U)$, respectively. Stochastic trajectories for this process were generated using the Gillespie algorithm [54]. The deterministic model can be recovered in the limit of a large number of molecules ($\Omega \rightarrow \infty$) (see Ref. [51] for details). In this limit, the change in U over time is described by an ODE determined by the propensities of both reactions

$$\frac{\dot{U}}{\Omega} = \frac{W_1(U)}{\Omega} - \frac{W_2(U)}{\Omega},$$

resulting in the deterministic model [Eq. (1)].

Intrinsic noise results in fluctuations of u , which can induce switching in the bistable zones of the mushroom [$s \in (s_{SN_2}, s_{SN_1})$ and $s \in (s_{SN_4}, s_{SN_3})$; see blue areas in Fig. 1(b)]. Thus, intrinsic noise leads to an expansion of the signal values over which gene expression levels can transition to the upper branch of the mushroom (starting with the initial condition $u = 0$). This noise-induced tipping is observed in any genetic bistable switch [31,55]. In our particular case, this implies that our study of transient times needs to expand to signals beyond

TABLE I. Intrinsic noise model. Protein U is produced and degraded with the propensities W_1 and W_2 , respectively.

Elementary process	Propensities	Stoichiometry	Description
$\emptyset \rightarrow U$	$W_1 = \Omega \left(r + \frac{a(s)U^2}{(\Omega^2 + U^2)} \right)$	+1	regulated protein production
$U \rightarrow \emptyset$	$W_2 = U$	-1	linear protein degradation

the regions studied in the deterministic scenario. Figure 7(a) shows the transition times calculated as first passage times towards the upper branch of the mushroom. For all values of signal and noise intensities, intrinsic noise accelerates the transient with respect to the deterministic scenario. This effect is more dramatic close to the SN bifurcations, where the deterministic times diverge while there is still a finite transient time for the stochastic case. For larger noise intensities (lower Ω) the transients are shorter. This leads to an almost constant transition time along the whole mushroom neck for high intrinsic noise ($\Omega = 10$). This behavior is in line with the fact that intrinsic noise allows the system to overcome the ghost, escaping the regime for which $\dot{u} \simeq 0$. Studying the coefficient of variation (the standard deviation of the mean first passage time divided by its mean), we observe that the precision of the distributions decreases with noise [Fig. 7(b)]. This suggests that intrinsic noise can be detrimental to encoding precise transient times.

As in the deterministic case, constricting the neck of the mushroom results in longer transients that are not erased by intrinsic noise, even in scenarios with high intrinsic noise [Fig. 7(c)]. Similarly, the trend of decreasing precision of transient times with noise intensity is also preserved.

B. Extrinsic noise

We can incorporate the effect of extrinsic noise as stochastic fluctuations around a given input signal, s_0 . To do so we describe the stochastic dynamics of signal, s , as an unregulated process with a constant production rate, R , and linear degradation rate, R/s_0 . Defining the total number of signaling molecules as $S = s\Psi$, the signal fluctuations result from

another birth-death process where the volume parameter Ψ controls the intensity of the extrinsic noise (see Table II). Stochastic trajectories for this process were also generated using the Gillespie algorithm. The deterministic limit for the extrinsic noise is recovered in the limit $\Psi \rightarrow \infty$ returning the ODE

$$\frac{\dot{S}}{\Psi} = \frac{V_1}{\Psi} - \frac{V_2}{\Psi} = R \left(1 - \frac{s}{s_0} \right),$$

which relaxes to the deterministic steady state $s = s_0$.

Interestingly, for signal regimes between the SNs, extrinsic noise significantly slows down the transient compared to the deterministic scenario [see Fig. 8(a)], in opposition to what was observed for intrinsic noise. This stochastic slowing down increases with increasing extrinsic noise intensity (lower values of Ψ). However, as in the intrinsic noise scenario, extrinsic noise also expands the signal region over which gene expression can transit toward the upper branch of the mushroom. Therefore, we find that for both intrinsic and extrinsic noise, the expanded signal regime for the transition is due to noise-induced tipping allowing the system to transition. Similarly, close to the SNs where the deterministic transient time diverges, extrinsic noise accelerates the transient. Analysis of the distribution of transition times under extrinsic noise shows a similar effect to intrinsic noise in which higher noise is translated into a higher coefficient of variation. Nevertheless, for a large range of noise intensities the coefficient of variation (CV) remains very low (below 0.1), suggesting that extrinsic noise does not have a high impact on the variability of the transients [see Fig. 8(b)]. High coefficients of variation (CV > 0.5) are only observed in limit cases with narrow mushroom necks and high extrinsic noise [see Fig. 8(c)].




FIG. 7. Transition times, τ , computed as the first passage times towards the upper branch of the mushroom $u_r = 1.4$, starting with initial condition $u_0 = 0$. Deterministic transients (black solid line) are compared with stochastic transients for different noise intensities: $\Omega = 10$ (blue), $\Omega = 10^2$ (red), and $\Omega = 10^3$ (yellow). Parameters are $r = 0.14$, $q = 3$, resulting in a mushroom neck $\Delta_{4,1} = 1.15$. (a) Transient times as a function of the signal. Positions of saddle nodes SN1 and SN4 ($\Delta s = 0$ and $\Delta_{4,1} = 1.1$), are indicated by dotted lines. (b) Histogram showing the probability distributions of the transition times for a fixed value of signal $\Delta s = \Delta_{4,1}/2$. (c) Histogram of transition times for a smaller mushroom neck $\Delta_{4,1} = 0.6$ ($q = 2.85$) at a signal $\Delta s = \Delta_{4,1}/2$. Simulations are performed using the Gillespie algorithm [54] using the reaction schemes described in Table I.

TABLE II. Extrinsic noise reactions. Both reactions with propensities V_1 and V_2 introduce fluctuations around the state $S = \Psi s_0$. Simulations along this article use $R = 1$.

Elementary Process	Propensities	Stoichiometry	Description
$\emptyset \rightarrow S$	$V_1 = \Psi R$	+1	constant signal production
$S \rightarrow \emptyset$	$V_2 = RS/s_0$	-1	linear signal degradation

C. Full stochastic model

Given the opposite effects of intrinsic and extrinsic noise within the neck of the mushroom $\Delta_{4,1}$, we sought to investigate the behavior of the system when both sources of noise are introduced simultaneously, a situation expected to happen in a real experimental setup. To do so, we analyzed the transient times for different combinations of intrinsic and extrinsic noise (Fig. 9). The results show that mean transient times can be faster or slower than the deterministic time, depending on the combination of intrinsic and extrinsic noise intensities. Notably, one can keep the same timing as the deterministic scenario by balancing both sources of noise over a wide range of values of noise intensities [Fig. 9(b)]. However, this relationship is different for the coefficient of variation of the timings, where the precision is primarily controlled by the intrinsic noise intensity [Fig. 9(c)]. This suggests a mechanism by which one can achieve a specific transient timing with a prescribed precision by balancing the intrinsic and extrinsic noise intensities.

Some of the timing dynamics for the noisy system for the mushroom bifurcation can be attributed to the existence of SN bifurcations independently of the mushroom bifurcation diagram. To understand the coupling effect of the SNs characteristic of the mushroom we compared the transient average time and precision with the case in which a single SN is present (Fig. 9). Specifically, we studied a bifurcation diagram corresponding to a traditional bistable switch built by keeping the mushroom bifurcation diagram up to its symmetry point $\Delta s = \Delta_{4,1}/2$ and fixing a constant steady state for higher values of the signal (see Appendix D for details). Before considering the case with mixed noises we can study the effect that intrinsic and extrinsic noise have independently on the transient times of the bistable switch. As expected, intrinsic noise transient is unaffected for values of $\Delta s < \Delta_{4,1}/2$, where

the transients of the mushroom and bistable switch are identical [Fig. 10(a)]. On the other hand, intrinsic noise transients are independent of the signal for $\Delta s > \Delta_{4,1}/2$ [Fig. 10(a)]. These results confirm that intrinsic noise effects only depend on the steady states available at a particular value Δs , independently of the bifurcation diagram. This is not true for extrinsic noise, where mushroom and bistable switch transients are different for all values of Δs , including values $\Delta s < \Delta_{4,1}/2$ for which both bifurcation diagrams are identical [Fig. 10(b)]. This is the result of extrinsic noise allowing the system to sample different signal values, and thereby be affected by localized slowing down effects near the SNs. Since the effect of the SNs is to slow the transients, extrinsic noise slows both the bistable switch and the mushroom bifurcation. This effect is stronger for the mushroom where two SNs contribute to the slowing down dynamics [Fig. 10(b)]. As the width of the neck increases, it is expected that the difference in transition times between both diagrams will decrease, as a signal will have a lower probability of attaining values close to the SNs.

Next, we aim to understand whether balancing intrinsic and extrinsic noise to control the mean transient time and precision is a specific property of the mushroom configuration as shown above, or if it can also be achieved in the bistable switch. Comparison of the transition times for different noise intensity combinations [Figs. 9(d)–9(f)] show that at $\Delta_{4,1}/2$, the transient times in the bistable switch diverge from the deterministic case with a similar trend to the one observed in the mushroom. However, the change in transition time is much larger for the mushroom bifurcation. In particular, the mushroom allows for a more dramatic slowing down of the transition, achieving transient timescales that can be more than double the deterministic time. Meanwhile, when comparing the coefficient of variation between the mushroom and the bistable switch, we find that the coefficient of variation does




FIG. 8. Transition times, τ , for the mushroom under the same parameters and conditions as in Fig. 7. Deterministic transients (black solid line) are compared with stochastic transients for different extrinsic noise intensities: $\Psi = 10$ (blue), $\Psi = 20$ (cyan), $\Psi = 50$ (red), and $\Psi = 100$ (yellow). (a) Transition times at different Δs . Panels (b) and (c) display histograms with probability distributions of transient times sizes $\Delta_{4,1} = 1.15$ and $\Delta_{4,1} = 0.6$, respectively. Simulations are performed using Gillespie algorithm [54] employing the reaction schemes described in Table II.




FIG. 9. Transition times and variability measures for mushroom bifurcations (a)–(c) and bistable switches (d)–(f), for $r = 0.14$ and $q = 3$ with initial expression $u_0 = 0$ to threshold $u_\tau = 1.4$ for variations in extrinsic stochastic noise, Ψ , and variations in intrinsic noise, Ω , at $\Delta_{4,1}/2 = 0.5575$. Note that for the deterministic system, the transition time is $\tau_{\text{det}} = 17.0439$. Panels (a) and (d) show schematic bifurcation diagrams for the mushroom system and the bistable switch, respectively. The dotted gray line at s_0 corresponds to $\Delta_{4,1}/2$, and the pink and blue arrows correspond to the effective direction of intrinsic and extrinsic noise, respectively, as controlled by Ω and Ψ . Panels (b) and (e) show the differences in mean transition times from deterministic time $\Delta\tau = \langle\tau\rangle - \tau_{\text{det}}$. Panels (c) and (f) display the coefficient of variation $\sigma/\langle\tau\rangle$. Simulations are performed using Gillespie algorithm [54] employing the reaction schemes described in Tables I and II.

not change significantly across the models [cf. Figs. 9(c) and 9(f)]. All in all this suggests that the mushroom bifurcation allows for a better control of the slowing down dynamics in stochastic scenarios.

V. CONCLUSIONS

Transients are known to become extremely long close to critical transitions in a phenomenon known as critical slowing down. This phenomenon has implications in physical and biological systems introducing dynamics much slower than the timescale of the rates composing the system [1,3–8,10,11]. The transient times, τ , near deterministic bifurcations typically follow universal scaling laws of the form $\tau \sim |\mu - \mu_c|^\beta$ [9,13]. Such exponents have been identified in both local [14–16] and global [24] bifurcations. Despite critical slowing down and transient scaling phenomena having been widely investigated over the past decades for single bifurcations (either local or global), the question of how dynamical systems behave close to nearby bifurcations remains unexplored.

In this article, we have studied critical slowing down in systems exhibiting mushroom bifurcation diagrams. This bifurcation diagram is usually found in mathematical models of gene regulation [42,44–47] and, for appropriate parameter values, it has four SN bifurcations. The simple model presented in this article considers a gene that self-activates and is regulated by an external signal. By tuning biochemical parameters, it is possible to arbitrarily control the distance of the two SN bifurcations forming the neck of the mushroom. By further reducing this distance, the neck of the mushroom eventually

closes until both SN bifurcations collapse into a transcritical one. For this system, we show that the slowing down found right after a SN bifurcation largely increases when the two SN bifurcations approach each other. Moreover, the inverse square-root scaling law remains preserved, as shown for SN bifurcations studied with delay differential equations [19]. We also provide analytical and numerical evidence of this synergistic effect by examining a normal form with the same phenomenology, i.e., two SN bifurcations merging into a transcritical one. In a nutshell, we have found that the critical slowing down obtained by the coupling of the two SN bifurcations with the transcritical one is much longer than the times expected from summing the transients of the transcritical and SN bifurcations alone, thus identifying a nonadditive effect arising from this coupling.

In the context of gene regulatory networks (GRNs), our findings provide a mechanism to control the timing of an event, achieving timescales slower than those of protein production and degradation. This suggests an approach to the construction of genetic timers. Traditionally, GRN transients have been overlooked in systems biology with a primary focus on steady states. However, the timing of transitions is crucial for understanding GRN functions. For example, differences in embryonic developmental timing between species often arise from the same reaction networks with nearly identical rates, highlighting the importance of understanding time control in developmental processes [56].

One of the defining features of GRNs is the significant impact of noise due to the low number of molecules involved. Our study demonstrates that specific bifurcation diagrams,




FIG. 10. Transient times at different signal values for the mushroom bifurcation (blue lines) and the bistable switch (red lines) with $r = 0.14$, $q = 2.85$, i.e., $\Delta_{4,1} = 0.6$. The symmetry axis of the mushroom where the bistable switch and mushroom start to differ is located at $\Delta s = \Delta_{4,1}/2 = 0.3$ (dotted gray line). Transients of the corresponding simulations in the deterministic case are also plotted for reference (dashed lines). (a) Comparison of deterministic and stochastic transition times for a system with intrinsic noise. (b) Comparison of deterministic and stochastic transition times for a system with extrinsic noise.

which can be achieved with minimal GRN topologies, not only exploit these transients but also buffer the effects of noise. Importantly, we show that intrinsic and extrinsic noise have opposite effects on timing. This discovery enables the achievement of specific timings with prescribed precision by controlling intrinsic noise (e.g., through plasmid number regulation in synthetic circuits) and extrinsic noise (e.g., by modulating signal noise).

While our study focused on the effects of coupled local bifurcations in critical slowing down due to bifurcation (B-tipping), and noise-induced tipping (N-tipping) [57], further mechanisms such as rate-induced tipping and overshooting [58–60] in nearby bifurcations offer an intriguing area for further study. As rate-induced tipping occurs when parameters change too quickly for a system to adapt to the changes, these dynamics may be of particular interest in determining the robustness of GRNs. This is especially the case in terms of overshooting tipping point thresholds [61], and in the

active control of GRN signaling. Concerning SN overshooting, recent research has explored the probability of tipping under parameter forcing, and provided an approximation of this probability in the case of the time-changing parameter reversing near the tipping point [62]. These authors observed that the critical curve relating the forcing in the deterministic system and the time for which the parameter forcing exceeded the fold bifurcation followed the inverse-square law. Similar investigations using random disturbances also converged to this law, provided that noise intensity was kept low [62]. Our results could imply that SN overshooting in close, local bifurcations may decrease the tipping probabilities and increase the times for which the forcing could surpass the fold bifurcation due to the synergies described here. One might also expect these critical curves to follow the scaling laws described here for the coupled bifurcations. Future research could investigate the overshooting phenomenon in nearby bifurcations.

Furthermore, while our study explores the one-dimensional case, it opens the door for future research to investigate scenarios involving multiple genes. Examining how different bifurcations interact could unveil new dynamical regimes not observed before, providing deeper insights into the complexity and robustness of GRNs in synthetic biology. These advancements not only enhance our understanding of biological timing mechanisms but also pave the way for innovative applications in the design of synthetic biological systems.

ACKNOWLEDGMENTS

We would like to thank Dr. Daniel O’Hanlon for his feedback on this work. We would also like to thank the anonymous referees for their comments. J.S. and T.A. have been supported by Grant No. PID2021-127896OB-I00 funded by MCIN/AEI/10.13039/501100011033 “ERDF A way of making Europe.” J.S. has also been supported by Ramón y Cajal Grant No. RYC-2017-22243 funded by MCIN/AEI/10.13039/501100011033 “ESF invests in your future.” R.P.-C. and S.M. acknowledge funding by the Leverhulme Trust Research Project Grant No. RPG-2023-085. A.H. has been supported by the Spanish Grant No. PID2021-125535NB-I00 (MCIU/AEI/FEDER, UE). This research has also been funded through the S. Ochoa and M. de Maeztu Program for Centers and Units of Excellence in R&D CEX2020-001084-M (A.H., T.A., and J.S.). We thank CERCA Programme/Generalitat de Catalunya for institutional support.

APPENDIX A: ISOLA AND TRANSCRITICAL BIFURCATIONS FROM MORSE THEORY

Consider the case where $q < q_c$, where q_c denotes the value of q at which the transcritical bifurcation occurs [Fig. 1(b), right panel]. For smaller values of q , an isola [41] appears. Equivalently, for Eq. (6), when considering $r < 0$, the isola appears and for $r = 0$, we locally observe a point in the bifurcation diagram, showing its collapse (see Fig. 2). The existence of this isola in the deterministic mushroom bifurcation diagram, together with the existence of the transcritical bifurcation, can be analytically demonstrated through the application of the Morse Lemma [63]:

Theorem A1 (Morse Lemma). Let $f : U \rightarrow \mathbb{R}$ be a C^3 function defined on an open set $U \subset \mathbb{R}^n$. Suppose $p \in U$ is a nondegenerate critical point (equilibrium point) of f with index s . Then, there exists a C^1 diffeomorphic immersion $G : V \rightarrow U$, where $V \subset \mathbb{R}^n$ is an open set, such that $g = f \circ G : V \rightarrow \mathbb{R}$ is of the form

$$g(x) = f(p) - x_1^2 - \cdots - x_s^2 + x_{s+1}^2 + \cdots + x_n^2.$$

From the Morse Lemma, the following corollary can be derived, which provides a criterion for distinguishing between the formation of isolas and transcritical bifurcations based on the sign of the determinant of the Hessian matrix at the critical point.

Corollary A1. (Isola formation). Let $f(x, \lambda)$ be a smooth function, such that

$$f(0, 0) = 0, \quad \frac{\partial f}{\partial x}(0, 0) = 0, \quad \frac{\partial f}{\partial \lambda}(0, 0) = 0, \\ \Delta = \frac{\partial^2 f}{\partial x^2}(0, 0) \frac{\partial^2 f}{\partial \lambda^2}(0, 0) - \left(\frac{\partial^2 f}{\partial x \partial \lambda}(0, 0) \right)^2 > 0,$$

where Δ denotes the determinant of the Hessian matrix of $f(x, \lambda)$.

Then, the following statements hold:

(1) If $\frac{\partial^2 f}{\partial x^2}(0, 0) > 0$, the point $(0, 0)$ is a local minimum, and the function $f(x, \lambda)$ can be expressed in the form

$$f(x, \lambda) = y_1^2(x, \lambda) + y_2^2(x, \lambda),$$

for certain smooth functions $y_1(x, \lambda)$ and $y_2(x, \lambda)$ defined in a neighborhood of $(0, 0)$.

(2) If $\frac{\partial^2 f}{\partial x^2}(0, 0) < 0$, the point $(0, 0)$ is a local maximum, and the function $f(x, \lambda)$ can be expressed in the form

$$f(x, \lambda) = -y_1^2(x, \lambda) - y_2^2(x, \lambda),$$

for certain smooth functions $y_1(x, \lambda)$ and $y_2(x, \lambda)$ defined in a neighborhood of $(0, 0)$.

Proof. By hypothesis, the Hessian matrix H of f at the origin is positive, therefore H is nondegenerate and $f = f(x, \lambda)$ has a nondegenerate critical point. By applying the Morse Lemma, we can assert that there exists a local diffeomorphism

$$h : (0, 0) \in J_1 \times J_2 \longrightarrow I_0 \times \Lambda_0 \ni (x_0, \lambda_0) \\ (j_1, j_2) \mapsto (x, \lambda),$$

such that $h(0, 0) = (x_0, \lambda_0)$ and $(f \circ h)(j_1, j_2) = j_1^2 + j_2^2$, where we assume, without loss of generality, that $\frac{\partial^2 f}{\partial x^2}(0, 0) > 0$. In case it was negative, we define $f \circ h(j_1, j_2) = -j_1^2 - j_2^2$. The bifurcation diagram for $f(x, \lambda)$ is reduced locally to a point.

The set of points satisfying the equation $f \circ h(j_1, j_2) = 0$ forms a set of isolas or closed curves in the j_1 - j_2 plane. Since the transformation h is a local diffeomorphism, these level curves, which are circumferences in variables j_1, j_2 defined by $f(x, \lambda) + c = 0$, map to a set of isolas in the x - λ plane.

The equation $(f \circ h)(j_1, j_2) = 0$ implies $j_1^2 + j_2^2 = 0$, which corresponds to the point $(0, 0)$. Now, if we look at a small neighborhood around $(0, 0)$, the equation $(f \circ h)(j_1, j_2) = c$ for small $c > 0$ describes a set of points forming a circle with radius $\sqrt{c}$ around the origin. ■

When these circles are transformed by h into the x - λ plane, they form isolas in a small neighborhood around the critical point (x_0, λ_0) . This is possible because the transformation h preserves the topology of the small neighborhood around the origin, and thus the isolas in the j_1 - j_2 plane for a fixed c parameter value, become isolas in the x - λ plane for this corresponding c value.

Corollary A2. (Transcritical bifurcation). Let $f(x, \lambda)$ be a smooth function such that $f(0, 0) = 0$, $\frac{\partial f}{\partial x}(0, 0) = 0$, $\frac{\partial f}{\partial \lambda}(0, 0) = 0$, $\frac{\partial^2 f}{\partial x^2}(0, 0) \neq 0$, and

$$\Delta = \frac{\partial^2 f}{\partial x^2}(0, 0) \frac{\partial^2 f}{\partial \lambda^2}(0, 0) - \left(\frac{\partial^2 f}{\partial x \partial \lambda}(0, 0) \right)^2 < 0,$$

where Δ denotes the determinant of the Hessian matrix of $f(x, \lambda)$ at $(0, 0)$.

Then, the local behavior of f around the origin can be expressed as

$$f(x, \lambda) = g_1(x, \lambda)g_2(x, \lambda),$$

where

$$g_1(x, \lambda) = 0, \quad g_2(x, \lambda) = 0,$$

describe the two distinct solution branches that intersect transversely at $(0, 0)$ forming the transcritical bifurcation for certain smooth $g_1(x, \lambda)$ and $g_2(x, \lambda)$ functions.

Proof. By the Morse Lemma, there exists a local diffeomorphism h :

$$h : G_1 \times G_2 \rightarrow I_0 \times \Lambda_0, \quad (0, 0) \in G_1 \times G_2, \\ (0, 0) \in I_0 \times \Lambda_0,$$

such that $h(0, 0) = (0, 0)$, as $(0, 0)$ is a nondegenerate critical point. Furthermore, by the Morse Lemma h satisfies

$$f(h(g_1, g_2)) = g_1 g_2.$$

Now, by taking $h^{-1}(g_1, g_2)$ we can express f as

$$f(x, s) = g_1(x, s)g_2(x, s),$$

such that, as the Hessian matrix of the system has a negative determinant because of the saddle behavior at the origin, with the level curves describing hyperbolas,

$$g_i(0, 0) = 0, \quad \frac{\partial g_i}{\partial x}(0, 0) \neq 0,$$

where g_i defines the behavior of the two branches for the transcritical bifurcation, and by the implicit function theorem applied to each branch g_i , we can consider $g_i(x, \lambda) = g_i(x(\lambda), \lambda)$. ■

APPENDIX B: COMPARISON OF THE TRANSIENT TIMES FOR ISOLATED AND MULTIPLE NEARBY BIFURCATIONS

In this appendix, we compare the transient times close to several bifurcations by inspecting six different cases. To facilitate this comparison, we define the distance to the bifurcation in a general way as $\Delta\delta = |p - p_*|$, where p is the control parameter and p_* denotes the bifurcation value. The transients have been computed choosing initial conditions passing through the delaying regions for each case [64], e.g.,

for SN bifurcations we have chosen initial conditions passing through the bottleneck of the ghosts. By doing so, we aim to compare such times to show that the dynamics close to multiple bifurcations, such as the ones studied for the mushroom bifurcation diagrams, are nonadditive. As we show below, not only are the effects on transients in the mushroom bifurcation nonadditive but they also show a coupling effect of both SN1 and SN4 when approaching the transcritical bifurcation. That is, a resonance effect of these bifurcations produces a synergistic slowing down.

We first show the results for transients close to a single SN bifurcation using the normal form $\dot{x} = a + x^2$ (blue triangles in Fig. 6) and the transients obtained by the sum of times for two independent and distant SN bifurcations (violet rectangles in Fig. 6). These later results will be compared with the multiple nearby bifurcations as a case to look for additivity in the resulting transient times. For this case, transients are measured from $x(0) = -0.1$ to $x = 0.1$. Then, the same results are obtained for the normal form of the transcritical bifurcation, $\dot{x} = ax - x^2$ (open blue circles in Fig. 6). The green data in Fig. 6, which appear overlapped to the results for the transcritical bifurcation, show the times for the transcritical case plus twice the time of a SN bifurcation. In a linear additive scenario, transient mushroom dynamics phenomena should follow such times. Then, we compute these transient times for the normal form Eq. (2) (orange circles in Fig. 6 and the gene circuit system given by Eq. (1) (black circles in Fig. 6). For these two models, we set $s = 0$ and adjusted the opening gap such that $\Delta\delta$ corresponds to the distance $\Delta\delta = |s - s_{\text{SN1}}| = |s_{\text{SN1}}|$. For Eq. (1) we used $r = 0.14$. For the normal form and model of the mushroom bifurcation, transient times were measured for $x(0) = -0.1$ to $x = 0.1$, and from $u(0) = 0.2$ to $u = 0.4$, respectively.

The results show that the longest transients are found for the self-activating gene model Eq. (1) and the normal form (2). These results also show that the slowing down obtained for the multiple bifurcation models departs from an additive effect of individual local bifurcations. For instance, the times obtained with the normal form of the mushroom model and the gene model are much longer than the times obtained by summing up the delays tied to the transcritical bifurcation plus the delays of two independent SN bifurcations. Finally, the times for the multiple bifurcations are about three to four orders of magnitude longer than those for two single and far away SN bifurcations, computed either separately or additively.

APPENDIX C: ANALYTICAL LOCAL APPROXIMATION OF THE MUSHROOM BIFURCATION AND SCALING LAWS

Theorem C1. (Implicit function theorem [65]). Let f be a function of class C^p (with $p > 0$) defined on an open set U of $\mathbb{R}^2$ and taking values in $\mathbb{R}$. Let (x_0, y_0) be a point in U such that $f(x_0, y_0) = 0$ and such that the partial derivative of f with respect to the second variable is nonzero at (x_0, y_0) . There exists a real function φ of class C^p , defined on an open real interval V containing x_0 , and an open neighborhood Ω of (x_0, y_0) in U such that, for all $(x, y) \in \Omega$,

$$[(x, y) \in \Omega \text{ and } f(x, y) = 0] \Leftrightarrow [x \in V \text{ and } y = \varphi(x)].$$

Applying Theorem C1 and, as a specific case of the Malgrange preparation theorem, we can state the following proposition to describe the local factorization of a function f in the neighborhood of a saddle-node bifurcation.

Proposition C1. Suppose $f(x, r)$ is a smooth function with respect to $x \in \mathbb{R}$ in a neighborhood of $x = 0$ such that $f(0, 0) = 0$, $\frac{\partial f}{\partial r}(0, 0) \neq 0$, $\frac{\partial f}{\partial x}(0, 0) = 0$, and $\frac{\partial^2 f}{\partial x^2}(0, 0) \neq 0$. Then, there exists a smooth function $p(r, x)$ which is nonzero in a neighborhood near the origin $N = [-\delta, \delta]^2$, such that f can be locally expressed as

$$f(x, r) = p(x, r)[r - a(x)], \quad (\text{C1})$$

where $a(x)$ satisfies $a(0) = a'(0) = 0$ and $a''(0) \neq 0$.

Proof. Given the condition $f(x, r) = 0$ and the orders of vanishing according to each variable, we can apply the implicit function Theorem to assert $r = r(x)$ and the Mean Value Theorem to proceed:

$$\begin{aligned} f(x, r) &= f(x, r) - f(x, r(x)) \\ &= \int_0^1 \frac{d}{dt} \{f[x, tr + (1-t)r(x)]\} dt \\ &= \int_0^1 \frac{\partial f}{\partial r}[x, tr + (1-t)r(x)] \frac{d}{dt} [tr + (1-t)r(x)] dt \\ &= \int_0^1 \frac{\partial f}{\partial r}[x, tr + (1-t)r(x)] dt [r - r(x)] \\ &= p(x, r)[r - r(x)]. \end{aligned} \quad (\text{C2})$$

This demonstrates the factorization of f . Furthermore, evaluating at the origin and considering that $a(0) = 0$, we get

$$\frac{\partial f}{\partial r}(0, 0) = \frac{\partial p}{\partial r}(0, 0)0 + p(0, 0)1 = p(0, 0).$$

Therefore, at the origin, the derivative of f concerning r equals the value of the function $p(x, r)$ at the same point. Moreover, the system's dynamics in the r direction at the origin are governed by the factor $p(x, r)$, as expected. The function $a(x)$ is a smooth function of degree 2, satisfying $a(0) = a'(0) = 0$ and $a''(0) \neq 0$. ■

In order to analytically study the SN and transcritical (T) scaling laws for Eq. (1), it is sufficient to prove that close to the bifurcation, the behavior of the model can be approximated by the following form, consisting of two saddle-node bifurcations colliding into a transcritical bifurcation.

Proposition C2. Let f be a smooth function with respect to x in a neighborhood of $x = 0$, defined in an open set $U \subseteq \mathbb{R}^3$ and taking values in $\mathbb{R}$, such that

$$\begin{aligned} f(0, 0, 0) &= 0, \quad \frac{\partial f}{\partial x}(0, 0, 0) = 0, \quad \frac{\partial f}{\partial s}(0, 0, 0) = 0, \\ \frac{\partial f}{\partial r}(0, 0, 0) &\neq 0, \quad \frac{\partial^2 f}{\partial x^2}(0, 0, 0) \neq 0, \\ \frac{\partial^2 f}{\partial x^2}(0, 0, 0) \frac{\partial^2 f}{\partial s^2}(0, 0, 0) - \left(\frac{\partial^2 f}{\partial x \partial s}(0, 0, 0) \right)^2 &< 0. \end{aligned}$$

Then, around the origin, the function can be approximated by

$$f(x, s, r) \approx c(x - sh_0)(x - sh_1) + rf_{1,0},$$

for certain nonzero constants $c, h_0, h_1, f_{1,0}$. Furthermore, if $f(x, s, r) = f(x, -s, r)$,

$$f(X, S, R) \approx X^2 - S^2 + R.$$

X, S, R being rescalings of x, s, r .

Proof. By the hypothesis, function f can be written as

$$f(x, s, r) = f_0(x, s) + r f_1(x, s, r),$$

where $f_1(0, 0, 0) \neq 0$. Note that for $r = 0$, $f_0(x, s)$ presents a transcritical bifurcation for $s = 0, x = 0$ and f_0 by Corollary 2 can be expressed as

$$\begin{aligned} f_0(x, s) &= g_1(x, s)g_2(x, s), \\ g_i(0, 0) &= 0, \quad \frac{\partial g_i}{\partial x}(0, 0) \neq 0. \end{aligned}$$

Now, let us apply Proposition C1, such that

$$g_i(x, s) = \tilde{c}_i(x, s)[x + h_{0,i}(s)],$$

for certain smooth functions $\tilde{c}_i(x, s), h_{0,i}(s)$ such that $\tilde{c}(0, 0) \neq 0$. Therefore,

$$f_0(x, s) = c(x, s)[x - h_{0,1}(s)][x - h_{0,2}(s)], \quad c(0, 0) \neq 0.$$

Now, as $h_{0,i}(0) = 0$, we know that we can write it as $h_{0,i}(s) = \tilde{h}_{0,i}(s)$. Therefore

$$f_0(x, s) = c(x, s)[x - \tilde{h}_{0,1}(s)][x - \tilde{h}_{0,2}(s)].$$

As in a neighborhood of the origin functions $c(x, s), \tilde{h}_{0,i}, f_1(x, s)$ are not zero; we can approximate them by the corresponding leading terms such that

$$f(x, s, r) \approx c(x - sh_0)(x - sh_1) + r f_{1,0}.$$

If the model presents symmetry of the branches, $\hat{h}_0 = -\hat{h}_{0,2}$, and properly doing a rescaling such that $x = \alpha X, s = \beta S, r = \gamma R$ we get

$$f(X, S, R) \approx X^2 - S^2 + R.$$

Therefore, for $r \neq 0$ we can study the behavior for the transient times τ close to the SN, describing its scaling law, and for $r \rightarrow 0^+$ we can study the transcritical scaling law as in Sec. II A, resulting in the same asymptotic behavior except for certain extra prefactors depending on rescaling factors $\alpha, \beta, \gamma, \delta$, applying Proposition C2.

For the SN scaling law, let us consider

$$\begin{aligned} \tau &= \frac{1}{\alpha} \int_{-\epsilon}^{\epsilon} \frac{1}{\left(\frac{x}{\alpha}\right)^2 - \left(\frac{s}{\beta}\right)^2 + \left(\frac{r}{\gamma}\right)} dx \\ &= \alpha \int_{-\epsilon}^{\epsilon} \frac{1}{x^2 + \frac{r}{\gamma\beta^2}(\beta^2 - A^2\gamma)} dx, \end{aligned}$$

where in the last equality we have used $s = A\sqrt{r}$. To study the asymptotic behavior of τ , let us consider $(\beta^2 - A^2\gamma) \rightarrow 0$, meaning we are considering its behavior in the vicinity of the SN bifurcations. Therefore

$$\tau(r, \beta, \gamma, \alpha) \rightarrow \frac{\alpha\beta\sqrt{\gamma}\pi}{\sqrt{\beta^2 - A^2\gamma}\sqrt{r}}.$$

For the transcritical scaling law, a similar analysis can be derived, resulting in

$$\begin{aligned} \tau(r, s, \alpha, \beta, \gamma) &= \lim_{\frac{r}{\alpha} \rightarrow 0^+} \frac{\alpha\pi}{\sqrt{\left(\frac{r}{\alpha}\right) - \left(\frac{s}{\beta}\right)^2}} \\ &= \lim_{r \rightarrow 0^+} \frac{\alpha\beta^2\gamma\pi}{(\sqrt{\sqrt{r}\beta - s\sqrt{\gamma}})(\sqrt{\sqrt{r}\beta + s\sqrt{\gamma}})} \\ &= \lim_{s_{sn}^- - s_{sn}^+ \rightarrow 0^+} \frac{\alpha\beta^2\gamma\pi}{(s_{sn}^- - s_{sn}^+)}. \end{aligned}$$

APPENDIX D: BISTABLE SWITCH MODEL

Similar to the mushroom model given in Sec. II, we here introduce a bistable switch model to help decoupling the difference between the effects of a single SN bifurcation and multiple bifurcations.

We use the same simple regulatory model as for the mushroom,

$$\frac{du}{dt} = r + a(s)H(u) - u, \quad (D1)$$

but with a modified external signal $a(s)$:

$$a(s) = \begin{cases} sq - s^2 & \text{if } s < q/2 \\ q^2/4 & \text{otherwise.} \end{cases} \quad (D2)$$

The stochastic models for the bistable switch are the same as for the mushroom bifurcation (see Tables I and II) with a change only in transition rate W_1 , which is dependent on the new external signal function. In the intrinsic noise case, the signal can be rewritten as

$$a(s) = \begin{cases} \frac{1}{\Psi^2}(SQ - S^2) & \text{if } S < Q/2 \\ \frac{Q^2}{4\Psi^2} & \text{otherwise.} \end{cases} \quad (D3)$$

- [1] C. Wissel, A universal law of the characteristic return time near thresholds, *Oecologia* **65**, 101 (1984).
- [2] F. Nazarimehr, S. Jafari, M. Perc, and J. C. Sprott, Critical slowing down indicators, *Europhys. Lett.* **132**, 18001 (2020).
- [3] J. Kramer and J. Ross, Stabilization of unstable states, relaxation, and critical slowing down in a bistable system, *J. Chem. Phys.* **83**, 6234 (1985).

- [4] S. T. Trickey and L. N. Virgin, Bottlenecking phenomenon near a saddle-node remnant in a Duffing oscillator, *Phys. Lett. A* **248**, 185 (1998).
- [5] M. Fitzpatrick, N. M. Sundaesan, A. C. Y. Li, J. Koch, and A. A. Houck, Observation of a dissipative phase transition in a one-dimensional circuit QED lattice, *Phys. Rev. X* **7**, 011016 (2017).

- [6] P. Brookes, G. Tancredi, A. D. Patterson, J. Rahamim, M. Esposito, T. K. Mavrogordatos, P. J. Leek, E. Ginossar, and M. H. Szymanska, Critical slowing down in circuit quantum electrodynamics, *Sci. Adv.* **7**, eabe9492 (2021).
- [7] M. J. Dunlavy and D. Venus, Critical slowing down in the two-dimensional Ising model measured using ferromagnetic ultrathin films, *Phys. Rev. B* **71**, 144406 (2005).
- [8] C.-J. Yang, K. Kliemt, C. Krellner, J. Kroha, M. Fiebig, and S. Pal, Critical slowing down near a magnetic quantum phase transition with fermionic breakdown, *Nat. Phys.* **19**, 1605 (2023).
- [9] S. H. Strogatz and R. M. Westervelt, Predicted power laws for delayed switching of charge density waves, *Phys. Rev. B* **40**, 10501 (1989).
- [10] S. Lade and T. Gross, Early warning signals for critical transitions: A generalized modeling approach, *PLoS Comput. Biol.* **8**, e1002360 (2012).
- [11] V. Dakos and J. Bascompte, Critical slowing down as early warning for the onset of collapse in mutualistic communities, *Proc. Natl. Acad. Sci. USA* **111**, 17546 (2014).
- [12] J. Sardanyés, F. Ivančić, and B. Vidiella, Identifying regime shifts, transients and late warning signals for proactive ecosystem management, *Biol. Conserv.* **290**, 110433 (2024).
- [13] S. H. Strogatz, *Nonlinear Dynamics and Chaos with applications to Physics, Biology, Chemistry, and Engineering* (Westview Press, Cambridge, MA, 2000).
- [14] E. Fontich and J. Sardanyés, General scaling law in the saddle-node bifurcation: a complex phase space study, *J. Phys. A: Math. Theor.* **41**, 015102 (2008).
- [15] R. M. N. Teixeira, D. S. Rando, F. C. Geraldo, R. N. Costa-Filho, J. A. Oliveira, and E. D. Leonel, Convergence towards asymptotic state in 1-D mappings: A scaling investigation, *Phys. Lett. A* **379**, 1246 (2015).
- [16] E. D. Leonel, Defining universality classes for three different local bifurcations, *Commun. Nonlinear Sci. Numer. Simulat.* **39**, 520 (2016).
- [17] B. Vidiella, E. Fontich, S. Valverde, and J. Sardanyés, Habitat loss causes long extinction transients in small trophic chains, *Theor. Ecol.* **14**, 641 (2021).
- [18] J. Sardanyés and R. V. Solé, Ghosts in the origins of life? *Int. J. Bifurcation Chaos* **16**, 2761 (2006).
- [19] J. Gimeno, À. Jorba, and J. Sardanyés, On the effect of time lags on a saddle-node remnant in hyperbolic replicators, *J. Phys. A: Math. Theor.* **51**, 385601 (2018).
- [20] J. Canela, L. Alsedà, N. Fagella, and J. Sardanyés, Dynamical mechanism behind ghosts unveiled in a map complexification, *Chaos, Solitons Fractals* **156**, 111780 (2022).
- [21] À. Calsina, S. Cuadrado, B. Vidiella, and J. Sardanyés, About ghost transients in spatial continuous media, *Chaos, Solitons Fractals* **166**, 112915 (2023).
- [22] J. Sardanyés, R. Martínez, C. Simó, and R. Solé, Abrupt transitions to tumor extinction: A phenotypic quasispecies model, *J. Math. Biol.* **74**, 1589 (2017).
- [23] J. Sardanyés, R. Martínez, and C. Simó, Trans-heteroclinic bifurcation: a novel type of catastrophic shift, *R. Soc. Open Sci.* **5**, 171304 (2018).
- [24] E. Fontich, A. Guillamon, J. T. Lázaro, T. Alarcón, B. Vidiella, and J. Sardanyés, Critical slowing down close to a global bifurcation of a curve of quasi-neutral equilibria, *Commun. Nonlinear Sci. Numer. Simul.* **104**, 106032 (2022).
- [25] J. Sardanyés, C. Raich, and T. Alarcón, Noise-induced stabilisation of saddle-node ghosts, *New J. Phys.* **22**, 093064 (2020).
- [26] J. T. Lázaro, T. Alarcón, C. P. Garay, and J. Sardanyés, Semi-classical theory predicts stochastic ghosts scaling, *Proc. R. Soc. A* **479**, 20220621 (2023).
- [27] W. Shou, S. Ram, and J. M. G. Vilar, Synthetic cooperation in engineered yeast populations, *Proc. Natl. Acad. Sci. USA* **104**, 1877 (2007).
- [28] P. Dai, D. Vorselen, K. S. Korolev, and J. Gore, Generic indicators for loss of resilience before a tipping point leading to population collapse, *Science* **336**, 1175 (2012).
- [29] D. E. Tufcea and P. François, Critical timing without a timer for embryonic development, *Biophys. J.* **109**, 1724 (2015).
- [30] E. Herrera-Delgado, R. Perez-Carrasco, J. Briscoe, and P. Sollich, Memory functions reveal structural properties of gene regulatory networks, *PLoS Comput. Biol.* **14**, e1006003 (2018).
- [31] R. Perez-Carrasco, P. Guerrero, J. Briscoe, and K. M. Page, Intrinsic noise profoundly alters the dynamics and steady state of morphogen-controlled bistable genetic switches, *PLoS Comput. Biol.* **12**, e1005154 (2016).
- [32] K. Exelby, E. Herrera-Delgado, L. Garcia, R. Perez-Carrasco, A. Sagner, V. Metzis, P. Sollich, and J. Briscoe, Precision of tissue patterning is controlled by dynamical properties of gene regulatory networks, *Development* **148**, dev197566 (2021).
- [33] T. Rayon, D. Stamatakis, R. Perez-Carrasco, L. Garcia-Perez, C. Barrington, M. Melchionda, K. Exelby, J. Lázaro, V. L. J. Tybulewicz, E. M. C. Fisher, and J. Briscoe, Species-specific pace of development is associated with differences in protein stability, *Science* **369**, eaba7667 (2020).
- [34] I. Barbier, R. Perez-Carrasco, and Y. Schaeferli, Controlling spatiotemporal pattern formation in a concentration gradient with a synthetic toggle switch, *Mol. Syst. Biol.* **16**, e9361 (2020).
- [35] X. Tan, J. H. Letendre, J. J. Collins, and W. W. Wong, Synthetic biology in the clinic: Engineering vaccines, diagnostics, and therapeutics, *Cell* **184**, 881 (2021).
- [36] T.-C. Tang, B. An, Y. Huang, S. Vasikaran, Y. Wang, X. Jiang, T. K. Lu, and C. Zhong, Materials design by synthetic biology, *Nat. Rev. Mater.* **6**, 332 (2021).
- [37] R. Perez-Carrasco, C. P. Barnes, Y. Schaeferli, M. Isalan, J. Briscoe, and K. M. Page, Combining a toggle switch and a repressilator within the AC-DC circuit generates distinct dynamical behaviors, *Cell Syst.* **6**, 521 (2018).
- [38] D. Sengupta and S. Kar, Deciphering the dynamical origin of mixed population during neural stem cell development, *Biophys. J.* **114**, 992 (2018).
- [39] H. Song, P. Smolen, E. Avron, D. A. Baxter, and J. H. Byrne, Bifurcation and singularity analysis of a molecular network for the induction of long-term memory, *Biophys. J.* **90**, 2309 (2006).
- [40] K. Sriram, M. Rodriguez-Fernandez, and F. J. Doyle III, A detailed modular analysis of heat-shock protein dynamics under acute and chronic stress and its implication in anxiety disorders, *PLoS ONE* **7**, e42958 (2012).
- [41] An isola is an isolated closed curve in a bifurcation diagram that represents a set of steady-state solutions or periodic solutions of a system. It is not connected to the primary branch of solutions but stands alone as a distinct island of solutions.

- [42] I. Otero-Muras, R. Perez-Carrasco, J. R. Banda, and C. P. Barnes, Automated design of gene circuits with optimal mushroom-bifurcation behavior, *iScience* **26**, 106836 (2023).
- [43] D. Dellwo, H. B. Keller, B. J. Matkowsky, and E. L. Reiss, On the birth of isolas, *SIAM J. Appl. Math.* **42**, 956 (1982).
- [44] A. Giri and S. Kar, Incoherent modulation of bi-stable dynamics orchestrates the Mushroom and Isola bifurcations, *J. Theor. Biol.* **530**, 110882 (2021).
- [45] A. Dey and D. Barik, Emergent bistable switches from the incoherent feed-forward signaling of a positive feedback loop, *ACS Synth. Biol.* **10**, 3117 (2021).
- [46] D. Barik and S. Das, Protocol for potential energy-based bifurcation analysis, parameter searching, and phase diagram analysis of noncanonical bistable switches, *STAR Protoc.* **4**, 102665 (2023).
- [47] S. Das and D. Barik, Origin, heterogeneity, and interconversion of noncanonical bistable switches from the positive feedback loops under dual signaling, *iScience* **26**, 106379 (2023).
- [48] J. M. Raser and E. K. O'shea, Noise in gene expression: Origins, consequences, and control, *Science* **309**, 2010 (2005).
- [49] J. M. Pedraza and A. van Oudenaarden, Noise propagation in gene networks, *Science* **307**, 1965 (2005).
- [50] M. Thattai and A. van Oudenaarden, Intrinsic noise in gene regulatory networks, *Proc. Natl. Acad. Sci. USA* **98**, 8614 (2001).
- [51] N. G. van Kampen, *Stochastic Processes in Physics and Chemistry*, 4th ed. (Elsevier BV, Amsterdam, 2007).
- [52] T. B. Kepler and T. C. Elston, Stochasticity in transcriptional regulation: origins, consequences, and mathematical representations, *Biophys. J.* **81**, 3116 (2001).
- [53] Transient times for the normal form of a single saddle-node bifurcation, given by $\dot{x} = s + x^2$, follow $\tau = \pi/\sqrt{s}$ close to bifurcation threshold, being s proportional to the distance from the bifurcation value.
- [54] D. T. Gillespie, Exact stochastic simulation of coupled chemical reactions, *J. Phys. Chem.* **81**, 2340 (1977).
- [55] M. Weber and J. Buceta, Stochastic stabilization of phenotypic states: The genetic bistable switch as a case study, *PLoS ONE* **8**, e73487 (2013).
- [56] C. L. Manser and R. Perez-Carrasco, A mathematical framework for measuring and tuning tempo in developmental gene regulatory networks, *Development* **151**, dev202950 (2024).
- [57] P. Ashwin, S. Wieczorek, R. Vitolo, and P. Cox, Tipping points in open systems: bifurcation, noise-induced and rate-dependent examples in the climate system, *Philos. Trans. R. Soc. A* **370**, 1166 (2012).
- [58] P. D. L. Ritchie, H. Alkhayuon, P. M. Cox, and S. Wieczorek, Rate-induced tipping in natural and human systems, *Earth Syst. Dyn.* **14**, 669 (2023).
- [59] S. Wieczorek, C. Xie, and P. Ashwin, Rate-induced tipping: Thresholds, edge states and connecting orbits, *Nonlinearity* **36**, 3238 (2023).
- [60] P. E. O'Keeffe and S. Wieczorek, Tipping phenomena and points of no return in ecosystems: beyond classical bifurcations, *SIAM J. Appl. Dyn. Syst.* **19**, 2371 (2020).
- [61] P. D. L. Ritchie, J. J. Clarke, P. M. Cox, and C. Huntingford, Overshooting tipping point thresholds in a changing climate, *Nature* **592**, 517 (2021).
- [62] P. D. L. Ritchie, Ö. Karabacak, and J. Sieber, Inverse-square law between time and amplitude for crossing tipping thresholds, *Proc. R. Soc. A* **475**, 20180504 (2019).
- [63] J. Milnor, *Morse Theory* (Princeton University Press, Princeton, NJ, 1963).
- [64] We notice that either the initial condition or the value of the state variable chosen to compute the transient times does not influence the slow passage times whenever the orbit passes through the ghosts or within the slowing down region of the transcritical case. The time an orbit spends until it is captured by the ghost or until it slows down close to the transcritical bifurcation is much faster than the delay experienced.
- [65] R. Roussarie and J. Roux, *Des Équations Différentielles Aux Systèmes Dynamiques* (EDP Sciences, Les Ulis, France, 2012).