

A 60-year international history of Antarctic subglacial lake exploration

Martin J. Siegert

Grantham Institute and Department of Earth Science and Engineering, Imperial College London, London SW7 2AZ, UK

In January 2013, the US WISSARD programme measured and sampled Lake Whillans, a subglacial water body at the edge of West Antarctica, in a clean and environmentally sensitive manner, proving the existence of microbial life beneath this part of the ice sheet. The success of WISSARD represented benchmark in the exploration of Antarctica, made possible by a rich and diverse history of events, discoveries and discussions over the past 60 years; ranging from geophysical measurement of subglacial lakes, to the development of scientific hypotheses concerning these environments and the engineering solutions required to test them. In this article, I provide a personal account of this history, from the published literature and my own involvement in subglacial lake exploration over the last 20 years. I show that our ability to directly measure and sample subglacial water bodies in Antarctica has been made possible by a strong theme of international collaboration, at odds with the media representation of a scientific ‘race’ between nations. I also consider plans for subglacial lake exploration and discuss how such collaboration is likely to be key to success of future research in this field.

Introduction

There has been much written about the history of scientific discovery in Antarctica, revealing how science, exploration and geopolitics have been closely intertwined since the late 19th Century (e.g., Naylor et al., 2008). Scientific advances in Antarctica can be attributed to major periods of collaboration and cooperation, such as during the four International Polar Years (IPYs 1882-3, 1932-3, 1957-8 and 2007-8). The 1957-8 IPY was regarded as a huge success and prompted the further integration of scientific discussion through the establishment of the Scientific Committee on Antarctic Research (SCAR) and, following this, the Antarctic Treaty. SCAR works to coordinate and facilitate international cooperation in Antarctic science and to provide the Antarctic Treaty with scientific evidence, which allows international oversight on matters such as environmental protection and adherence to the Treaty's other rules. Thus, Antarctic science, particularly large programmes that require considerable logistical support and technological development over many years, often develops as a consequence of scientific planning, multinational collaboration and international scrutiny. One example of such research is the exploration of Antarctic subglacial lakes, bodies of water at the bed of the ice sheet, which SCAR has acted to facilitate over the past twenty years, and which culminated in field programmes to measure and sample three individual lakes in 2012/13; Lake Vostok led by a Russian team, Lake Ellsworth led by the British and Lake Whillans led by the US. A recurring theme of the media interest surrounding these programmes was the notion of a scientific ‘race’ between apparently competing nations, each attempting to be the first to discover the secrets that lie within these dark, cold and isolated environments (e.g. The Guardian, February 14th 2012 headline “Antarctic lake race sees scientists dash for life's secrets in subglacial world”). So, which is true? Has Antarctic subglacial lake exploration been a model of noble internationalism in polar science, or have individual nations, and scientists, been in it for themselves? To answer this question, I provide a personal view of the history of subglacial lake exploration, from which one can retrospectively understand whether it has been helped, or not, by an international approach to

scientific discovery and whether any individual nation has had, or can have, the ability to independently 'race' with another.

Subglacial lake identification, measurement and distribution

Shortly after the end of the Second World War, Australian physicist Gordon Robin, through his PhD investigations as part of the Norwegian-British-Swedish expedition to Dronning Maud Land in East Antarctica, perfected the use of seismic sounding to measure ice thickness (Robin, 1958). Seismic waves (e.g. sound) travel well through dense ice, but are attenuated by soft firn and snow at the surface. Consequently, to increase the signal to noise ratio, boreholes (~50 m deep) need to be drilled; one for the charge, one for the receiver(s). The experiment is simple; a small explosion sets off a sound wave, which travels down to the ice base where it is reflected and subsequently recorded by the receiver. The two-way travel time is noted, and converted into distance as the speed of sound in ice is known reasonably well. Thus, a measure of ice thickness is possible using a simple seismic reflection test, adapted for harsh Polar field conditions by Robin. While the process of data acquisition is time-consuming (two boreholes for each data point, meaning that a single datum would need at least a day to record) by aligning measurements along a survey line a profile of ice-sheet thickness, and therefore bed topography, could be derived. In this way, the first 2-dimensional cross-section measurements of the Antarctic ice sheet and its subglacial landscape were obtained.

The mission statement of the 3rd IPY (also known as the International Geophysical Year) was "*to observe geophysical phenomena and to secure data from all parts of the world; to conduct this effort on a coordinated basis by fields, and in space and time, so that results could be collated in a meaningful manner*". This inclusive approach led to several exploratory scientific missions across Antarctica, using the seismic techniques described by Robin a few years earlier. Two overland traverses were most notable. A US expedition crossing West Antarctica, involving a young glaciologist from the University of Wisconsin named Charles Bentley, and a Russian survey from the coast to the centre-point of East Antarctica (the Pole of Relative Inaccessibility), which had among its party Andrei Kapitsa from Moscow State University. The data collected by these surveys transformed our knowledge of the continent, proving it to be a single landmass, showing the ice to be several kilometres thick (at Vostok Station for example it was measured as ~3.7 km) and, in large parts of West Antarctica, revealing the bed to be over a kilometre below sea level.

In the early 1960s UK physicist Stan Evans and Robin, by now Director the Scott Polar Research Institute (SPRI) in Cambridge, began experiments to understand the electrical properties of cold ice and how VHF radio waves could be used to measure ice thickness. VHF radio waves (50-150 MHz) travel well through cold ice (<-10°C) but reflect off boundaries of large dielectric contrast (such as at the bed). Radio-echo sounding (RES), as it was known (essentially ice-penetrating radar), is able to chart ice thickness, therefore, in an analogous way to seismic sounding. The major advantages of RES over seismic sounding was that it did not require the drilling of boreholes and could be deployed on a moving platform to obtain cross-section information during transit. The most significant innovation by Evans and Robin was to consider how RES could be mounted and used effectively on aircraft. In the late 1960s their Cambridge team, supported by funding and logistics from the US Antarctic Research Programme, demonstrated the use of airborne RES with instant and revolutionary success.

Using RES on an aircraft, the rate and quality of data acquisition improved enormously. What took at least a day to get a data point now took less than a second and with equal accuracy (an improvement by 5 orders of magnitude). What took a season to build a transect now took a single

sortie; and where a profile of the ice sheet may have been constructed with a few dozen seismic data-points now it could be put together with many thousands of RES reflections. Early RES trial flights, using a Super Constellation L-1049 aircraft, were targeted at the very centre of the East Antarctic ice sheet, where Russian traversing had covered a decade before. In so doing, Robin and his team proved continental-wide coverage by aircraft mounted with RES was feasible, and that the data were remarkable. So followed one of the world's key decades in Antarctic glaciological and continental discovery.

Systematic profiling of the Antarctic ice sheet took place in four field seasons 1971/72, 1974/75, 1977/78 and 1978/79. Over the decade, further advances in RES equipment were made, primarily through physicist Preben Gudmundsen from the Technical University of Denmark, and navigation was improved (the early flights used 'dead reckoning', which was replaced by an Inertial Navigation System or 'INS'). The aircraft of choice by now was a reliable long-range Hercules C130 transporter, supplied by the US Navy. Thus, a US-UK-Danish collaboration surveyed about 40% of East Antarctica and 80% of West Antarctica, defining the subglacial landscape for the first time and making profound discoveries about the continent and the way in which ice flowed over it (Drewry, 1983). For example, the first subglacial lake was discovered from data collected on one of the first long-range survey flights in 1969. An unusually flat subglacial radio-echo surface beneath the Russian base at Sovetskaya Station in central East Antarctica was received and attributed to a "thick layer of water beneath the ice" (Robin et al., 1970). Shortly afterwards, the first inventory of seventeen subglacial lakes was documented from East Antarctic RES data in 1971/72 (Oswald and Robin, 1973) (Figure 1). Lake Vostok, the gigantic subglacial lake in East Antarctica, was detected by RES in 1974 (Robin et al. 1977), although its true extent was not established at this time. These early discoveries showed that the bed of the Antarctic Ice Sheet to be wet in many places, that water could be stored in 'lakes' and that the distribution of subglacial lakes was widespread about the surveyed regions of the continent.

Although subglacial lakes were undeniably discovered using RES by Robin and his team, the first mention of 'lakes' in Antarctica was made by a Russian pilot (Robinson, 1960) who, as part of an experiment to determine ice-surface landmarks to aid flight orientation, identified "oval depressions with gentle shores" on the ice surface. Although these features were referred to as 'lakes' by pilots who observed them, such as Robinson, there was no connection made between these features and water beneath the surface.

It is interesting to note that despite the published discovery of subglacial lakes in the 1970s, hardly any research was conducted on them during the 1980s. This was almost certainly due to the cessation of long-range surveying of the ice sheet at the expense of targeted smaller-scale observations (Turchetti et al., 2008). This changed, however, in the early 1990s with the introduction of satellite observations of ice-surface altimetry; specially the European Remote Sensing satellite (ERS-1). Ridley et al. (1993) analysed ERS-1 data from East Antarctica, and noticed a remarkably flat surface at and to the North of Vostok Station. This surface, which pilots may have observed in the 1950s, coincides remarkably well with RES evidence for a large subglacial lake established by Robin et al. (1977), and delineates the outline of the lake beneath (the ice surface above large subglacial lakes is flat due to the frictionless contact between ice and water at the ice-sheet base). The combination of satellite and RES data confirmed that this lake, Lake Vostok, was over 240 km long and more than 50 km wide. These data also showed the existence of the lake directly beneath Vostok Station; where Kapitsa and others had collected seismic ice thickness measurements around thirty years previously.

It thus became apparent that the old seismic data may contain a previously unidentified signal of the lake floor given that, unlike radio waves, sound waves travel well in water. To see if this was the case, SCAR convened a meeting at SPRI in 1994 with members of the early exploration programmes including, among others, Robin and Kapitsa. At the meeting, it was agreed that Kapitsa and Robin would work together in analysing the seismic data, drawing in a number of others including the Russian glaciologist Igor Zotikov who had previously worked on determining the temperature distribution of ice masses. This international collaboration led to the discovery that Lake Vostok, hidden beneath ~4 km of ice, had a water-depth of over 500 m (Kapitsa et al., 1996), making it one of the World's top ten largest freshwater lakes. Consideration of the glacial history of East Antarctica suggested that the lake may have been in continual existence for as long as thick ice cover occupied the continent, up to fourteen million years ago. Unlike previous publications on subglacial lakes during the 1970s Kapitsa et al. (1996), which featured on the front cover of *Nature* magazine, was met by world-wide media attention and considerable new scientific interest, particularly from the microbiological community, which instantly regarded the lake to be an extreme yet viable habitat for life, cut off from the rest of the planet for sufficient time to allow novel adaptations to have developed (Ellis-Evans and Wynn-Williams, 1996).

The actual writing of the paper by Kapitsa et al. (1996) was a challenging process, due mainly to the different views between Kapitsa, Robin and Zotikov on who was the first to discover subglacial lakes. Robin argued that his team were the first to publish evidence for subglacial lakes, which was undeniable. However, Kapitsa claimed that his 1961 PhD thesis included the first mention of subsurface water (referred to as 'meltwater lenses') in Antarctica. Zotikov disagreed with Kapitsa, however, because his analysis lacked appreciation of ice-sheet thermal dynamics (Zotikov, 1963) and was, consequently, unsubstantiated. Kapitsa had obtained the seismic data, however, and within those data was the evidence of basal water. Unfortunately, Kapitsa had not spotted the lake-floor reflections, never considered the seismic data to contain information other than ice thickness and so never published on it. He often referred to his failure to tie his ideas on basal water with the seismic data as a scientific regret. At one point in the writing process, this disagreement resulted in the abandonment of the paper until a precise wording of Kapitsa's early ideas was eventually accepted by all three. Thus, in the paper it states, "owing mainly to the high noise level, seismic shooting studies...produced no evidence...of the meltwater lenses beneath the ice sheet that had been suggested by Kapitsa on morphological grounds" (Kapitsa et al., 1996). This seminal paper was saved, therefore, by an important degree of international scientific diplomacy.

The establishment of the geographical scale of Lake Vostok in 1996, and speculative ideas on its potential contents, was supplemented in the same year by a second inventory of 145 Antarctic subglacial lakes (Figure 2), which was collated from a reanalysis of the 1970s RES data collected by Robin's team twenty years earlier (Siegert et al., 1996). The revised inventory demonstrated that scientific interest in subglacial lakes need not necessarily be restricted to Lake Vostok; the physical conditions in that lake, and therefore the environment to support life, being likely similar in many other subglacial water bodies. Nine years later, as more RES data were collected by US, UK, Russian and Italian nations, the inventory of subglacial lakes grew to 145 (Siegert et al., 2005) (Figure 3).

Then, another remarkable finding was made. Analysis of ERS-1 time-series data revealed an area of the central East Antarctic ice sheet to have reduced in elevation by more than 3 meters, far in excess of the instrumental error, over 14 months between 1997 and 1999. During the same period, three areas of the ice sheet more than 200 km away rose up by more than a meter. Given that the regions of uplift coincided with the positions of known subglacial lakes at the mouth of a major subglacial valley known as the Adventure Trench, and that the area of subsidence was located at the head of

this trench, the measurements were interpreted as evidence of an outburst of subglacial lake water, flowing over 200 km along the axis of the trench into a series of other lakes (Wingham et al., 2006). This discovery revealed a previously unappreciated highly active basal hydrology in Antarctica, and demonstrated that, far from being totally isolated, some subglacial lakes could be connected over large distances through a hierarchical hydrological chain (Siegert et al., 2007). Further analysis of satellite interferometry (Gray et al., 2005) and laser altimetry revealed more evidence for basal water flow and subglacial lake outbursts, particularly beneath Whillans Ice Stream (Fricker et al., 2007), leading to an inventory of 130 so-called 'active' subglacial lakes (Smith et al., 2009). Combined with lakes detected from newly acquired RES data, these active lakes pushed the tally within the fourth version of the inventory of Antarctic subglacial lakes to 381 (Wright and Siegert, 2011, 2012) (Figure 4), scattered throughout the continent, confirming wet-based conditions over around half the ice sheet. Following subsequent discoveries, by 2016 the tally of known, discrete subglacial lake locations stood at 402 (Siegert et al., 2016a).

Scientific interest in Lake Vostok and the coordinating role of SCAR

As a consequence of a fragmented thirty-year period of investigation between the mid-1960s and 1990s, and the data collected by Russian, British, US and European scientists, by 1996 subglacial lakes had been discovered in Antarctica and their viability as habitats for life had begun to provoke scientific interest among an interdisciplinary and international community. With Lake Vostok as a focus of attention, and with direct exploration of the lake in the minds of many, a new multidisciplinary field of investigation began to emerge in which microbiologists, geologists and glaciologists discussed the nature of the science they might hope to realise from direct measurement and sampling of the water and, indeed, how such research would be logistically possible and technically feasible.

Two international meetings were held in 1998 to initiate plans for the exploration of Lake Vostok. The first, in Washington DC was led by Robin Bell from Columbia University, and involved mainly US scientists, including participation from NASA, and a few from the UK and elsewhere. This was followed quickly by a Russian-led meeting, hosted at the Arctic and Antarctic Research Institute in St. Petersburg by Valery Lukin, again involving some overseas scientists but without significant US participation. It quickly became apparent to those in SCAR that the developing plans were in danger of being fragmented between Russia and the US at this early stage. For this reason, it convened a third meeting in 1999 in Cambridge (led by Cynan Ellis-Evans from the British Antarctic Survey, BAS) to build international consensus on how exploration of Lake Vostok (and other subglacial lakes, though despite attempts they were rarely given attention) could be achieved, and how international collaboration could support the ambitions. For those few present at all three meetings, three things became clear; (1) that drilling into the lake and acquiring direct measurement and sampling in a clean way was necessary to test whether life existed in the lake (2) that such an experiment had to be conducted cleanly and (3) that no equipment existed to meet these requirements. It was also apparent that the physical conditions of none of the 145 known subglacial lakes (i.e. the size and shape and physiographical setting) was understood in sufficient detail to constrain an exploration plan.

In 1999, to enable the purposeful development of scientific objectives and their delivery, SCAR commissioned an international 'group of specialists' to encourage sharing of data and plans between individual nations, including the acquisition of geophysical data. Within this framework of SCAR support, a series of advances in our knowledge of Lake Vostok and other subglacial lakes was made.

First, Italian geophysicists made the first airborne RES survey of Lake Vostok for twenty years (Tabacco et al., 2002). In the Austral summer of 1999–2000, 12 new RES transects were collected, including a continuous line along the 240 km axis of the lake, which confirmed the conclusion of Kapitsa et al. (1996) of a large, continuous lake, with margins defined by the satellite-derived flat ice surface. This survey was followed in the 2000–2001 season by 20,000 km worth of line geophysical data, collected by the US Support Office for Aerogeophysical Research (SOAR), defining both the lake margin in finer (<10 km) resolution (Studinger et al., 2003a) and its geological setting (Studinger et al., 2003b).

By around the same time Russian geophysicists had collected a series of ground-based measurements, including seismic data, revealing the shape of the lake cavity and confirming the water depth to be more than 1 km at its centre (Masolov et al., 2001). These advances in the physiography of Lake Vostok allowed several teams to utilise numerical modelling, to comprehend the likely physical conditions within the lake, including temperature, salinity, density and the flow of water (Mayer et al., 2003; Wüest and Carmack, 2000; Williams, 2001).

Further SCAR involvement and project development

When in 1996 it was confirmed that Vostok Station was located over the southern edge of Lake Vostok, it became instantly apparent that the base of the Vostok ice core, which had recovered a unique record of Earth's atmospheric composition dating back 420,000 years (Petit et al., 1999), was only ~150 m metres from the roof of the lake. Studies of the basal units of the ice core, expecting to take the climate record further back, revealed virtually no gas content, however. Rather than being formed by ice accumulating at the surface, this gas-poor ice was formed instead by lake water freezing to the ice sheet underside creating over 200 m of 'accreted ice' at the ice sheet base. Thus, without knowing it, the ice core had collected a frozen sample of lake water (Jouzel et al., 1999). The accreted ice was distributed to several teams, as part of a Russian, French, US understanding, and was quickly processed for biogeochemical signatures. In the same volume of *Science* magazine that published evidence of the accretion came two further papers revealing evidence in the accreted ice for microbial life, some of which was culturable (Karl et al., 1999; Priscu et al., 1999). One problem with the accreted ice samples is that their extraction from the ice sheet involved being subjected to the ice-core antifreeze (in this case kerosene). As the samples were not obtained cleanly, and were thus potentially contaminated, the findings of life within the accreted ice were open to challenge. Considerable, sometime acrimonious, debate on the fidelity of the accretion ice microbial content followed, but with the realisation of these disagreements being made redundant if and when direct, clean samples of the water were acquired.

Given that the ice core was positioned so close to the lake ceiling, Russian scientists focused on using the technology to drill further down and into the lake, to recover direct water samples. National plans for subglacial lake exploration were, subsequently, starting to take shape, prompting SCAR to reform its 'group of specialists' in to a formal 'scientific research programme' named Subglacial Antarctic Lake Environments (SALE), which had a remit to coordinate and plan the international exploration of these subglacial systems.

Around 2000, Lake Vostok was seen by many as a special focus for scientific investigation, given its huge size versus other known lakes (as evidenced by the discussion at the international meetings around that time). For some, this focus ignored the facts that the lake was hidden under particularly thick, cold ice, and was located at the very centre of the East Antarctic ice sheet – all of which would

contribute to making its exploration more challenging than for other lakes. This focus also ignored the science that might be achieved through exploration of other subglacial lakes; the question of life in the lake water could potentially be answered by any of the lakes, for example. Given the potentially significant cost of both the technological development and fieldwork, it was appropriate to design an objective set of criteria to comprehend which of the Antarctic subglacial lakes was best suited to exploration in the first instance (Siegert, 2002). Six criteria were drawn as follows: (1) does the lake provide the greatest likelihood for attaining the scientific goals? (2) Can the lake be characterized in a meaningful way (e.g., size, postulated structure)? (3) Is the lake representative of other lakes and settings? (4) Is the geological/glaciological setting understood? (5) Is the lake accessible (what is the closest infrastructure)? (6) Is the program feasible within cost and logistical constraints? Siegert (2002) concluded that one of the relatively small (~10 km long) subglacial lakes, beneath the thinner, warmer ice in West Antarctica would be best suited for the first clean direct exploration of a subglacial lake. The advantage such a lake would have over lakes beneath relative thick cold ice is that it made the prospect of using hot-water drilling, allowing potentially clean access to the lake, relatively feasible. This analysis was quickly followed by the SALE group's evaluation of the programme, and its timeframe, that would be needed for such exploration (Priscu et al. 2003), the discovery of Lake Ellsworth (which had not featured in the 1996 inventory) as a "candidate for subglacial lake exploration" (Siegert et al., 2004) and the updated inventory of 145 lakes (Siegert et al. 2005). As a consequence, by 2005, Lake Vostok was no longer the prime focus of attention for many, though it remained key to Russian scientific plans given the infrastructure already in place at Vostok Station, and because the ice core borehole was so close to the surface of the lake.

In 2006, a fifth international meeting, organised by SALE, was held at Grenoble, France. Its purpose was to shift the discussion from the scientific drivers for subglacial lake exploration (as these had become clear from previous meetings) to the practical plans necessary to deliver an exploration programme. By this time, a UK team had established Lake Ellsworth as a preferred candidate for exploration, having funding in place to conduct a comprehensive geophysical investigation of the lake to determine its dimensions and physiographical setting. Hence, with Russian scientists maintaining interests in Lake Vostok, two separate programmes began to form. In the US, a National Academy of Sciences (NAS) undertook an independent assessment of the exploration of subglacial aquatic environments with the intention of defining a set of standards for responsible exploration of these systems (National Research Council, 2007). SALE evaluated the report and accepted its findings, promoting cleanliness of any direct measurement and sampling experiment to be paramount over other considerations. While it failed to offer a view of which technology was best suited to clean exploration, the NAS report made a clear and thoughtful critique of the various techniques available, including their potential pitfalls on this key issue, from which an individual could reach an objective understanding of the form of experiment needed.

As a consequence, and through discussions held previously by SALE, many concluded that hot-water drilling was the only feasible means by which 'clean' direct access to a subglacial lake environment could be achieved, with the notable exception of members of the Russian programme on Lake Vostok whose focus remained on using the on-site ice-coring technology. The Russian position drew criticism from some (for example the Committee on Environmental Protection, CEP, of the Antarctic Treaty Consultative Committee, ATCM), over continued plans to use the kerosene-filled borehole to access Lake Vostok and, potentially, deploy instruments.

In 2007, following the identification of 'active' lakes within Whillans Ice Stream in West Antarctica, plans began to form surrounding the direct investigation of the one of the lakes, named Lake

Whillans, the ice-sheet grounding zone downstream of the lake and the subglacial ice shelf environment. Hence, by the end of the first decade of the 21st Century, three programmes had been initiated: a trilogy of US programmes on the Whillans ice stream system, the UK programme on Lake Ellsworth and the Russian programme on Lake Vostok.

Plans for exploration

A ground-based geophysical survey of the Lake Ellsworth region, over two seasons in 2007-8 and 2008/9, revealed the lake to be buried beneath 3-3.3 km of ice, around 14 km long, 2-3 km wide and up to 160 m deep; thus confirming it as a deep-water subglacial lake and an ideal candidate for direct measurement sampling (Woodward et al., 2010). The data revealed the likely presence of sediments on the lake floor, and that the lake was contained within a subglacial topographic valley, formed by the action of dynamic glacial erosion when the West Antarctic Ice Sheet was restricted to regional highlands (Ross et al., 2014). The data also identified the best location to attempt direct access, based on maximising scientific value whilst minimising experimental cost. As a consequence of the first field-seasons results, and through an extensive proposal to design, build and deploy bespoke equipment to cleanly drill through 3.3 km of ice, and to measure and sample the lake water and sediment, funding was awarded to undertake the exploration mission in late 2008.

Lake Whillans was first identified through analysis of time-series satellite altimetry; the ice surface was observed to rise and lower twice as a consequence of two cycles of water input and discharge, respectively (Fricker et al. 2011). The boundaries of the surface change were well defined, implying a single lake had received and issued the water, and that the lakes boundaries were robust to the inferred hydrological changes. Radar confirmed the outline of the lake, and measured the ice thickness over it to be ~800 m. Seismic studies over the lake revealed no evidence for its depth, however, implying that it was either likely shallow (of the order of the wavelength of the sound wave in ice, ~4 m), or was completely drained during data acquisition. Indeed there was an understanding that the depth of water would be likely to change in time due to inputs of water from upstream and discharges. As a consequence of the US Obama Stimulus Package of funding for science, following the 2008 financial crash, the trilogy of projects aimed at investigating the Whillans ice stream system were funded, but in a way that merged the projects to form the WISSARD programme, with Lake Whillans being a priority.

Hence, by 2008, funding was in place to support the exploration of Lake Ellsworth and Lake Whillans, as well as continued Russian work at Lake Vostok. As a consequence, the SCAR SALE programme concluded that it had met its original terms of reference and, therefore, the final meeting was held in 2009 to formally close the programme.

To continue international dialogue between the three most prominent nations, and others, an AGU Chapman Conference was convened in Baltimore in March 2010 (Siegert et al., 2011); the sixth international meeting on subglacial lake exploration. Uniquely, the conference gave an opportunity for engineers to discuss the design of hot-water drills, equipment for water measurement and sampling and sediment recovery, and protocols for experimental cleanliness and environmental stewardship.

Following this meeting, SCAR drafted a 'code of conduct' on Antarctic subglacial lakes exploration, ratified at the 2011 ATCM (held in Buenos Aires, Argentina), which explained the scientific basis for cleanliness and the requirement for this to be achieved under in situ measurement and sampling. For the Lake Ellsworth programme, a prerequisite for exploration was a comprehensive

environmental evaluation (CEE) (as recommended by the 2007 NAS report) agreed by the ATCM CEP. The CEE was submitted to the Buenos Aires Meeting, and was ratified the following year, paving the way for the exploration of Lake Ellsworth to take place. The WISSARD programme did not complete a CEE because, in the opinion of US authorities, its exploration could be covered by existing permits, as it was located at the edge of the grounded ice sheet where direct access had occurred several times in the 1990s for glaciological purposes. Nonetheless, both the Lake Ellsworth and Whillans programmes were designed to be fully compliant with the SCAR code of conduct on subglacial lake access. By the end of 2011 subglacial access programmes were ready to commence at Lake Vostok, Lake Ellsworth and Lake Whillans.

Subglacial lake Exploration

Lake Vostok

In February 2012, the Vostok ice core was continued to base of the ice sheet and into the surface water of Lake Vostok. In so doing the Russian programme became the first to access a subglacial lake. If there was ever a 'race' the Russian scientists had certainly won.

As the ice core's antifreeze drilling fluid, kerosene, is lighter than water, at the point of lake penetration lake water rushed into the borehole leading to a hydrological shock in the lower borehole, leading to the development of cracks within the ice core walls and damage to the drill bit and causing ~1.5 m³ of drill fluid to overspill at the surface. On return to the surface, the drill bit was shown to have ice around it, which would have been from the lake water coating the metal and freezing to it (the drill bit retaining its cold temperature as it entered the lake). The lake water within the borehole froze, and was subsequently re-cored the following season to recover a lake sample.

The use of a kerosene-filled ice core to access Lake Vostok was controversial, with several negative comments made about the experiment at the ATCMs in 2010 and 2011; the issue of environmental protection and sample cleanliness being difficult to guarantee with the experiment. Nonetheless, sample of frozen lake water were acquired, on which laboratory investigations could be made.

Lake Ellsworth

A purpose built, clean hot-water drill, and sterile probes and sediment corers were deployed into Antarctica in November 2012 (Siegert et al., 2012). In December, the equipment was primed and drilling commenced. A number of issues prevented the drill from working properly, however. First, a component of the drill's electronic burner control unit failed, as did its replacement. The component was found subsequently to be faulty, but since the system was being run below its operating temperature (the unit was switched on at -17°C; ideally it should have been warmed to around 0°C) this could not be guaranteed as the reason for its failure. Nonetheless, this setback caused considerable delay as a new part (a reformed circuit board with a totally new component) had to be flown in. This delay led to the unplanned use of fuel, as the system had to be kept warm to avoid freezing. Although the new component worked well, pre-recorded data that helped optimise the boiler performance had been lost, meaning the desired water temperature could not be achieved. This led to drilling at a much reduced speed, and further unplanned loss of fuel. The drilling procedure required a subsurface reservoir, which had been developed successfully, to be linked by the main bore hole. Despite over 24 hours of trying, this link was not made, due to the boreholes not being drilled perfectly vertically (due to bending of the drill head as consequence of low melting

rates). Without this connection, drilling could not continue, and despite the boiler finally attaining the required temperatures, drilling was ceased on 24 December, due to the lack of fuel to drill to the lake and to the loss of surface water needed to continue drilling.

Once back in the UK, the Natural Environment Research Council (NERC) led a formal independent failure review board, in which the causes of the failure were confirmed, and recommendations made for their remedy. Details of this review, the issues encountered during fieldwork, and the modifications to the drill needed to achieve deep, clean measurement and sampling of subglacial environments, can be found in Siegert et al. (2014).

Lake Whillans

In January 2013, Lake Whillans became the first subglacial lake to be accessed using a clean hot water drill, and measured and sampled using clean instruments following agreed protocols (Fricker et al., 2011). As a consequence, samples retrieved became the first from a subglacial system in which contamination issues could be assured, adding confidence to the scientific results.

Drilling into the lake was successful, although several technical issues had to be overcome – demonstrating the difficulty inherent in this work. Upon access to the lake, instruments deployed measured the water column at ~1.5 m. Samples of water and sediment were taken, and returned to the surface for both immediate inspection and transfer to laboratories in the US. They showed that water within Lake Whillans contained “metabolically active” microorganisms, and that it was derived primarily from glacial ice melt with a minor component of seawater (Christner, 2014; Michaud et al., 2016) making it unique among known subglacial environments within Antarctica.

That the subglacial environment of Whillans ice stream contains a small level of sea water will not be a surprise to glaciologists as the region has been subject to sea water inundation and marine sedimentation during previous interglacials. The seawater influence is unlikely to be contemporary, as tidal pumping of modern ice-shelf-cavity sea water is limited to a distance of ~10km upstream of the grounding line, whereas Lake Whillans is ~100 km upstream (Michaud et al., 2016). An observed increase in the proportion of sea water with sediment depth suggests that Antarctic groundwater flow may well be important to Whillans ice stream dynamics, as has been demonstrated by numerical ice flow and hydrology modelling (Christoffersen et al., 2014).

Seventh international meeting on subglacial lake exploration

Activity in the months following the fieldwork in 2012-13 varied considerably between respective nations. While US colleagues were inspecting the samples obtained, UK engineers were involved in examining what went wrong with their drill and how best to put it right and Russian scientists were busy considering how to reactivate the ice core to repeat the lake access. At the same time, SCAR organised a global initiative to identify the most important scientific challenges that need to be addressed by 2035 – a 20-year horizon scan.

The ‘scan, led by former SCAR President Chuck Kennicutt from Texas A & M University, with support from the Tinker Foundation, was a community-led activity. It worked by offering an open call for the most important questions to be faced in 20 years’ time, with a second call to refine these questions and gather support for those deemed most important. These questions, numbering several hundred, were then attributed to scientific themes, to provide a loose organisational framework for

discussion. A retreat was then held in New Zealand in March 2014, involving ~75 scientists, nominated by the community, to consider the top 80 of these questions (details can be found in Kennicutt et al., 2014, 2015). Following this exercise the Council of Managers of National Antarctic Programmes (COMNAP) began a matching exercise, to understand the logistics and engineering challenges that answers to each question will require.

In March 2015, the seventh international meeting on subglacial lake exploration was held at the Royal Society's Chicheley Hall, to discuss the first results and lessons learned from the three exploratory missions (Siegert et al., 2016b). It also focused on the future development of research. To this end, attendees completed a questionnaire relating to their scientific ambitions and preferred programme arrangements required to meet them. Attendees were first asked to list which of the horizon scan questions they plan to address. The top four answers were: (1) How does subglacial hydrology affect ice sheet dynamics, and how important is this linkage? (2) How do the characteristics of the ice sheet bed, such as geothermal heat flux and sediment distribution, affect ice flow and ice sheet stability? (3) How will the sedimentary record beneath the ice sheet inform our knowledge of the presence or absence of continental ice? And (4) How do subglacial systems inform models for the development of life on Earth and elsewhere? Collectively, these questions demonstrate the multidisciplinary nature of the research that can be gained from subglacial lake exploration using a combination of geophysical survey, clean subglacial lake access measurement and sampling, down borehole measurement and sediment drilling; all of which have considerable logistical and engineering requirements.

A second set of questions related to the location where research is thought best conducted in terms of scientific deliverables and logistical ease. Although there were numerous responses for subglacial lakes Vostok, Ellsworth and Whillans, the largest number of respondents commented that a variety of settings was required to fully answer the questions, and not restricting only to subglacial lakes. Indeed, clean sampling of sedimentary material away from subglacial lakes was described by several attendees as being an interesting way of answering the top four questions. That the community did not focus on one particular lake, indicates that there is no single agreed 'best suited' lake for exploration at this stage, testifying to a wide variety of unique subglacial lake environments (Siegert, 2016). Only by the exploration of multiple subglacial targets, across the Antarctic continent, can the full diversity of these systems at the ice sheet bed be comprehended.

The third set of questions concerned the technological advances needed for measurement and sampling, lake access and cleanliness and environmental stewardship. Given that numerous lake exploration probes have already been designed, built and tested, the majority of responses focused on equipment not yet configured, such down borehole monitoring systems, long-term *in situ* measurement, and the deep sampling of benthic sediments. On lake access, the consensus was far clearer; that clean, reliable deep-ice hot-water drilling is required. In terms of cleanliness, most respondents commented that procedures for clean subglacial lake access have now been developed using hot-water drilling. Some remarked that procedures and protocols for monitoring cleanliness of boreholes, and devices passed within them, need to be further established.

The final set of questions focused on whether international collaboration is required to undertake subglacial lake exploration in future and, if it is, what the nature of such collaboration should be. While some level of cross-national collaboration was almost unanimously regarded as being desirable, only half the responses thought it essential. Although some favoured the idea of a single major international programme, financially supported by several nations, the majority of respondents spoke to the advantages of retaining a multiple target approach. Instead of a single managed programme, international collaboration should be enhanced through academic and knowledge exchanged between programmes, and through sharing of samples (where possible) to

ensure reproducibility of results. With the emphasis on informal cooperation rather than on managed collaboration, there was an agreement that SCAR can, and should, retain a role in promoting and coordinating subglacial lake exploration research.

Future research

Testing the hypotheses of life in the extreme environments of subglacial lakes, and the climate records held in sediments across their floors, continues to drive future research planning. If we speculate that within 50 years' time, subglacial lake exploration will be common, then between now and then reliable, clean access to deep subglacial environments needs to be established. In doing so, scientific targets in addition to subglacial lakes become possible. For example, we know that there are several deep sedimentary basins in Antarctica, which heavily influence the flow of ice above. Such sediments are likely to be permeable, offering opportunity for water storage (groundwater) (Siegert et al., 2016a), geological records and, potentially, the build-up of methane due to biogeochemical processes (Wadham et al., 2014).

Medium-sized deep-water lakes at the ice-sheet centre, such as Lake Ellsworth, remain well suited to exploration. However, of the >400 known lakes, others are emerging as equally appealing candidates. For example, a large (>100 km long) subglacial lake has been proposed in Princess Elizabeth Land in East Antarctica (Jamieson et al., 2016), which is around 100 km from a logistics hub of the Chinese Antarctic programme. If confirmed, it will be one of four subglacial lakes that are controlled by the tectonic setting of the region, the others being Lakes Vostok, Sovetskya (the very first lake to be discovered), and 90E (Bell et al., 2006), which all occupy topographic troughs with long-axes parallel to one another and roughly orthogonal to ice flow. Exploration of any of these deep, likely ancient, systems may yield substantial scientific advances. Exploring the easiest first is sensible, which places the new discovery as a feasible and realistic target for exploration in the coming years.

Summary and discussion

A time-line of selected key events within the history of subglacial lake exploration is provided in Figure 5. Sixty years ago, glaciologists were perfecting the field measurement of deep subglacial environments through pioneering developments in seismic and radar sounding. This early work led directly (i.e., provided data as well as methods) to the discovery of Lake Vostok as a huge deep-water body, buried beneath 4 km of ice for potentially millions of years. As a consequence, it was hypothesized to represent a unique and ancient environment for microbial life and a recorder of climate change. As these hypotheses are testable with direct measurement and sampling, the exploration of subglacial lakes became a serious proposition from the mid-1990s.

International discussion on how and where to explore in Antarctica was supported by SCAR, which convened a group of specialists, a scientific research programme and several international meetings. It also oversaw the development of protocols to protect these pristine from unnecessary contamination and disturbance. Slowly, three programmes began to take shape; a Russian plan to use the ice core facility at Vostok Station to break through the ice, a US programme to sample the hydrologically active bed of Whillans ice stream, including Lake Whillans, and a UK-led project to investigate Lake Ellsworth.

As it is easy to see the similarities between the projects – they are subglacial lakes – the notion of a ‘race’ between nations to explore these environments was reported regularly in the international

media. However, while research is an inherently competitive business, participants of the three programmes did not view the competition as a race, in large part because of the distinctions between the three environments: Lake Vostok being huge and in East Antarctica, Lake Ellsworth being medium-sized in West Antarctica, Lake Whillans being shallow and potentially ephemeral. The fact that there was regular correspondence between the programmes, via SCAR and through seven international symposia, testifies to the collegiately and openness across the programmes, in line with the long-held spirit of Antarctic research.

In January 2012, Lake Vostok was penetrated by the Russian Ice Core, allowing frozen lake water to be recovered by subsequent re-coring. In December 2012, the UK-led mission to Lake Ellsworth was halted due to technical issues with the deep-ice drill. In January, 2013, the US programme to Lake Whillans successfully recovered samples, demonstrating the ice base here to contain viable micro-organisms. While these twelve months saw considerable advances in our knowledge of subglacial lake environments, and our ability to explore them, the main drivers for the exploration, namely the two hypotheses on life in ancient deep-water subglacial environments and on climate records, remain strictly untested. When one considers this, and the fact that there are over 400 known subglacial lakes, one can see why subglacial lakes research remains in its infancy as a topic.

Acknowledgements

I wish to thank the organisers and convenors of the XII International Symposium on Antarctic Earth Sciences in Goa for the opportunity to present material within this chapter as a keynote lecture, and to the UK Natural Environment Research Council for funding the Lake Ellsworth exploration programme (NE/G00465X/1, 2 and 3).

References

- Bell, R.E., Studinger, M., Fahnestock, M.A. & Shuman, C.A. 2006. Tectonically controlled subglacial lakes on the flanks of the Gamburtsev Subglacial Mountains, East Antarctica, *Geophys. Res. Lett.*, v. 33, L02504.
- Christner, B.C., Priscu, J.C., Achberger, A.M., Barbante, C., Carter, S.P., Christianson, K., Michaud, A.B., Mikucki, J.A., Mitchell, A.C., Skidmore, M.L., Vick-Majors, T.J., & the WISSARD Science Team. A microbial ecosystem beneath the West Antarctic ice sheet. *Nature*, 512, 310–313, doi:10.1038/nature13667. (2014).
- Christoffersen P, Bougamont M, Carter SP, Fricker HA and Tulaczyk S (2014) Significant groundwater contribution to Antarctic ice streams hydrologic budget. *Geophys. Res. Lett.*, **41**, doi:10.1002/2014GL059250
- Drewry, D.J., 1983. *Antarctica: glaciological and geophysical folio*. Scott Polar Research Institute, University of Cambridge.
- Ellis-Evans, J.C. & Wynn-Williams, D. 1996. Antarctica: A great lake under the ice. *Nature*, v. 381, p. 644-646.
- Fricker, HA, Powell R, Priscu J, Tulaczyk S, Anandakrishnan S, Christner B, Fisher AT, Holland D, Horgan H, Jacobel R, Mikucki J, Mitchell A, Scherer R, Severinghaus J. (2011). Siple Coast subglacial aquatic environments; the Whillans ice stream subglacial access research drilling project. In Siegert, M.J., Kennicutt, M.C. and Bindshadler, R., eds. *Subglacial Antarctic Aquatic Environments*, American Geophysical Union Geophysical Monograph. 192:199-219.
- Fricker HA, Siegfried MR, Carter SP, Scambos TA. 2016 A decade of progress in observing and modelling Antarctic subglacial water systems. *Phil. Trans. R. Soc. A* 374: 20140294. <http://dx.doi.org/10.1098/rsta.2014.0294>
- Fricker, H.A., T. Scambos, R. Bindshadler and L. Padman, 2007, An active subglacial water system in West Antarctica mapped from space. *Science*, **315**, 1544-1548.
- Gray, L., I. Joughin, S. Tulaczyk, V.B. Spikes, R. Bindshadler and K. Jezek, Evidence for subglacial water transport in the West Antarctic Ice Sheet through three-dimensional satellite radar interferometry. *Geophysical Research Letters*, **32** (L03501), doi:10.1029/2004GL021387.
- Jamieson, S.S.R., Ross, N., Greenbaum, J.S., Young, D.A., Aitken, A.R., Roberts, J.L., Blankenship, D.D. & Siegert, M.J. 2016. A large subglacial lake and canyon system in Princess Elizabeth Land, East Antarctica. *Geology*, v. 44, p. 87–90.
- Jouzel, J., J. R. Petit, R. Souchez, N. I. Barkov, V. Y. Lipenkov, D. Raynaud, M. Stievenard, N. I. Vassiliev, V. Verbeke, and F. Vimeux (1999), More than 200 meters of lake ice above subglacial Lake Vostok, Antarctica, *Science*, 286, 2138–2141.
- Kapitsa, A., Ridley, J.K., Robin, G. de Q., Siegert, M.J., and Zotikov, I., 1996. Large deep freshwater lake beneath the ice of central East Antarctica. *Nature*, 381: 684-686.
- Karl, D. M., D. F. Bird, K. Bjorkman, T. Houlihan, R. Shackelford, and L. Tupas (1999), Microorganisms in the accreted ice of Lake Vostok, Antarctica, *Science*, 286, 2144–2147.

Masolov, V. N., V. V. Lukin, A. N. Sheremetyev, and S. V. Popov (2001), Geophysical investigations of the subglacial Lake Vostok in Eastern Antarctica (in Russian), *Dokl. Akad. Nauk*, 379 (5), 680–685. (*Dokl. Earth Sci.*, 379A(6), 734–738.)

Mayer, C., K. Grosfeld, and M. J. Siegert (2003), Salinity impact on water flow and lake ice in Lake Vostok, Antarctica, *Geophys. Res. Lett.*, 30(14), 1767, doi:10.1029/2003GL017380.

Michaud, A.B., Skidmore, M.L., Mitchell, A.C., Vick-Majors, T.J., Barbante, C., Turetta, C., van Gelder, W. & Priscu, J.C. 2016. Solute sources and geochemical processes in Subglacial Lake Whillans, West Antarctica. *Geology*, 44, 347-350. doi:10.1130/G37639.1

National Research Council, 2007. Exploration of Antarctic Subglacial Aquatic Environments: Environmental and Scientific Stewardship, US National Academy of Sciences, Washington DC, 152 p.

Naylor, S., Dean, K. & Siegert, M.J. The IGY and the ice sheet: surveying Antarctica. *Journal of Historical Geography*, 34, 574-595 (2008).

Needle, A. 2000. Science, Cold War and the American State, Smithsonian Institution, Washington DC.

Oswald, G.K.A, and Robin, G. de Q., 1973. Lakes beneath the Antarctic Ice Sheet. *Nature*, **245**: 251-254.

Petit, J. R., et al. (1999), Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429–436.

Priscu J.C., Bell R.E., Bulat S.A., Ellis-Evans J.C., Kennicutt M.C. II, et al. 2003. An international plan for Antarctic subglacial lake exploration. *Polar Geogr.* 27(1):69--83

Priscu, J. C., et al. (1999), Geomicrobiology of subglacial ice above Lake Vostok, Antarctica, *Science*, 286, 2141–2144.

Priscu, J.C., Bell, R.E., Bulat, S.A., Ellis-Evans, J.C., Kennicutt, M.C. II., Lukin, V.V., Petit, J.R., Powell, R.D., Siegert, M.J. and Tabacco, I.E. An international plan for Antarctic subglacial lake exploration. *Polar Geography*, 27(1), 69-83. (2003).

Ridley, J. K., W. Cudlip, and S. W. Laxon (1993), Identification of subglacial lakes using ERS-1 radar altimeter, *J. Glaciol.*, 39, 625–634.

Robin G de Q, Drewry DJ, Meldrum DT. 1977. International studies of ice sheet and bedrock. *Philos. Trans. R. Soc. London* 279:185--96

Robin, G. de Q., 1958. Seismic shooting and related investigations. *Glaciology III*, Published by the Norsk Polarinstitut, Oslo 1958.

Robin, G. de Q., Swithinbank, C.W.M., and Smith, B.M.E., 1970. Radio echo exploration of the Antarctic ice sheet. *International symposium on Antarctic glaciological exploration (ISAGE), Hanover, New Hampshire, USA, 3-7 September, 1968.*

Robinson, R.V., 1960. Experiment in visual orientation during flights in the Antarctic. *International Bulletin of the Soviet Antarctic Expeditions* 18, 28–29.

Ross, N., Jordan, T.A., Bingham, R.G., Corr, H.F.J., Ferraccioli, F., Le Brocq, A., Rippin, D.M., Wright, A.P., Siegert, M.J. The Ellsworth Subglacial Highlands: inception and retreat of the West Antarctic Ice Sheet. *Geological Society of America Bulletin*, 126, 3-15 doi: 10.1130/B30794.1 (2014).

- Siegert, M.J., Makinson, K., Blake, D., Mowlem, M. and Ross, N. An assessment of deep-hot-water drilling as a means to undertake direct measurement and sampling of Antarctic subglacial lakes: experience and lessons learned from the Lake Ellsworth field season 2012-13. *Annals of Glaciology*, 55, 59-73 doi: 10.3189/2014AoG65A008 (2014).
- Siegert, M.J., Dowdeswell, J.A., Gorman, M.R. & McIntyre, N.F. An inventory of Antarctic sub-glacial lakes. *Antarctic Science*, 8, 281-286. (1996).
- Siegert M.J., Hindmarsh, R., Corr H., Smith, A., Woodward, J., King, E., Payne, A.J., Joughin, I. Subglacial Lake Ellsworth: a candidate for *in situ* exploration in West Antarctica. *Geophysical Research Letters*, 31 (23), L23403, 10.1029/2004GL021477. (2004).
- Siegert, M.J., Carter, S., Tabacco, I., Popov, S. and Blankenship, D. A revised inventory of Antarctic subglacial lakes. *Antarctic Science*, 17 (3), 453-460. (2005).
- Siegert, M.J., Le Brocq, A. And Payne A. Hydrological connections between Antarctic subglacial lakes and the flow of water beneath the East Antarctic Ice Sheet. In, *Glacial Sedimentary Processes and Products* (Hambrey, M.J., Christoffersen, P., Glasser, N.F. and Hubbard, B.P). Special Publication #39, International Association of Sedimentologists, 3-10. (2007).
- Siegert, M.J., Kennicutt, M, Bindschadler, R. (eds.). Antarctic Subglacial Aquatic Environments. AGU Geophysical Monograph 192, American Geophysical Union, Washington DC, USA, 246pp. (2011).
- Siegert, M.J., Clarke, R.J., Mowlem, M., Ross, N., Hill, C.S., Tait, A., Hodgson, D., Parnell, J., Tranter, M., Pearce, D., Bentley, M., Cockell, C, Tsaloglou, M.N., Smith, A., Woodward, J., Brito, M.P. and Waugh, E. Clean access, measurement and sampling of Ellsworth Subglacial Lake: a method to explore deep Antarctic subglacial lake environments. *Reviews of Geophysics*, 50, RG1003, doi:10.1029/2011RG000361 (2012).
- Siegert, M.J., Priscu, J. Alekhina, I., Wadham, J. and Lyons, B. Subglacial Antarctic Lake Exploration: first results and future plans. *Transactions of the Royal Society of London*, A. 374, issue 2059 (2016).
- Siegert M.J., Hindmarsh, R., Corr H., Smith, A., Woodward, J., King, E., Payne, A.J., Joughin, I. Subglacial Lake Ellsworth: a candidate for *in situ* exploration in West Antarctica. *Geophysical Research Letters*, 31 (23), L23403, 10.1029/2004GL021477. (2004).
- Siegert, M.J. A wide variety of unique environments beneath the Antarctic ice sheet. *Geology*, 44, 399–400, doi:10.1130/focus052016.1 (2016). <http://hdl.handle.net/10044/1/31391>
- Siegert, M.J. Which are the most suitable Antarctic subglacial lakes for exploration? *Polar Geography*, 26, 134-146 (2002).
- Siegert, M.J., Ross, N. & Le Brocq, A. 2016a. Recent advances in understanding Antarctic subglacial lakes and hydrology. *Phil. Trans. Roy. Soc. Lon A*, v. 374: 20140306.
- Smith, B.E., Fricker, H.A., Joughin, I.R. & Tulaczyk, S. 2009. An inventory of active subglacial lakes in Antarctica detected by ICESat (2003-2008), *J. Glaciol.*, v. 55, p. 573-595.
- Studinger, M., Karner, G.D.; Bell, R.E., Levin, V., Raymond, C.A., Tikku, A.A. (2003b), Geophysical models for the tectonic framework of the Lake Vostok region, East Antarctica, *Earth Planet. Sci. Lett.*, 216(4), 663–677.

- Studinger, M., Bell, R.E., Karner, G.D., Tikku, A.A., Holt, J.W., Morse, D.L., Richter, T.G., Kempf, S.D., Peters, M.E., Blankenship, D.D., Sweeney, R.E., Rystrom, V.L. (2003a), Ice cover, landscape setting, and geological framework of Lake Vostok, East Antarctica, *Earth Planet. Sci. Lett.*, 205(3–4), 195–210.
- Tabacco IE, Bianchi C, Zirizzotti A, Zuccheretti E, Forieri A, et al. 2002. Airborne radar survey above Vostok region, east Antarctica: ice thickness and Lake Vostok geometry. *J. Glaciol.* 48:62–69
- Turchetti, S., Dean, K., Naylor, S. & Siegert, M. Accidents and Opportunities: A History of the Radio Echo Sounding (RES) of Antarctica, 1958-1979. *British Journal of the History of Science*, 41, 417-444
- Wadham, J. L., S. Arndt, S. Tulaczyk, M. Stibal, M. Tranter, J. Telling, G.P. Lis, E. Lawson, A. Ridgwell, A. Dubnick, M.J. Sharp, A.M. Anesio & C.E.H. Butler. 2014. Potential methane reservoirs beneath Antarctica. *Nature*, v. 488, p. 633–637.
- Williams, M.J.M. (2001), Application of a three-dimensional numerical model to Lake Vostok: An Antarctic subglacial lake, *Geophys. Res. Lett.*, 28, 531–534.
- Winberry, J.P., Anandakrishnan, S., Wiens, D.A., Alley, R.B., & Christianson, K. 2011. Dynamics of stick–slip motion, Whillans Ice Stream, Antarctica. *Earth Plan. Sci. Lett.*, v. 305, p. 283–289.
- Wingham, D.J., Siegert, M.J., Shepherd, A.P. & Muir, A.S. Rapid discharge connects Antarctic subglacial lakes. *Nature*, 440, 1033-1036 (2006).
- Woodward J., Smith, A., Ross, N., Thoma, M., Grosfeld, C., Corr, H., King, E., King, M., Tranter, M., Siegert, M.J. Location for direct access to subglacial Lake Ellsworth: An assessment of geophysical data and modelling. *Geophysical Research Letters*, 37, L11501, doi:10.1029/2010GL042884. (2010).
- Wright, A., Siegert, M.J. The identification and physiographical setting of Antarctic subglacial lakes: an update based on recent geophysical data. In, *Subglacial Antarctic Aquatic Environments* (M. Siegert, C. Kennicutt, B. Bindschadler, eds.). AGU Geophysical Monograph 192. Washington DC. 9-26. (2011).
- Wright, A.P. and Siegert, M.J. A fourth inventory of Antarctic subglacial lakes. *Antarctic Science*, 24, 659–664. doi:10.1017/S095410201200048X (2012).
- Wüest, A., and E. Carmack (2000), A priori estimates of mixing and circulation in the hard-to-reach water body of Lake Vostok, *Ocean Modell.*, 2, 29–43.
- Zotikov, I. A. (1963), Bottom melting in the central zone of the ice shield of the Antarctic continent and its influence upon the present balance of the ice mass, *Bull. Int. Assoc. Sci. Hydrol.*, 8(1), 36.

Figure Captions

Figure 1. The inventory of **17** Antarctic subglacial lakes, 1973. Taken from Oswald and Robin (1973).

Figure 2. The inventory of **77** Antarctic subglacial lakes, 1996. Taken from Siegert et al. (1996).

Figure 3. The inventory of **145** Antarctic subglacial lakes, 2005. Yellow lakes are those identified from SPRI data, green lakes are from the Italian dataset, pink lakes were located in US RES data and red lakes have been identified from Russian data. Insets illustrate the coverage of RES data used in the inventory (note that the US and Italian surveys comprise multiple RES transects within boxed areas) and enlargements of two regions of subglacial lakes around Dome C. Abbreviations to place names are as follows: AB (Astrolabe Subglacial Basin); AL (Aurora Lake); CL (Concordia Lake); DA (Dome A); DC (Dome C); DF (Dome F); DML (Dronning Maud Land); EM (Ellsworth Mountains); GVL (George V Land); HD (Hercules Dome); LV (Lake Vostok); MRL (Mac Robertson Land); OL (Oates Land); RB (Ridge B); SP (South Pole); TAD (Talos Dome); TID (Titan Ice Dome); VL (Vincennes Lake); WM (Whitmore Mountains). Taken from Siegert et al. (2005).

Figure 4. The inventory of **381** Antarctic subglacial lakes, 2012. Colours/shapes indicate the type of investigations undertaken at each site: Black/triangle = RES, yellow = seismic sounding, green = gravitational field mapping, red/circle = surface height change measurement, square = shape identified from ice surface feature. Lake Vostok is shown in outline. Taken from Wright and Siegert (2012).

Figure 5. A timeline of subglacial lake discoveries showing key events between 1956 and 2016. Horizontal bars show the total number of known subglacial lakes in published inventories. Adapted from Wright and Siegert (2012).